

Full Length Research Paper

## Biopreservation of tomato paste and sauce with *Leuconostoc* spp. metabolites

Djadouni Fatima, Kihal Mebrouk and Heddadji Miloud

Laboratory of Applied Microbiology, Department of Biology, Faculty of Sciences, Es-Senia University, Oran, Algeria.

Received 14 March 2015; Accepted 5 June 2015

The aim of this study was to evaluate 40 lactic acid bacteria (LAB) and 20 *Bacillus* strains isolated from the fermented tomato (*Solanum lycopersicum*) for their capacity to produce antimicrobial activities against several bacteria and fungi. The strain designed LBc03 has been selected for advanced studies. The supernatant culture of this strain inhibits the growth of *Escherichia coli*, *Staphylococcus aureus* and *Aspergillus* sp. Based on the cultural, morphological, physiological and biochemical characteristics, LBc03 was identified as *Leuconostoc* spp. Its antimicrobial compound was determined as a proteinaceous substance, but it is possible that the bacteriocin may also be bound to other molecules like a lipid or a carbohydrate moiety. Metabolite extracts from selected LAB were more effective in preserving tomato paste and sauce stored at 4°C against spoilage bacteria like *E. coli* and the application of bio-preservative should be encouraged in food processing industries.

**Key words:** Biopreservation, tomato (*Solanum lycopersicum*), bacteriocins, lactic acid bacteria.

### INTRODUCTION

Microbial spoilage of fruits and vegetable is known as rot, which manifests as loss of texture (soft rot), changes in color (black or grey) and often off odor (Trias et al., 2008). Also, the high water content in tomatoes makes it very susceptible to spoilage bacteria and fungi during storage, harvesting and transportation (Spadaro and Gullino, 2004). Fresh food like fruits and vegetables, are normal part of the human diet and are consumed in large quantities in most countries. These products are rich in carbohydrates and poor in proteins with pH value from slightly acidic to 7.0 and provide a suitable niche to several bacteria, yeasts and moulds (Wiessinger et al., 2000; Trias et al., 2008; Ogunbanwo et al., 2014).

Tomato (*Solanum lycopersicum*) is one of the highly nutritious food ingredient used in the preparation of food all over the world (Ogunniyi and Oladejo, 2011; Ogunbanwo et al., 2014). Its utilization as an ingredient in vegetable salads, other dishes and its processing into different products like puree, ketchups and juice is well documented. Nutritionally, it contains a large amount of water, niacin, calcium and vitamins especially A, C, E which are important in the metabolic activities of man and protects the body against diseases (Taylor, 1987). Lycopene (acarotene) an essential component of tomato contributes in the prevention of cardiovascular disease and cancer of the prostate (Clinton, 1998; Bernard et al., 1999).

\*Corresponding author. E-mail: houarisalwa@yahoo.fr.

**Table 1.** S01, S02, S03, and S04 tomatoes *Solanum lycopersicum* samples.

Samples	Origin	Tomato Variety	Characteristics	Additives added	Temperature of storage (°C)	Storage time
S01 <sup>C1</sup>	Mascara, Algeria	<i>Solanum lycopersicum</i>	Fresh-cut tomato	Absence	25	24 h
S02 <sup>C2</sup>	Mascara, Algeria	<i>Solanum lycopersicum</i>	Tomato paste was boiled for 90°C for 02 h	Absence	25	24 h
S03	Mascara, Algeria	<i>Solanum lycopersicum</i>	Fresh-cut tomato	Absence	-20	06 months
S04	Mascara, Algeria	<i>Solanum lycopersicum</i>	Tomato paste was boiled for 90°C for 02 h	05% NaCl and oil	+4	06 months

C1 and C2 are control samples.

The characteristic flavor of tomato is produced by the complex interaction of the volatiles and non-volatile components (Petro-Turza, 1987; Buttery, 1993). The nutritional value of tomato products is a topic attracting much attention, particularly regarding the effects resulting from food processing and storage treatments (Capanoglu et al., 2010). Among the common post-harvest fungal pathogens of tomatoes are *Penicillium expansum*, *Monilinia laxa* and *Rhizopus stolonifer* (Ogawa et al., 1995; Pla et al., 2005). Many LAB strains are able to produce protein compounds with efficient antimicrobial effect, which are known as bacteriocins (Davidson and Harrison, 2002). In recent times, the understanding of the preservation mechanisms of LAB is being exploited for industrial production of foods (Trias et al., 2008) because of their natural acceptance as Generally Recognized as Safe (GRAS) for human consumption and exhibit antimicrobial property (Aguirre and Collins, 1993). There is a complementary effect by the production of acid and antimicrobial compounds that increases inhibition of both pathogen and spoilage bacteria (Edwards et al., 1983; Artés et al., 1999). Although many efforts have been made to develop bio-protective lactic acid bacteria strains, the application of these strains in fresh fruits and vegetables have not been developed yet (Toivonen and DeEll, 2002).

This work is designed to investigate the effectiveness of lactic acid bacteria and *Bacillus* metabolites in preserving tomato paste and sauce against *S. aureus*, *E. coli*, *Clostridium* sp., *Aspergillus* sp., and *Penicillium* sp.

## MATERIALS AND METHODS

### Samples collection

Fresh tomato samples *Solanum lycopersicum* (Table 1) were purchased from markets in Mascara, Algeria. The samples were collected in separate sterile polythene bags and immediately transported to the laboratory for analysis.

### Determination of the physical-chemical characteristics of the tomato samples (S01, S02, S03, and S04)

#### Moisture content

The moisture content of the samples (S01, S02, S03, and S04) was determined before storage time by weighing into moisture cans then weighed and then placed in an oven at 80°C for 24 h to dry to a constant weight. Then, brought out and allowed to cool in desiccators and then reweighed (A.O.A.C., 1995).

$$\text{Thus, moisture content} = \frac{(\text{initial weight} - \text{final weight})}{\text{initial weight}} \times 100$$

#### Dry matter content

Five grams of each sample were obtained and placed into pre-weighed crucibles and dried in at 100°C for 12 h. The dried samples were weighed after cooling in the desiccators (A.O.A.C., 1984).

#### Determination of lactic acid produced by lab isolates

This was achieved based on the methods described (Ogunbanwo et al., 2008; Bamidele et al., 2011). For these measurements lactic acid was determined by transferring 25 ml of the tomato juice (S01, S02, S03, and S04) into conical flasks and 3 drops of phenolphthalein were added as indicator. From a burette, 0.1 M NaOH was slowly added to the samples until a pink color appeared. Each ml of 0.1 M NaOH is equivalent to 90.08 mg of lactic acid.

#### Determination of the pH

pH was determined after and before fermentation of the tomato samples (S01, S02, S03, and S04) by the used of pH meter (Inolab MLM) (Lehninger, 1981).

#### Detection of pathogens, contaminants, LAB, and bacillus

The main objective of this experiment was to detect microbial contaminants in tomato samples (S01, S02, S03, and S04) in addition to other related hygiene tests. Total aerobic bacterial count,

total coliform count, Total anaerobic bacteria count, *Salmonella*, *S. aureus*, *Clostridium*, lactic acid bacteria, and *Bacillus* were detected (Delarras et al., 2006).

### Isolation and selection of LAB strains

LAB strains were isolated from the tomato samples (S01, S02, S03, and S04). The samples were plated directly on MRS as detailed by De Man, Rogosa and Sharpe (Merck, Germany) and M17 agar (Merck, Germany) at 30°C for 2-3 days (pH6.5) under aerobiosis and anaerobiosis conditions. They were routinely propagated and stored at - 20°C supplemented with glycerol (20%, v/v, final concentration). Working cultures were sub-cultured twice (1 inoculum, 24 h, 30°C) prior to use (Djadouni and Kihel, 2012; Ogunbanwo et al., 2014).

### Identification of the LAB isolates

The pure isolate selected as a potential bacteriocin - producer was identified on the basis of its cultural, morphological, physiological and biochemical characteristics. The selected LAB isolates were characterized by Gram stain, absence of spores and catalase test. Gram+, catalase and spores negative strains were maintained frozen until needed for the antimicrobial activity testing. Confirmation of the identification was based on the use of Bergey's manual of systems bacteriology (Sneath et al., 1986).

### Isolation and Identification of bacillus strains

Samples of tomato were weighed as 1 g portions and thoroughly homogenized in sterile distilled water; serial dilutions were plated on LB agar (Luria-Bertani media, Merck, Germany) plates were incubated at 30°C for 2-3 days (Kalil et al., 2009). The pure isolate was examined macroscopically and microscopically and identified with reference to Holt et al., (1994). Isolates were identified by colonial appearance, gram positive, and presence of spores, the presence or absence of  $\beta$ -haemolysis, lecithinase activity, motility, penicillin susceptibility and biochemistry (Abriouel et al., 2010).

### The indicator strains

The indicator strains (*S. aureus*, *E. coli*, *Clostridium* sp., *Aspergillus* sp., and *Penicillium* sp.) used in this work was provided by the Laboratory of Bacteriology, Microbiology Department at the Faculty of Sciences, Es-Senia, Oran University, Algeria. For the antimicrobial assay, the pathogenic cultures were grown in the nutrient agar media (NA) at pH, 7.4 (Ogunbanwo et al., 2014).

For antifungal activities determination, *Aspergillus* sp. and *Penicillium* sp. were grown in potato dextrose agar (PDA, Merck, Germany) for 7 days at 30°C. Spores were collected in sterile distilled water and then concentrated to  $10^4$  spores  $ml^{-1}$  (Smaoui et al., 2010).

### Preparation of cell-free filtrate

MRS broth (1000  $\mu$ l) were inoculated separately with isolates (LAB or *Bacillus*) previously characterized and incubated at 30°C for 72 h. After incubation, a cell free supernatant was obtained by centrifuging (Spectrafuge 24D, Labnet, USA) the bacterial culture at 10.000 rpm for 45 min, followed by filtration of the supernatant through 0.2  $\mu$ m pore size filter paper thus obtaining cell free filtrate (Khalil et al., 2009; Djadouni and Kihel, 2013).

### Antagonistic activity of LAB and bacillus metabolites against spoilage microorganisms

Sixty isolates (40 LAB and 20 *Bacillus*) were grown in MRS broth for 72 h at 30°C and the broth cultures were centrifuged at 10.000 rpm for 30 min and the supernatant containing the metabolites were obtained and 100  $\mu$ L of the supernatant was transferred into wells (6 mm diameter) bored in Muller Hinton and potato dextrose agar previously seeded with the spoilage bacteria cells and fungi spores. The culture plates were incubated at 30 °C for 48 h and 7 days respectively and observed for zones of inhibition (Ogunbanwo et al., 2014).

### Characterization of the antimicrobial substance

The isolated crude antimicrobial substance was characterized with respect to the effect of proteolytic enzymes on the bacteriocin activity. Selected enzymes were tested on the cell free supernatant (Bizani and Brandelli, 2002). Proteolytic enzymes including trypsin, pepsin, and papain were dissolved in 40 mM Tris-HCl (pH, 8.2), 0.002 M HCl (pH, 7), and 0.05 M sodium phosphate (pH, 7.0) respectively to a final concentration of 0.1 mg  $ml^{-1}$ . Other enzymes such as lipase and  $\alpha$ -amylase were dissolved in 0.1 M potassium phosphate (pH, 6.0), and 0.1 M potassium phosphate (pH, 7.0) respectively to a final concentration of 0.1 mg/ml. Equal aliquots of both filter sterilized of each test strain and each enzyme solution were mixed, incubated at 30°C for each enzyme for 2 h and heated in boiling water for 5 min to inactivate the enzymes. These sample mixtures and the controls (without enzyme treatment) were inoculated with the indicator strains as previously mentioned and tested for antimicrobial activity by the optical density method (ODM) (Smaoui et al., 2010; Djadouni and Kihel, 2013).

### Shelf life study of tomato paste and sauce

The tomato paste and sauce were boiled for 10 min and dispensed in 20 g amount separately into three pre-sterilized containers. Twenty milliliters of the crude bacteriocin-like substance of *Leuconostoc* spp. were added (v/v) differently to the tomato paste and sauce inoculated with *E. coli*  $10^7$  CFU  $g^{-1}$  (2 ml) and stored at 4°C for 72 h. Microbial load of each treatment was monitored by determining the colony forming unit (CFU  $ml^{-1}$ ) of *E. coli* on the hektoen medium (Safdar et al., 2010; Cottaz et al., 2008).

### Statistical analysis

Data were expressed as mean  $\pm$  standard deviation. Statistical significance was determined using one-way analysis of variance on the replicates, where a p-value of  $\leq 0.05$  was considered significant.

## RESULTS AND DISCUSSION

The results of the physical and chemical analyses of tomato samples are shown in Table 2. The percentage of moisture contents decreased in S02 and S04 samples to 60.30 and 62.50%; this may be due to the boiling at 90°C for 2 h and the production process steps of tomato paste, that is made by cooking tomatoes for several hours to reduce moisture, straining them to remove the seeds and skin (Adam, 1998; Ife Fitz and Bas Kuipers, 2003; Trgera et al., 2007; Souci et al., 2008). Dry matter content was

**Table 2.** Physical-chemical characteristics of tomatoes samples (S01, S02, S03, and S04).

Samples	Moisture content %	Dry matter content (g)	lactic acid content (°D)	pH	Firmness, taste and color
S01	94.20 ± 0.1	5.9 ± 0.1	0.99 ± 0.02	04.20±0.03	No change
S02	60.30 ± 0.1	5.7 ± 0.2	0.75 ± 0.02	04.43±0.03	Change in color
S03	94.20 ± 0.1	5.8 ± 0.1	0.89 ± 0.02	04.28±0.02	No change
S04	62.50 ± 0.1	5.9 ± 0.1	01.58 ± 0.02	03.70±0.01	Change in color, taste, and production of CO <sub>2</sub>

stable along the different time and temperature of storage in S01, S02, S03 and S04 from 5.9 g to 5.7 g, but the pH values decreased to 03.70 in sample S04 that was stored at 4°C for six months, and lactic acid concentration increased to 01.58 D° during this storage period with an important change in the color, taste, and CO<sub>2</sub> production; this changes may be due to the thermal treatment, fermentation process and organic acids (citric acid, ascorbic acid, and lactic acid) produced by LAB during storage in anaerobic conditions in presence of oil and NaCl (Jean-Louis, 2007).

The fresh-cut tomato S03 was not affected during cold storage and shelf-life. On the other hand, S01 and S02 did not show significant differences among them. Color of fresh-cut tomatoes was significantly affected by the antimicrobial treatments and the storage time (Ayala-Zavala et al., 2008). Previous experiments demonstrated that firmness losses of fresh-cut tomato slices were probably linked to the ripening stage of whole fruit at processing and storage temperature. In addition, processing operations may have triggered important losses in membrane integrity due to mechanical stressing of plant tissues. In this regard, diminishing membrane integrity could cause a loss of texture related enzymes and their substrates, leading to a rise in Xuids and solute exchanges as well as an increase in the enzymatic activity (Ayala-Zavala et al., 2008). Results obtained in this study show that transformation process, temperature and storage period can affect the tomato components and quality (Goodman et al., 2002; Chong et al., 2009; Ife Fitz and Bas Kuipers, 2003). During transport and storage, intact fruits and vegetables are prone to deleterious changes induced by respiratory, metabolic and enzymatic activities, as well as by desiccation, pests, microbial spoilage, and temperature-induced injury. Many of these changes adversely affect the antioxidant status of the tomato products (Lindley, 1998). Capanoglu et al. (2010), worked closely with a Turkish tomato paste factory and obtained samples from each step during the paste production process, from the tomatoes arriving at the factory gate to the final canned product. Both targeted (for carotenoids) and untargeted (LC-MS of semi-polar, hydrophilic extracts) metabolic approaches were used to follow biochemical changes after each step.

Results show that a multitude of modifications took place, involving both increases and decreases in individual

components. Those steps causing the greatest changes were identified and predictions were made as to how these steps could be tackled in a modified processing strategy to improve the antioxidant capacity of the end product. Gahler et al. (2003) investigated home preparation methods such as peeling, tomato soup preparation, etc., and three different steps of tomato juice production including sieving, homogenization, sterilization, filling, and pasteurization. Results suggested that homogenization increased the hydrophilic antioxidant capacity of the different tomato products. However, the exact mechanism still remains unclear. Similarly, in industrial processing, Capanoglu et al. (2010) also showed the “breaker” or homogenization step most significantly altered the biochemical composition of tomato paste. Abushita et al. (2000) also analyzed samples taken from three steps of paste processing (raw tomato, crushed sieved puree, and pasteurized paste) which were obtained from a canning factory in Hungary. The results show that the contents of ascorbic acid and tocopherols decreased during processing while carotenoids either remained unchanged or were found to increase.

Absence of *Salmonella*, *S. aureus*, and *Clostridium* in the tomatoes samples (Table 3) showed that thermal treatment and cold storage of tomato inhibit the growth and developments of pathogens and spoilage microorganisms. However, thermal treatments applied during industrial preparation of tomato products may involve various chemical reactions leading to the degradation of these antioxidants. Besides, addition of vegetable oil for the preparation of tomato sauce may lead to lipid oxidation contributing to the micro-constituent instability (Chanforan et al., 2010a and b). The presence of some aerobic mesophilic microorganisms, coliforms, and anaerobic bacteria compared to control samples was related to the averments conditions and preparations methods of tomatoes. Ayala-Zavala et al. (2008) suggest that mechanical damage during minimal processing enhance contamination by epithelial microflora, and promote leaking of nutrients, which are rich substrates to microorganisms, supporting the fast microbial development in contrast with intact tissue.

The highest count of LAB and *Bacillus* in S04 sample was explained by the storage of tomato paste for 06 month at 4°C in the presence of oil and NaCl, which favor the fermentation and organics acids production that

**Table 3.** Detection of pathogens, contaminants, LAB, and *Bacillus* in the tomato samples (S01, S02, S03, and S04).

Samples	Total aerobic bacterial count (CFU ml <sup>-1</sup> )	total coliform count (CFU ml <sup>-1</sup> )	Total anaerobic bacteria count (CFU ml <sup>-1</sup> )	<i>Salmonella</i> (CFU ml <sup>-1</sup> )	<i>Clostridium</i> (CFU ml <sup>-1</sup> )	<i>S. aureus</i> (CFU ml <sup>-1</sup> )	LAB (CFU ml <sup>-1</sup> )	<i>Bacillus</i> (CFU ml <sup>-1</sup> )
S01	86.10 <sup>5</sup>	36.10 <sup>5</sup>	30.10 <sup>5</sup>	ABS	ABS	ABS	ABS	20.10 <sup>5</sup>
S02	55.10 <sup>5</sup>	17.10 <sup>5</sup>	34.10 <sup>5</sup>	ABS	ABS	ABS	20.10 <sup>5</sup>	25.10 <sup>5</sup>
S03	76.10 <sup>5</sup>	ABS	ABS	ABS	ABS	ABS	47.10 <sup>5</sup>	29.10 <sup>2</sup>
S04	ABS	25.10 <sup>5</sup>	36.10 <sup>5</sup>	ABS	ABS	ABS	60.10 <sup>6</sup>	72.10 <sup>5</sup>

\* ABC, absence.

decreased the pH of products and increased also the CO<sub>2</sub> production; these conditions inhibit the growth of contaminants in the tomato paste (Buta and Moline, 1998; Benkeblia, 2004). Also, this LAB and *Bacillus* have antibacterial and antifungal activities against a variety of Gram-negative and Gram-positive bacteria and fungi. Related to fungal development, the highest counts were observed in control samples (Siboukeur, 2011).

A total of 60 isolates were obtained from fermented tomato paste and screened for antimicrobial spectrum against the Gram-positive, Gram-negative bacteria, and fungi using the well diffusion method. The average diameter of the inhibition zone measured ranged from 2 to 4 mm size. One strain of LAB was selected for further studies, because LBc03 contained antimicrobial compound with wide spectrum that inhibited the growth of three indicator strains *E. coli*, *S. aureus*, and *Aspergillus* sp but did not inhibited *Penicillium* sp. and *Clostridium* sp. On the basis on their positive Gram reaction, non-motility, absence of catalase activity and of spore formation, the rod or cocal shape, physiological and biochemical characters as well as sugar utilization pattern, LBc03 was identified as *Leuconostoc* spp. (Table 4). The isolation of LAB species from healthy tomato fruits is in accordance with findings of Sajur et al. (2007) and Settanni and Corsetti, (2008). The presence of LAB in tomato fruit is attributed to their high survival in post harvest conditions of tomatoes (Trias et al., 2008; Ogunbanwo et al., 2014).

The LAB ability to produce antimicrobial compounds is due the absence of true catalyses to break down hydrogen peroxide generated which accumulates and becomes inhibitory to some organisms. The antagonistic activity of LAB metabolites against the spoilage bacteria and fungi agrees with the findings of Trias et al. (2008). The inhibitory effect of lactic acid is due to undissociated forms of the acids which penetrates the pathogen's membrane and liberate hydrogen ion in the neutral cytoplasm thus inhibiting vital cell functions (Corleh and Brown, 1980; Adeniyi et al., 2006). Diacetyl is known to be very effective against fungi and this is due to the interference with the utilization of arginine (De Vyust and Vandamme, 1994) and in addition to a strong oxidizing effect on the organisms cell especially bacteria (Condon,

1987; Sangorin et al., 2014).

The *Leuconostoc* spp. (LBc03) bacteriocin-like substance produced was inactivated by the tested enzymes (Table 5). However, since the activity of the filtrate was not completely inhibited, it is possible that the bacteriocin may also be bound to other molecules like a lipid or a carbohydrate moiety. These data clearly show that the bacteriocin-like substance was of a proteinaceous nature.

Similar results were obtained (Joshi et al., 2006; Diop et al., 2008; khalil et al., 2009; Chanforan et al., 2010a and b; Rakshita, 2011).

Bacteriocin-like substance produced by LBc03 resulted in the decrease of *E. coli* from an initial population of 10<sup>7</sup> to 05.10<sup>3</sup>CFU mL<sup>-1</sup> and 1.10<sup>3</sup> CFU mL<sup>-1</sup> in tomato paste and sauce respectively (Figure 1). This reveals a possible potential of the LAB metabolites in the retardation of food spoilage which agrees with the findings of Ogunbanwo et al. (2008).

The bio-preservative potential of LAB metabolites has been tested on other food product like suya (Adesokan et al., 2008) and chicken meat (Ogunbanwo and Okanlawon, 2006). A major advantage in the use of lactic acid bacteria and their metabolites is that they are considered as generally recognized as safe (GRAS) and comply often with the recommendations for food products (Stiles and Holzapfel, 1997). Unlike some chemical preservatives, LAB metabolites have not been reported to have residual effect on the food product or the consumer's health.

## Conclusion

Our bacteriocin-like substance revealed interesting properties that justifies its importance regarding food safety and protection.

## Conflict of interests

The author(s) did not declare any conflict of interest.


**Table 4.** Morphological, physiological and biochemical properties of LAB isolate (LBc03).

Test	Isolate LBc03
<b>Morphology</b>	<b>Rod, circular and white colonies 3.0mm</b>
<b>Growth at temperature (°C)</b>	
10	+
15	+
30	+
37	+
45	+
<b>Growth at pH</b>	
3.5	-
4.5	+
5.5	+
6.5	+
7.5	+
8.5	+
9.5	w
<b>Growth at NaCl %</b>	
4	+
6.5	+
10	+
15	+
CO <sub>2</sub> from glucose	+
CO <sub>2</sub> from gluconate	+
ADH	+
Citrate	+
Thermoresistance at 60°C for 30 min at 45°C	+
Fermentation Type	He
Fermentation of: glucose, saccharose, and lactose.	+
Lait Sherman (1% BM) at 42°C	+
Lait Sherman (3% BM) at 42°C	-
Hydrolyse of caseine	+
Fructose	+
Mannane	+
Maltose	+
Trehalose	+
Manose	+
Melibiose	-
Palmitine	-
Raffinose	+
Xylose	-
Mannitol	+

+, Growth; -, no growth; w, weak growth; He, hetero-fermentation.

**Table 5.** Effect of enzymes treatment (1 mg/mL) on bacteriocin activity against *S. aureus*. Results are expressed as % of means values of growth reduction (n= 3) ± standard deviations.

Enzymes	Enzyme concentration (1 mg mL <sup>-1</sup> )
Pepsin	38.50 ± 0.5
Trypsin	40.30 ± 0.5
α- Amylase	41.80 ± 0.3
Lipase	40.80 ± 0.5



**Figure 1.** Reduction of *E. coli* population in paste and sauce tomato treated with natural antimicrobial of LBc03 isolate and stored at 4°C for 03 days.

## ACKNOWLEDGEMENT

This research work was supported by Microbiology Research Laboratory, Faculty of Sciences, Es-Senia University, Oran, Algeria.

## REFERENCES

- A.O.A.C. (1984). Official methods of analysis, 14<sup>th</sup> revised edition, Association of Official Analytical Chemists. Washington. D.C. USA G. Eason, B. Noble, and I. N. Sneddon, "On certain integrals of Lipschitz-Hankel type involving products of Bessel functions."
- A.O.A.C. International (1995). Official Methods of Analysis of AOAC International. 16<sup>th</sup> revised edn. Association of Official Analytical Communities. Arlington, VA, USA.
- Abriouel H, Franz C, Ben Omar N, Galvez A (2010). Diversity and applications of *Bacillus* bacteriocins. *FEMS Microbiol. Rev.* 35:201-232.
- Abushita AA, Daood HG, Biacs PA (2000). Change in carotenoids and antioxidant vitamins in tomato as a function of varietal and technological factors. *J. Agric. Food Chem.* 48: 2075–2081.
- Adam E (1998). Solar drying of sliced onion and quality attributes as affected by the drying process and storage conditions. Dissertation. Stuttgart: Universität Hohenheim, Institut für Agrartechnik in den Tropen und Subtropen.
- Adeniyi BA, Ayeni FA, Ogunbanwo ST (2006). Antagonistic activities of lactic acid bacteria isolated from Nigerian fermented dairy food against organisms implicated in urinary tract infection. *Biotechnology* 5(2): 183-188.
- Adesokan IA, Odetoyinbo BB, Olubamiwa AO (2008). Biopreservative activity of lactic acid bacteria on suya produced from poultry meat. *Afr. J. Biotechnol.* 7(20): 3799-3803.
- Aguirre M, Collins MD (1993). Lactic acid bacteria and human clinical infection. *J. Appl. Bacteriol.* 75(2): 95-107.
- Artés F, Conesa MA, Hernandez S, Gil MI (1999). Keeping quality of fresh-cut tomato. *Postharvest Biol. Technol.* 1: 153-162.
- Ayala-Zavala JF, Oms-Oliu G, Serrano IO, González A GA, Alvarez-Parrilla E, Martín-Belloso O (2008). Biopreservation of fresh-cut tomatoes using natural antimicrobials. *Eur. Food. Res. Technol.* 226: 1047-1055.
- Bamidele TA, Adeniyi BA, Ogunbanwo ST, Smith SI, Omonigbehin EA (2011). Antibacterial activities of lactic acid bacteria isolated from selected vegetables grown in Nigeria: A preliminary report Sierra Leone. *J. Biomed. Res.* 3(3): 128-132.
- Benkeblia N (2004). Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium sativum*) *Lebensm. Wiss. Technol.* 37(2):263-268.
- Bernard OE, Obioma EN, Fabian JO (1999). Growth inhibition of tomato-rot fungi by phenolic acids and essential oil extracts of pepper fruit (*Dennetiatripetala*). *Food Res. Int.* 32: 395-399.
- Bizani D, Brandelli A (2002). Characterization of a bacteriocin produced by a newly isolated *Bacillus* sp. strain 8 A. *J. Appl. Microbiol.* 93: 512-519.
- Buta IT, Moline HE (1998). Methyl jasmonate extends shelf life and reduces microbial contamination of fresh-cut celery and peppers. *J. Agric. Food Chem.* 46: 1253-1256.
- Buttery RG (1993). Quantitative and sensory aspects of flavor of tomato and other vegetables and fruits. *In: Flavor Science: Sensible Principles And Techniques.* Acree, T. and Teranishi, R. (eds.). Am. Chem. Soc. Books, Washington, D.C. pp. 259-286.
- Capanoglu E, Beekwilder J, Boyacioglu DR, Hall RD (2010). The Effect of industrial food processing on potential ly health-beneficial tomato antioxidants. *Crit. Rev. Food. Sci. Nutr.* 50: 919–930.
- Chanforan C, Caris-Veyrat C, Dufour C (2010b). Evaluation of the stability of tomato and sunflower oil microconstituents in systems modelling the processing of tomato paste into tomato sauce. UMR 408 UAPV-INRA d'Avignon.
- Chanforan C, Caris-Veyrat C, Dufour C (2010a). Stability of tomato microconstituents (phenolic compounds, carotenoids, vitamins C and E) during processing: studies in model systems, development of a stoichio-kinetic model and validation for the unit step of tomato sauce preparation. UMR 408 UAPV-INRA d'Avignon.
- Chong HH, Simsek S, Reuhs BL (2009). Analysis of cell-wall pectin from hot and cold break tomato preparations. *Food Res. Int.* 42: 770-772.
- Clinton SK (1998). Lycopene; chemistry, biology and implications for human health and diseases. *Nutr. Rev.* 56(2): 35-51.
- Condon S (1987). Responses of lactic acid bacteria to oxygen. *FEMS Microbiol. Lett.* 46(3): 269-280.
- Corleh DA, Brown, MH (1980). pH and activity *In: (Col) Microbial Ecology of Foods,* Silkier. J. H. Academic Press, New York.
- Cottaz A, Oulalal N, Sebti I (2008). Effet des ultra-violets sur les microorganismes de contamination de la tomate. Laboratoire d'InfoRmatique en Image et Systèmes d'information
- Davidson PM, Harrison MA (2002). Resistance and adaptation to food antimicrobials, sanitizers, and other process controls. *Food Technol.*

- 56(11):69-78.
- De Vyust L, Vandamme EJ (1994). Bacteriocins of lactic acid bacteria: Microbiology, genetics and applications. Blackie Academic and Professional, London. pp. 151-221.
- Delarras C (2006). Microbiologie pratique pour le laboratoire d'analyses ou de contrôle sanitaire. ISBN: 978-2-7430-0945-8, Lavoisier 2007, Paris.
- Diop MB, Dauphin DR, Dortu C, Destain J, Tine E, Thonart P (2008). *In vitro* detection and characterization of bacteriocin-like inhibitory activity of lactic acid bacteria (LAB) isolated from Senegalese local food products. Afr. J. Microbiol. Res. 2: 206-216.
- Djadouni F, Kihal M (2013). Characterization and determination of the factors affecting anti-listerial bacteriocins from *Lactobacillus plantarum* and *Pediococcus pentosaceus* isolated from dairy milk products. Afr. J. Food. Sci. 7(2): 35-44.
- Djadouni, F, Kihal M (2012). Antimicrobial Activity of Lactic Acid Bacteria and the Spectrum of their Biopptides against Spoiling Germs in Foods. Braz. Arch. Biol. Technol. 5 (3): 435-443.
- Edwards J, Saltveit M E, Henderson WR (1983). Inhibition of lycopene synthesis in tomato pericarp tissue by inhibitors of ethylene biosynthesis and reversal with applied ethylene. J. Am. Soc. Hort. Sci. 108: 512-514.
- Sangorrín MP, Lopes C A, Vero S, Wisniewsk, M (2014). Cold-adapted yeasts as biocontrol agents: Biodiversity, adaptation strategies and biocontrol potential. Book Part. pp. 441-464.
- Gahler S, Otto K, Bohm V (2003). Alterations of vitamin C, total phenolics, and antioxidant capacity as affected by processing tomatoes to different products. J. Agric. Food Chem. 51: 7962-7968.
- Goodman CL, Fawcett S, Barringer SA (2002). Flavor, viscosity, and color analyses of hot and cold break tomato juices. J. Food. Sci. 67: 404-408.
- Holt JG, Krieg NR, Sneath PHA, Staley JT, Williams ST (1994). Editors. In: *Bergey's Manual of Determinative Bacteriology*. 9<sup>th</sup> ed. Baltimore: Williams and Wilkins. p. 559.
- Jean-Louis I (2007). Les relations aliments, micro-organismes, et consommateurs. Microbiologie alimentaire-STIA 2, Université Montpellier II.
- Joshi VK, Sharma S, Rana N (2006). Production, purification, stability and efficacy of bacteriocin from isolates of natural Lactic acid fermentation of vegetables. Ind. J. Food Technol. Biotechnol. 3 (44): 435-439.
- Khali R, El Bahloul YA, Djadouni F, Hamed Omar S (2009). Isolation and characterization of a bacteriocin produced by a newly isolated *Bacillus megaterium* 19 strains. Pak. J. Nutr. 03 (8): 242-250.
- Lehninger AL (1981). Biochimie bases moléculaires de la structure et des fonctions cellulaires. 2<sup>ème</sup> Ed, Flammarion médecine-science.
- Lindley MG (1998). The impact of food processing on antioxidants in vegetable oils, fruits and vegetables. Trend Food Sci. Technol. 9: 336-340.
- Ogawa JM, Zehr EI, Bird GW, Ritchie DF, Uriu K, Uyemoto JK (1995). Compendium of stone fruit diseases. APSS press. St. Paul, Minnesota. p. 57.
- Ogunbanwo ST, Adebayo AA, Ayodele MA, Okanlawon B M, Edema MO (2008). Effects of lactic acid bacteria and *Saccharomyces cerevisiae* co-culture starters on the nutritional contents and shelf life of cassava-wheat bread. J. Appl. Biosci. 12: 612-622.
- Ogunbanwo ST, Fadahunsi IF, Molokwu A J (2014). Thermal stability of lactic acid bacteria metabolites and its application in preservation of tomato pastes. Malays. J. Microbiol. 10 (1):15-23.
- Ogunbanwo ST, Okanlawon BM (2006). Microbial and sensory changes during the cold storage of chicken meat treated with bacteriocin from *L. brevis* OG1. Pak. J. Nutr. 5(6): 601-605.
- Ogunniyi LT, Oladejo JA (2011). Technical efficiency of tomato production in Oyo State Nigeria. Agric. Sci. Res. J. 1(4):84-91.
- Petro-Turza M (1987). Flavor of tomato and tomato products. Food Rev. Int. 2(3):309-351.
- Pla M, Rodriguez-Lazaro D, Badosa E, Montesinis E (2005). Measuring microbiological contamination in fruit and vegetables. In: Improving The safety of fresh fruit and vegetables. Jongen, W. (ed.). CRC Press, Abington. pp. 147-155.
- Rakshita PK (2011). Antimicrobial activity of lactic acid bacteria and partial characterization of its bacteriocins. Master thesis, New Delhi, Mysore. pp. 14-20.
- Safdar MN, Mumtaz A, Amjad M, Siddiqui N, Hameed T (2010). Development and quality characteristics studies of tomato paste stored at different temperatures. Pak. J. Nutr. 9(3): 265-268.
- Sajur SA, Saguir FM, Manca de Nadra MC (2007). Effect of dominant species of lactic acid bacteria from tomato on the natural development in tomato puree. Food Control 18(5): 594-600.
- Settanni L, Corsetti A (2008). Application of bacteriocins in vegetable food biopreservation. Int. J. Food. Microbiol. 121:123-138.
- Siboukeur A (2011). Etude de l'activité antibactérienne des bactériocines (type nisine) produites par *Lactococcus lactis* subsp *lactis*, isolée à partir du lait camelin. Magister en Biologie. Option: Microbiologie Appliquée. Ouargla. pp. 12-15.
- Smaoui S, Elleuch L, Bejar W, Karray-Rebai I, Ayadi I, Jaouadi B, Mathieu F, Chouayekh H, Bejar S, Mellouli L (2010). Inhibition of Fungi and Gram-Negative Bacteria by Bacteriocin BacTN635 Produced by *Lactobacillus plantarum* sp. TN635. Appl. Biochem. Biotechnol. 9: 88217.
- Sneath PH, Mair NS, Sharpe ME, Holts JG (1986). *Bergey's Manual Systematic Bacteriology*. Vol. 2. Willam and Wilkins, Baltimore.
- Souci, Fachman, Kraut. (2008). La composition des aliments. Tableaux des valeurs nutritives, 7<sup>ème</sup> édition. MedPharm Scientific Publishers / Taylor & Francis, ISBN 978-3-8047-5038-8.
- Spadaro D, Gullino ML (2004). State of the art and future prospects of the biological control of post-harvest fruit diseases. Int. J. Food. Microbiol. 91(2): 185-194.
- Stiles EM, Holzapfel WH (1997). Lactic acid bacteria of foods and their current taxonomy. Int. J. Food. Microbiol. 36(1): 1-2.
- Taylor JH (1987). Text of lectures delivered at the national workshop on fruit and vegetable seedlings production held at NIHORT 9-13 Technicon Instrument Corporation (1975). Industrial method. pp. 155-171.
- Toivonen PMA, DeEl JR (2002). Physiology of fresh-cut fruits and vegetables. In: Lamikanra (ed) Fresh-cut fruits and vegetables: Science, technology and market. CRC Press, Boca Raton. pp. 91-1238.
- Trgera K, Oliver H, Andreas B (2007). Conservation of onion and tomato in Niger assessment of post-harvest losses and drying methods. Conf. Int. Agric. Res. pp.1-7.
- Trias R, Baéras L, Montesinos E, Badosa E (2008). Lactic acid bacteria from fresh fruit and vegetables as biocontrol agents of phytopathogenic bacteria and fungi. Int. J. Microbiol. 11 (4):231-236.
- Wiessinger WR, Chantaranon W, Beuchat LR (2000). Survival and growth of *Salmonella bairdoin* shredded lettuce and diced tomatoes, and effectiveness of chlorinated water as a sanitizer. Int. J. Food. Microbiol. 62 (1-2):123-131.