

Full Length Research paper

# Natural occurrence of AFB<sub>1</sub> in maize and effect of traditional maize flour production on AFB<sub>1</sub> reduction in Malawi

L. Matumba<sup>1</sup>, M. Monjerezi<sup>2\*</sup>, E. Chirwa<sup>3</sup>, D. Lakudzala<sup>4</sup> and P. Mumba<sup>5</sup>

<sup>1</sup>Chitedze Agricultural Research Station, P. O. Box 158, Lilongwe, Malawi.

<sup>2</sup>University of Malawi, Chemistry Department, Chancellor College, P. O. Box 280, Zomba, Malawi.

<sup>3</sup>University of Malawi, Department of Mathematical Sciences, Chancellor College, P. O. Box 280, Zomba, Malawi.

<sup>4</sup>University of Malawi, Department of Physical and Biochemical Sciences, The Malawi Polytechnic, P/bag 303, Chichiri, Blantrye 3, Malawi.

<sup>5</sup>University of Malawi, Department of Basic Sciences, Bunda College of Agriculture, P. O. Box 219, Lilongwe, Malawi.

Accepted 2 November, 2009

**Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) is one of the most harmful mycotoxins to humans, animals, and crops that result in illnesses and economic losses. Maize, the most important staple food in most southern African countries, is susceptible to these fungi infections throughout its growth, harvest, transport and storage. In this study, we report on the incidence of AFB<sub>1</sub> (determined by indirect competitive ELISA) contamination in maize from Malawi and the effect of traditional maize-flour production procedures on the final AFB<sub>1</sub> levels. AFB<sub>1</sub> was detected in 45.3% of the maize samples with 12.3% of them exceeding 5 µg/kg (FAO median AFB<sub>1</sub> MTL). The traditional flour production procedures reduced AFB<sub>1</sub> significantly in the order: soaking of dehulled maize (72.4±5.4, 75.4±3.5 and 80.9±5.3% for 24, 48 and 72 h soaking periods, respectively) > dehulling of maize (mean 29.3±5.4%) > sun drying (11.7% max). Sun drying followed pseudo-first order kinetics in AFB<sub>1</sub>. A maximum AFB<sub>1</sub> reduction of 88.1 ± 3.1% was achieved using a sequence of dehulling, soaking for 72 h and sun drying the flour for 4.5 h.**

**Key words:** Aflatoxin B<sub>1</sub>, maize, Malawi.

## INTRODUCTION

Maize (*Zea mays*) is the major staple food in the developing world, mostly in the Sub-Saharan Africa. In Malawi, maize crop is cultivated on more than 70% of the total arable land and contributes significantly to diets of more than 80% of the population, with per capita consumption of 182 kg per year (Pingali, 2001). However, maize is susceptible to contamination by aflatoxins, fungal metabolites that contaminate agricultural products and threaten food safety. They are produced by three species of *Aspergillus*, namely *A. flavus*, *A. parasiticus* and *A. nomius*. Aflatoxins have been shown in many studies to be hepatotoxic, teratogenic, mutagenic, genotoxic and hepatocarcinogenic, depending on the

duration and level of exposure (Hendrickse, 1997; Peraica et al., 1999; Fung and Clark, 2004).

A series of studies on aflatoxin biomarkers have indicated that extensive exposure to aflatoxins can occur in humans of all ages in Africa (Coulter et al., 1984, 1986; Lamplugh et al., 1988; De Vries et al., 1989; Wild et al., 1991, 1992; Zarba et al., 1992; Jonsyn et al., 1995; Oyelami et al., 1996; El-Sayed et al., 2000; Gong et al., 2002, 2003, 2004). Further, there have been various reported outbreaks of human aflatoxicosis in Africa (Muraguri et al., 1981; Nangindu et al., 1982; Siboe and Muriuki, 1995; Lewis et al., 2005; Probst et al., 2007). In general, the prevention of aflatoxin contamination is not possible, hence salvage mechanisms that reduce the toxin content should be utilised to the maximum extent feasible. The effect of several industrial processing methods on the reduction of aflatoxins in corn has been reported (Scott, 1984; Rustom et al., 1993; Samarajeewa et al., 1990; Rustom, 1997). Overall, varying degrees of

\*Corresponding author. E-mail: [mmonjerez@chanco.unima.mw](mailto:mmonjerez@chanco.unima.mw). Tel: +265 1 524 222/+265 995 491 785. Fax: +265 1 524 046.

**Table 1.** Sections and number of participating households in the survey on incidence of aflatoxin in maize and flour. Refer to Figure 1 for locations.

Location in Mpingu EPA	Number of participating households	
	Maize	Milled maize (flour)
Kalima	11	-
Sinumbi	12	14
Masiye	13	15
Kadzio	13	13
Kafisi	13	-
Likuni	14	-
Kakule	14	-
CARS residence	16	-
Katate	-	15
Kawanda	-	15
Kagwanipenya	-	16
Total	106	88

aflatoxin reduction are reported, most being directly related to the nature of the specific physical or chemical processing technique and commodity.

Traditional methods used in the preparation of the final consumed product have also been reported to reduce aflatoxin content to varying degrees, depending on the process (Sennapa and Nyagahungu, 1982; Njapau et al., 1998; Fandohan et al., 2005; Mutungi et al., 2008). In Southern Africa, the final consumed product from maize is mostly a thick porridge; popularly known as *ugali* (e.g. in Tanzania, Kenya), *nshima* (e.g. in Zambia), *sadza* (e.g. in Zimbabwe) or *nsima* (in Malawi).

In Malawi, little work has been done to assess status of aflatoxin contamination in foodstuffs and this has mainly focused on groundnuts, due to their importance in trade. However, contaminated maize has been blamed for aflatoxic hepatitis outbreaks in Kenya and India (Bulatao-Jayme et al., 1976). This is supported by studies conducted in Thailand, Philippines and USA that revealed that dietary intake of aflatoxins from groundnuts was lower than that from maize (Mehan et al., 1991). A recent survey in Makuani District (Kenya) showed that 35.5% of maize had aflatoxin levels above the WHO MTL of 20 µg/kg with 20.1% presenting aflatoxin above 100 µg/kg (Mwihia et al., 2008). There is therefore need to assess the levels of aflatoxins in maize, as a staple food. The four major naturally occurring aflatoxins are aflatoxin B<sub>1</sub> (AFB<sub>1</sub>), aflatoxin B<sub>2</sub> (AFB<sub>2</sub>), aflatoxin G<sub>1</sub> (AFG<sub>1</sub>) and aflatoxin G<sub>2</sub> (AFG<sub>2</sub>). AFB<sub>1</sub> is the most toxic in this series and has been classified as Group I human carcinogen by the International Agency for Research on Cancer (IARC) in 1987 (IARC, 1987). In this study, we therefore provide a first report on the incidence of AFB<sub>1</sub> contamination in maize from Malawi and the effect of traditional maize-flour production procedures on the final AFB<sub>1</sub> levels.

## MATERIALS AND METHODS

### Description of study site

Malawi is a small developing country situated between Zambia, Mozambique and Tanzania, in Southern Africa (Figure 1). In common with greater part of South Central Africa, the country has distinct dry and wet seasons. The wet season lasts from November to May and the remainder of the year is dry, with temperatures increasing until the onset of the next rains. The study was conducted in Mpingu Agricultural Extension Planning Area (EPA), located about 20 km West of the Capital City, Lilongwe (Figure 1). The EPA is under Lilongwe Agricultural Development Division, one of the main maize production regions in Malawi. It has a mean annual temperature of 20°C, a mean annual relative humidity of 68% and receives a mean annual rainfall of 892 mm, 85% of which falls between November and March. The EPA has a total area of 26, 406 ha, 85.5% of which is arable land with about 14,800 farming families (Government of Malawi, 2008).

### Sampling of flour and stored maize


Ninety households from seven of eighteen randomly selected sections of Mpingu EPA and sixteen households from Chitedze Agricultural Research Station (CARS) staff residence participated in the study (Table 1). From all households that participated in the survey, we collected a 5 kg shelled maize sample (farmers own produce) between 13 and 25 March, 2008. We also collected 51 whole-maize flour (locally known as *mgaiwa* flour) and 37 soaked dehulled maize flour (locally known as *woyera* flour) samples (200 g each) from 6 randomly selected maize mills located in 6 different sections of the EPA between 15 and 18 April, 2008 (Table 1). The samples were collected in khaki paper bags and stored at 0°C at CARS laboratory until analysis for aflatoxin.

### Assessment of effect of traditional maize flour processes on AFB<sub>1</sub> levels

#### Artificial contamination of maize

The results of our survey on aflatoxin incidence in maize showed high heterogeneity and in most cases, the levels of AFB<sub>1</sub> were low to demonstrate statistically significant differences between treatments. We therefore, opted to contaminate Open Pollinated Variety (OPV623) maize acquired from a farmer from the study area in order to increase homogeneity and contamination level. 42.6 kg of maize (11.2% grain moisture content) was soaked in water for 6 h and then air-dried for 2 h, after draining the water. A one-litre suspension of aflatoxinogenic *aspergillus* species, prepared by washing heavily naturally contaminated de-husked maize, was spread over the maize and thoroughly mixed by hand. We then divided the inoculated maize into 10 equal portions, put the portions in moistened gunnysacks (Njapau et al., 1998) and spread them on racks of an incubator (Gallenkamp, Loughborough, UK) to a thickness of 4 - 5 cm. The maize was incubated at 29°C (Njapau et al., 1998) and on the seventh day, 200 mL of water was evenly sprayed over each maize portion to replenish lost moisture. Incubation was terminated after 14 days by air-drying the maize to 11.2% at 45°C. The 10 portions were recombined and thoroughly mixed.

In order to obtain practical levels of natural aflatoxin contamination, the incubated maize was divided into 10 portions by mass of 0.775, 1.549, 2.323, 3.098, 3.872, 4.647, 5.422, 6.196, 6.971 and 7.754 kg (corresponding to mass ratios of 1:1, 1:2, 1:3, 1:4, 1:5, 1:6, 1:7, 1:8, 1:9, 1:10, respectively). To each portion, we added healthy OPV623 maize to make a final mass of 22 kg. To



**Figure 1.** Map of Africa showing the location of Malawi (a); map of Malawi showing location of *Mpingu* EPA (b) and sampled areas (c).

maximize homogeneity the incubated and healthy maize portions were simultaneously poured slowly into a locally fabricated rotating mixer and mixed for 6 min.


#### Preparation of different types of flour

We established through interviews that households in the study area employed three methods of processing the maize into flour. These include (1) simply milling the maize

to make whole meal flour (locally known as *mgaiwa* flour); (2) dehulling, soaking, drying and milling the grit into flour, followed by further drying (locally known as *woyera* flour) and (3) dehulling and then known as *gramil* flour). The three methods are summarized in Figure 2.

In order to simulate the flour production procedures of the village, we ground 4.54 kg from each of the ten portions (of the 22 kg-contaminated maize from §2.3.1 above) in a laboratory mill (Christy and Norris Ltd, Suffolk, UK) to pass sieve #20. From this, we prepared two

different 200 g sub-samples. For the first sub-sample, we took the flour immediately after grinding and stored it at 0°C until aflatoxin assay (corresponding to sequence 1 in Table 2). The second one was taken from a 4½ h sun dried (mean temperature 27.6°C) flour that was evenly spread on 0.558 m<sup>2</sup> new sack surface and turned hourly before storing it at 0°C until aflatoxin analysis (sequence 2 in Table 2). 15 kg of each of the remainder from the ten contaminated maize (portions) was dehulled in a hammer mill located near CARS, using the expertise of three



**Figure 2.** A flow sheet showing processes for production of the three flour types from dry maize, as is the practice in the study area.

**Table 2.** Sequence of procedures for flour production and mean cumulative percentage reduction in AFB<sub>1</sub> concentration during flour production. The AFB<sub>1</sub> levels and reduction percentages displayed are means of 10 replicates. Treatments with different letters in the *Remark* column have significantly ( $p < 0.05$ ) different means according to Tukey's HSD test.

Sequence	Process	Final AFB <sub>1</sub> (µg/kg)	Induced percentage reduction in AFB <sub>1</sub> (%)	Remark
1	Milling	48.41 ± 26.75	0	Represents initial AFB <sub>1</sub> content in Maize
2	Milling→ sun drying (4½ h)	42.1 ± 22.4	11.7 ± 3.2	e
3	Dehulling→ milling	33.9 ± 18.9	29.3 ± 5.4	d
4	Dehulling→ milling→ sun drying (4½ h)	29.4 ± 15.3	37.3 ± 5.6	c
5	Dehulling→ soaking (24 h) → milling	9.5 ± 6.1	80.9 ± 4.4	b
6	Dehulling→soaking (24 h) →milling→sun drying (4½ h)	8.6 ± 5.4	82.6 ± 3.5	ab
7	Dehulling→ soaking (48 h) → milling	8.6 ± 5.5	82.4 ± 3.6	b
8	Dehulling→ soaking (72 h) → milling	7.2 ± 5.3	86.5 ± 3.5	a
9	Dehulling→soaking (72 h) →milling→sun drying (4½ h)	6.4 ± 5.3	88.1 ± 3.1	a

hammer mill located near CARS, using the expertise of three women. Immediately before dehulling, the women added 2 L of water to the maize to enhance the dehulling process, in accordance with everyday village practice. The dehuller separated hulls from endosperm grits with  $78 \pm 4\%$  efficiency. Further separation was achieved manually by use of a winnow. The two sets of bran (from dehuller and winnow) were combined, weighed and vigorously mixed by hand before taking a 2.5 kg sample. The bran sample was ground in a laboratory mill and 200 g of the resultant bran flour was stored at 0°C until aflatoxin assay.


We then divided 10 kg of the dehulled maize into 4 portions of 2.5 kg each using a sample divider (Burrows, Evanston, Illinois). Three of these samples were soaked in 2.5 L of groundwater in separate containers for 24, 48 and 72 h. The pH of soaking water was determined at the beginning of soaking and every 24 h for 72 h. At the end of the soaking period, the water was drained and the

maize washed in a basin until wash water became clear. The maize was then sun dried for 1½ h. After this, we separately ground the three soaked portions and the fourth (non-soaked) portion in a laboratory mill, from which we drew two 200 g samples (one sun dried for 4.5 h and the other non-dried) for each portion. The samples were then stored at 0°C until aflatoxin assay (sequences 3 - 9 in Table 2).

#### DETERMINATION OF AFB<sub>1</sub> AND TOTAL AFLATOXINS

##### AFB<sub>1</sub> by indirect competitive ELISA Method

Indirect competitive ELISA Method (Waliyar et al., 2005) was used to determine AFB<sub>1</sub> in the maize and flour samples. Moisture content of test ground sample was determined by drying in an oven (ISO, 6540: 1980). A sample of flour, equivalent to 20 g dry matter, was



**Figure 3.** Incidence (%) of aflatoxin contamination in stored maize samples collected from farmers own produce in *Mpingu* EPA, Lilongwe, Malawi. ND = not detected

mixed with 100 mL of a mixture of methanol and water (7:3 v/v) containing 0.5% KCl. The mixture was blended at high speed for 2 min and then transferred to a conical flask, shaken for 30 min at 300 rpm and then filtered through Whatman No. 41 filter paper. The filtrate was diluted in PBS-Tween (1:10). 150 µL of Aflatoxin B<sub>1</sub>-Albumin bovine serum (AFB<sub>1</sub>-BSA) conjugate in carbonate coating buffer (100 ng/mL) was dispensed into wells of a 96-well ELISA plate (Immuno 96 Microwell, Nunc-Maxisorp, Denmark) and incubated at 37°C for 1 h. The plate was washed in three changes of PBS-Tween allowing 3 min for each wash. Then 200 µL of 0.2% BSA in PBS-Tween was added to each well and incubated for 1 h. Blocked plate was washed with PBS-Tween only as described above. AFB<sub>1</sub> standard (Sigma AF-1, Sigma-Aldrich Co., St. Louis, MO, USA) was diluted at concentrations ranging from 25 µg/L to 0.0488 µg/L (using 1:2 diluted health maize extract).

Antiserum (50 µl) was added to each dilution of AFB<sub>1</sub> standard (100 µL) and test sample extract (100 µL) in triplicate and incubated at 37°C for 1 h. The plate was emptied and washed again as previously described. To each well 150 µL of goat anti-rabbit IgG labelled with alkaline phosphatase, in PBS-Tween containing 0.2% BSA (1:1000) was added and incubated for 1 h at 37°C. Plates were emptied and washed 3 times with PBS-Tween allowing 3 min for each wash. p-nitrophenyl phosphate (PNPP) (150 µl) prepared in 10% diethanolamine buffer (pH 9.8) was added to each well and plate was kept in dark at 25°C for 1 h. Absorbance was measured at wavelength of 405 nm by means of an automatic ELISA reader (Multiskan EX, Thermo Electron Cooperation, France). AFB<sub>1</sub> content was calculated using a reference curve of optical density (OD) values for AFB<sub>1</sub> standards plotted using graphical software Sigma Plot 8.0 (SPSS Inc., Chicago, IL) with AFB<sub>1</sub> concentration on the x-axis and optical density values on the y-axis. The method's limit of detection (LOD) was interpolated at 1.0 µg/kg. The amount of AFB<sub>1</sub> (ng/g) was calculated as follows

$$AFB_1(ng/g) = \frac{A \times D \times E}{G} \quad (1)$$

Where *A* = AFB<sub>1</sub> concentration in sample extract (ng/mL); *D* = Dilution factor with buffer; *E* = Extraction solvent volume used (mL); and *G* = Dry matter weight equivalent (g).

#### Total aflatoxins by Vicam Fluorometer method

Further, 20 maize samples (from the maize survey) that were analyzed by indirect ELISA procedure for AFB<sub>1</sub> were also tested for total aflatoxins using the fluorometer aflatest immunoaffinity

procedure (VICAM Inc., 1999). A maize flour sample, equivalent to 50 g dry matter, was blended with 5 g of sodium chloride and 100 mL mixture of methanol and water (8:2 v/v) at high speed for 1 min using a blender. The mixture was filtered through fluted filter paper (Whatman 2V, Whatman, Middlex). The filtrate (15 mL) was diluted (1:4) with distilled water, re-filtered through glass microfibre filter paper (Whatman, Middlex, UK). The filtrate (10 mL) was passed through an aflatest immunoaffinity column (VICAM, Watertown, MA, USA) at a rate of 1-2 drops/second. Distilled water (10 mL) was passed through the column at 1-2 drops/second and repeated once more until no bubbles come through the column. Toxin was eluted from the column with 1 mL HPLC grade methanol at the rate of 1-2 drops/second into a glass cuvette, mixed with freshly made 1 mL aflatest developer and its fluorescence measured in a pre-calibrated fluorometer (VICAM V1 series 4, VICAM, Watertown, MA). LOD was interpolated at 1.0 µg/kg.

#### Statistical data analysis

The data from the survey of aflatoxin content in stored maize and flour samples were analysed using Statistical Package for Social Scientists (SPSS) for Windows version 12 (SPSS inc., Chicago, IL, USA). Differences between mean aflatoxin levels were tested by *t*-test. Descriptive statistics such as percentages and frequencies were used to summarize the data. Analysis of data from experiments on effect of different flour making procedures was performed using Stastisca 6.0 (StatSoft Inc.) software. The difference between treatment means was tested by ANOVA and Turkey's HSD test was used as a post ANOVA technique for pairwise comparison of treatments.

## RESULTS AND DISCUSSION

### Occurrence of AFB<sub>1</sub> in stored maize and maize flour

AFB<sub>1</sub> was detected in 45.3% of the 106 stored maize samples analyzed, with 12.3 % of the maize exceeding the median AFB<sub>1</sub> Maximum Tolerable Limit (MTL) of 5 µg/kg (FAO, 2004) (Figure 3). The average AFB<sub>1</sub> in samples from village and CARS households was  $1.76 \pm 3.36$  µg/kg and  $1.48 \pm 1.85$  µg/kg respectively, without significant difference ( $p > 0.05$ ) between them. The combined mean of the CARS and village household samples was  $1.71 \pm 3.17$  µg/kg and the highest contaminated village and CARS sample registered AFB<sub>1</sub> content of 16.9 µg/kg and 5.3 µg/kg, respectively.

Out of the 20 samples analyzed for both AFB<sub>1</sub> and total aflatoxin, twelve samples (60%) had both detectable AFB<sub>1</sub> and total aflatoxin. We applied stepwise linear regression, using SPSS for Windows version 12 to the data for these samples to find a relationship between AFB<sub>1</sub> and total aflatoxin in stored maize from the study area. We obtained a strong and significant ( $F = 68.76$ ,  $p < 0.0001$ ,  $r^2 = 0.78$ ) linear relationship (Figure 4) given by

$$\text{Total aflatoxin} = 1.3838 \times AFB_1 \quad (2)$$

Based on this relationship, we believe the toxin production to be dominated by *A. flavus* species. Malawi has regulation for total aflatoxin only. FAO has expressed need for regulatory authorities in countries, where total aflatoxin regulatory limits apply, to inspect available data




Figure 4. Relationship between AFB<sub>1</sub> and total aflatoxin in stored maize samples.

Table 3. A summary of average reduction (%) in AFB<sub>1</sub> by a single process (dry matter basis). The AFB<sub>1</sub> levels and reduction percentages displayed are means of 10 replicates. Different letters within a row indicate that means are significantly ( $p < 0.05$ ) different according to Tukey's HSD test.

Process	Initial AFB <sub>1</sub> (µg/kg)	Final AFB <sub>1</sub> (µg/kg)	AFB <sub>1</sub> reduction (%)
Soaking grit for 72 h and washing	33.9 ± 18.9	7.2 ± 5.3	80.9 ± 5.3 <sub>a</sub>
Soaking grit for 48 h and washing	33.9 ± 18.9	8.6 ± 5.5	75.4 ± 3.5 <sub>ab</sub>
Soaking grit for 24 h and washing	33.9 ± 18.9	9.5 ± 6.1	72.4 ± 5.4 <sub>b</sub>
Dehulling maize	48.4 ± 26.8	33.9 ± 18.9	29.3 ± 5.4 <sub>c</sub>
Sun drying whole meal flour ( <i>mgaiwa</i> )	48.4 ± 26.8	42.1 ± 22.4	11.7 ± 3.2 <sub>d</sub>
Sun drying non soaked dehulled flour ( <i>gramil</i> )	33.9 ± 18.9	29.4 ± 15.3	11.4 ± 4.2 <sub>d</sub>
Sun drying 72 h soaked grit flour ( <i>woyera</i> )	9.5 ± 6.1	6.4 ± 5.3	11.4 ± 3.3 <sub>d</sub>
Sun drying 24 h soaked grit flour ( <i>woyera</i> )	9.5 ± 6.1	8.6 ± 5.4	9.4 ± 3.6 <sub>d</sub>


on aflatoxins and determine the ratio of AFB<sub>1</sub> to total aflatoxin in order to protect consumers (FAO, 2004). Due to the toxic nature of AFB<sub>1</sub>, setting a separate (lower) level for AFB<sub>1</sub> offers an extra assurance for public safety. This study therefore sets the right direction for AFB<sub>1</sub> regulation in Malawi.

In the survey conducted on the occurrence of AFB<sub>1</sub> in flour, 37 and 27% of *mgaiwa* and *woyera* flour, respectively, presented a detectable AFB<sub>1</sub> level of contamination (Figure 5). The highest contaminated *mgaiwa* sample had 7.1 µg/kg while the highest contaminated *woyera* flour sample had 2.3 µg AFB<sub>1</sub>/kg. The average AFB<sub>1</sub> content in *mgaiwa* samples was 1.00 ± 1.60 µg/kg, which was significantly ( $p < 0.05$ ) higher than in *woyera* flour (0.38 ± 0.69 µg/kg). The low AFB<sub>1</sub> levels in flour samples compared to levels in maize samples point towards the

efficiency of the flour processing methods in reducing aflatoxins.

#### Effect of maize flour production procedures on AFB<sub>1</sub> level in flour

Aflatoxin concentration in the artificially contaminated maize ranged from 8.9 to 87.1 µg/kg. Our results indicate that dehulling of maize, soaking of dehulled maize and drying reduced AFB<sub>1</sub> significantly in the order: soaking of dehulled maize > dehulling of maize > sun drying (Table 3). On average 29.3 ± 5.4% (range 16.6 – 34.9%) of the AFB<sub>1</sub> in the original maize was removed by simple bran removal (dehulling) (Figure 6). The bran represented 26.3% of the original mass of raw maize and contained



**Figure 5.** Incidence (%) of aflatoxin contamination in flour samples collected from hammer mills in Mpingu EPA, Lilongwe, Malawi.

on average  $87.6 \pm 7.8\%$  more AFB<sub>1</sub> than the *mgaiwa* flour (Figure 6 (a)). Our results are in general agreement with previous studies (Bennet and Anderson, 1978; Njapau et al., 1998). Bennet and Anderson (1978) observed slightly higher (30 - 38%) AFB<sub>1</sub> reduction through dehulling of maize.

Siwela et al. (2005) reported as high as 92% reduction of total aflatoxin through dehulling of maize. Ideally, the dehulling process aims at removing all the bran leaving only the endosperm, but this is only achievable in sophisticated commercial systems. The difference in the results could be due to differences in the efficiencies of dehullers used and extent of fungal and toxin penetration in the grains. We did not obtain significant relationship between initial AFB<sub>1</sub> concentration and AFB<sub>1</sub> percentage reduction caused by dehulling process (Figure 7 (a) and Table 4). This is because the dehulling process involves only mechanical removal of bran, the rate of which is independent of the aflatoxin level of the maize. In Malawi, the bran is widely used as animal feed and in times of food shortage people collect it from maize mills and prepare *nsima* from the resulting bran flour.

In these cases, both animals and humans are therefore at risk of aflatoxin toxicity. Further, risk-benefit analysis studies need to be carried out on aflatoxin decontamination through dehulling before it is extensively promoted

as a precautionary measure against aflatoxin exposure as it is associated with nutrient loss. Moeser et al. (2002) observed the loss of neutral detergent fiber (NDF) (61%) and acid detergent fiber (ADF) (32%), lysine (33%) and all other essential amino acids except methionine and tryptophan, following dehulling and de-germing of maize.

Soaking and subsequent washing of dehulled maize removed substantial amounts of AFB<sub>1</sub> content. Soaking dehulled maize for 24, 48 and 72 h reduced AFB<sub>1</sub> content significantly ( $p < 0.05$ ) by  $72.4 \pm 5.4$ ,  $75.4 \pm 3.5$  and  $80.9 \pm 5.3\%$  respectively (Table 3 and Figure 6 (c)). This observation is in agreement with the findings of Bennet and Anderson (1978) and Se (2002) who reported a recovery of 39 - 42% in aflatoxins in steep water and 48% reduction in aflatoxin due to loss of steep water, respectively. Our results of the treatment experiments show no significant difference between soaking for 24 and 48 h and between 48 and 72 h on aflatoxin removal. However, soaking for 72 h yielded significantly ( $p < 0.05$ ) higher AFB<sub>1</sub> reduction than soaking for 24 h (Table 3).

During soaking, pH dropped from 6.7 at the beginning of the soaking period to 5.8 after 24 h, 4.6 after 48 h and 4.3 after 72 h. The drop in pH may be attributed to microbial activity. We did not find a significant relationship between initial AFB<sub>1</sub> concentration and percentage AFB<sub>1</sub> reduction caused by the 24 and 48 h soaking processes,



**Table 4.** A summary of regression equations obtained using stepwise regression analysis using SPSS for Windows 12. The regressions equations are plotted together with experimental data in Figures 7 and 8.

Figure	Plot	Regression equation	Sign.	S.E	F	R <sup>2</sup>	R <sup>2</sup> <sub>adjusted</sub>
Figure 7	Dehulling, Soaking for 24 and 48 h and sun drying <i>woyera</i> flour		No significant regression				
	Soaking for 72 h.	AFB <sub>1</sub> reduction (%) = -0.234[AFB <sub>1</sub> ] <sub>o</sub> + 88.81	0.002	3.005	19.599	0.71	0.67
	Sun drying <i>mgaiwa</i> flour	AFB <sub>1</sub> reduction (%) = 0.106[AFB <sub>1</sub> ] <sub>o</sub> + 6.56	0.001	1.543	30.345	0.791	0.765
	Sun drying <i>gramil</i> flour	AFB <sub>1</sub> reduction (%) = 0.197[AFB <sub>1</sub> ] <sub>o</sub> + 4.70	0.001	2.165	26.652	0.769	0.740
Figure 8	Sun drying <i>Mgaiwa</i> flour	log rate <sub>o</sub> = 1.344log [AFB <sub>1</sub> ] <sub>o</sub> - 2.15	<0.001	0.7767	279.702	0.972	0.969
	Sun drying <i>Gramil</i> flour	log rate <sub>o</sub> = 1.383log [AFB <sub>1</sub> ] <sub>o</sub> - 2.18	<0.001	0.10262	159.052	0.952	0.946
	Sun drying <i>woyera</i> flour (24 h soaking)	log rate <sub>o</sub> = 1.176 log [AFB <sub>1</sub> ] <sub>o</sub> - 1.87	<0.001	0.19411	35.846	0.818	0.795
	Sun drying <i>woyera</i> flour (72 h soaking)	log rate <sub>o</sub> = 1.023log [AFB <sub>1</sub> ] <sub>o</sub> - 2.11	<0.001	0.13657	86.202	0.915	0.904

but for the 72 h soaking process (Figure 7 (b) and Table 4). This suggests that the microorganisms, although present, may not have been involved in reduction of AFB<sub>1</sub> during the 24 and 48 h soaking processes. On the other hand, the 72 h soaking process may have provided sufficient time for degradation of AFB<sub>1</sub> by microbial activity, as a certain time may be needed (lag phase) for cells to adjust to the environment. The reduction of AFB<sub>1</sub> may therefore be attributable to partial solubility of AFB<sub>1</sub> in water (Grant and Phillips, 1998) and microbial activity, which leads to reduced pH.

In this study, a difference in soaking time of 48 h was required to obtain a significant difference in AFB<sub>1</sub> reduction (24 and 72 h soaking periods). This period may have allowed both more AFB<sub>1</sub> to dissolve in the steep-water and for microbial action, hence more AFB<sub>1</sub> removal during the draining of the steep water.

We established, through interviews during the study, that people use steep-water for cooking sour porridge, especially for patients. The steep-water would be a significant source of dietary AFB<sub>1</sub> exposure. Further, interviews conducted with maize mill operators at two maize mills

located in *Biwi* and *Area 18* townships (Lilongwe City) revealed that the majority of the urban dwellers prefer preparing *gramil* to the *woyera* flour.

Sun drying for 4.5 h achieved mean AFB<sub>1</sub> reduction of 11.7 ± 3.2% for *mgaiwa*, 11.4 ± 4.2% for *gramil*, 11.4 ± 3.3% for *woyera* flour (72 h soaking) and 9.4 ± 3.6 % for *woyera* (24 h soaking), without significant differences between the flour types (Table 3 and Figure 6 (d)). The AFB<sub>1</sub> percentage reduction achieved through sun drying increased linearly with initial AFB<sub>1</sub> content in *gramil* (r<sup>2</sup> = 0.79) and *mgaiwa* (r<sup>2</sup> = 0.77) (Figure 7 (d) and Table 4). However, we did not find a significant relationship for sun-drying *woyera* flour (Figure 7 (c) and Table 4) probably due to low initial AFB<sub>1</sub> content and high moisture content. The experimental procedure for the sun-drying processes entailed exposing flour samples with different initial AFB<sub>1</sub> concentration to the sun for similar periods (4.5 h), corresponding to varying initial concentration of AFB<sub>1</sub> in flour samples, keeping the reaction time constant at 4.5 h. We therefore applied the method of initial rates (incorporating the isolation method) to determine the order of reaction for the reduction of AFB<sub>1</sub> with

respect to AFB<sub>1</sub> through the sun-drying process. Firstly, we calculated the rates of reactions from the experimental data using:

$$\text{rate } (\mu\text{g/kg hr}) = \frac{[\text{AFB}_1]_o - [\text{AFB}_1]_{4.5\text{hr}}}{4.5\text{hr}} \quad (3)$$


where [AFB<sub>1</sub>]<sub>o</sub> and [AFB<sub>1</sub>]<sub>4.5hr</sub> denote the initial and final concentration of AFB<sub>1</sub> in the flour sample. The empirical initial rate law for the conversion of AFB<sub>1</sub> may be written as:

$$\text{rate}_o = k [\text{AFB}_1]_o^\alpha \prod_i [X]_{oi}^{\gamma_i} \quad (4)$$

where  $k$  is the rate constant,  $\alpha$  is the order of reaction with respect to AFB<sub>1</sub>, [AFB<sub>1</sub>]<sub>o</sub> is the initial concentration of AFB<sub>1</sub> in the flour sample, [X]<sub>o,i</sub> is the initial concentration of other species  $i$  that may be present to react with AFB<sub>1</sub> and  $\gamma_i$  is the order of reaction with respect to species  $i$ .

Supposing that the concentration of the other species is in excess, (i.e. AFB<sub>1</sub> is isolated) and hence their concentrations do not vary significantly with time, equation 5 leads to:





**Figure 6.** Plots of initial and final AFB<sub>1</sub> content in different flour products produced using traditional methods outlined in Figure 2, depicting the efficiency of the methods in AFB<sub>1</sub> reduction. Dehulling (a) and (b); soaking for 24, 48 and 72 h (c); and sun-drying (d). For all plots, points are experimental data and lines are regression lines. For (c), 24 h soaking: AFB<sub>1</sub>(woyera) = 0.288 x AFB<sub>1</sub>(gramil), R<sup>2</sup> = 0.966; 48 h soaking: AFB<sub>1</sub>(woyera) = 0.261 x AFB<sub>1</sub>(gramil), R<sup>2</sup> = 0.972; 72 hour soaking: AFB<sub>1</sub>(woyera) = 0.227 x AFB<sub>1</sub>(gramil), R<sup>2</sup> = 0.924.


$$\log \text{rate}_o = \alpha \log [AFB_1]_o + \log K \quad (4)$$

where log K is a constant given by

$$\log K = \log k + \log \prod_i^N [X]_{o_i}^{\gamma_i}$$

To find the order of reaction, we plotted log rate<sub>o</sub> as a

function of log [AFB<sub>1</sub>]<sub>o</sub> (equation 5). Further, we applied stepwise linear regression analysis using SPSS for Windows 12.0 to the calculated pairs of log rate<sub>o</sub> and log [AFB<sub>1</sub>]<sub>o</sub>. We found strong and significant linear regressions expressing log rate<sub>o</sub> as a function of log [AFB<sub>1</sub>]<sub>o</sub> for all the flours, with α ≈ 1, indicating a first order reaction in AFB<sub>1</sub> concentration (Table 4 and Figure 8). We note that, in Malawian villages, mgaiwa and gramil



**Figure 7.** Variation of percentage AFB<sub>1</sub> reduction with initial AFB<sub>1</sub> content. Dehulling (a); soaking for 24, 48 and 72 h (b); and sun drying *woyera* flour produced after 24 and 72 h soaking periods (c) and sun drying *mgaiwa* and *gramil* flour (d). For all plots, points represent experimental data and lines are plots of significant linear regression equations summarized in Table 4.


are not sun-dried because they already contain low moisture levels, which are less likely to support microbial growth to spoil the flour.

When the combined effect of the process is considered, the final AFB<sub>1</sub> level in flour samples from our experiments compare very well with those obtain in the maize flour survey. The maximum AFB<sub>1</sub> reduction ( $88.1 \pm 3.1\%$ ) was achieved from combined processes of dehulling, soaking

(72 h) and sun drying of flour (4.5 h) (Table 2 and Figure 9).

## Conclusion

This study has shown that aflatoxin contamination in stored maize is wide spread in *Mpingu* EPA, reflecting




**Figure 8.** Plots of log initial rate and log initial AFB<sub>1</sub> content for sun drying of mgaiwa flour (a), gramil flour (b), woyera flour (24 h soaking) (c), and woyera flour (72 h soaking) (d). The dots represent data from experiment, whilst the straight lines are plots of significant linear regression equations summarized in Table 4.

wide spread contamination in Malawi. Further, we have also shown that despite the prevalence of aflatoxin contamination in stored maize, the traditional methods of maize flour production are effective in reducing AFB<sub>1</sub> content in the final flour and hence in *nsima*, thick porridge made from the flour. The processes of dehulling, soaking, milling and sun drying the resulting flour achieved a combined AFB<sub>1</sub> reduction of 88.1%. The efficiencies of the individual processes were in the order: soaking and drying > dehulling > sun drying. However, the aflatoxin reduction during processing may not necessarily translate into low exposure and risk due to

high consumption.

## ACKNOWLEDGEMENTS

We wish to thank Chitedze Agricultural Research Station (CARS) for granting leave and financial support to L. Matumba to undertake research activities under the Master of Science in Environmental Sciences programme at Chancellor College, University of Malawi. L. Matumba wishes to extend further gratitude to the Director of Agricultural Research Services and colleagues at



**Figure 9.** Variations in AFB<sub>1</sub> content during processing for the production of woyera flour using the village processing techniques. Letters (a, b ... j) denote different levels of contamination in the starting maize.

Chitedze Agricultural Research Station for the support rendered to him during this study.

## REFERENCES

- Bennet GA, Anderson RA (1978). Distribution of aflatoxin and/or zearalenone in wet-milled corn products: a review. *J. Agr. Food Chem.* 26: 1055-1060.
- Bulatao-Jayme J, Almero EM, Salamat L (1976). Epidemiology of primary liver cancer in the Philippines with special consideration of a possible aflatoxin factor. *J. Philippine Med. Assoc.* 52: 129-150.
- Coulter JBS, Suliman GI, Lamplugh SM, Mukhtar B, Hendrickse R.G (1986). Aflatoxins in liver biopsies from Sudanese children. *Am. J. Trop. Med. Hyg.* 35: 360 - 365.
- Coulter JBS, Lamplugh SM, Suliman GI, Omer MIA, Hendrickse RG (1984). Aflatoxins in human breast milk. *Ann. Trop. Paediatr.* 4: 61-66.
- De Vries HR, Maxwell SM, Hendrickse RG (1989). Foetal and neonatal exposure to aflatoxins. *Acta Paediatr. Scand.* 78:373-378.
- European Commission (EC) (2002). Commission Regulation (EC) No. 257/2002 of 12 February 2002 amending Regulation (EC) No. 194/1997 setting maximum levels for certain contaminants in foodstuffs and Regulation (EC) No. 466/2001 setting maximum levels for certain contaminants in foodstuffs. *Official J. Eur. Union.* L41: 12-15.
- European Food Safety Authority (EFSA) (2005). Opinion of the Scientific Committee on a request from EFSA related to A Harmonized Approach for Risk Assessment of Substances, which are both Genotoxic and Carcinogenic. *EFSA J.* 282: 1-31.
- European Food Safety Authority (EFSA) (2007). European Food Safety Authority. Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to the potential increase of consumer health risk by a possible increase of the existing maximum levels for aflatoxins in almonds, hazelnuts and pistachios and derived products. *EFSA J.* 446: 1-127.
- El-Sayed AMAA, Neamat-Allah AA, Soher EA (2000). Situation of mycotoxins in milk, dairy products and human milk in Egypt. *Mycotoxin Res.* 16: 91-100.
- Fandohan P, Zoumenou D, Hounhouigan DJ, Marasas WF, Wingfield MJ, Hell K (2005). Fate of aflatoxins and fumonisins during the processing of maize into food products in Benin. *Int. J. Food Microbiol.* 98: 249-59.
- FAO (2004). Worldwide regulations for mycotoxins in food and feed in 2003. *Food Nutr. Papers.* 81. Food and Agriculture Organisation (FAO).
- Fung F, Clark RF (2004). Health effects of mycotoxins: a toxicological overview. *J. Toxicol. Clin. Toxicol.* 42: 217-234.
- Gong YY, Hounsa A, Egal S, Turner PC, Sutcliffe AE, Hall AJ, Cardwell Gong YY, Cardwell K, Hounsa A, Egal S, Turner PC, Hall AJ, Wild CP (2002). Dietary aflatoxin exposure and impaired growth in young children from Benin and Togo: cross sectional study. *Br. Med. J.* 325

- (7354): 20-21.
- Gong YY, Egal S, Hounsa A, Turner PC, Hall AJ, Cardwell KF, Wild CP (2003). Determinants of aflatoxin exposure in young children from Benin and Togo, West Africa: the critical role of weaning. *Int. J. Epidemiol.* 32(4): 556-562.
- Grant PG, Phillips TD (1998). Isothermal adsorption of aflatoxin B<sub>1</sub> on HSCAS clay. *J. Agr. Food Chem.* 46: 599-605.
- Hendrickse RG (1984). The influence of aflatoxins on child health in the tropics with particular reference to Kwashiorkor. *Trans. R. Soc. Trop. Med. Hyg.* 78: 427-435.
- Hendrickse RG (1997). Of sick turkeys, kwashiorkor, malaria, prenatal mortality, heroin addicts and food poisoning: research on the influence of aflatoxins on child health in the tropics. *Ann. Trop. Med. Parasitol.* 91: 787-793.
- IARC (1987). Overall evaluations of carcinogenicity: An updating of IARC monographs, Vol. 1-42. IARC Scientific Publication, (Suppl. 7). Lyon: IARC
- International Standards Organisation (ISO) (1980). ISO 6540:1980. Maize. Determination of moisture content (on milled grains and on whole grains). ISO.
- Jonsyn FE, Maxwell SM, Hendrickse R G (1995). Ochratoxin A and aflatoxins in breast milk samples from Sierra Leone. *Mycopathol.* 131: 121-126.
- Lamplugh SM, Hendrickse RG, Apeagyei F, Mwanmut DD (1988). Aflatoxins in breast milk, neonatal cord blood, and serum of pregnant women. *Br. Med. J.* 296: 968
- Lewis L, Onsongo M, Njapau H, Schurz-Rogers H, Lubner G, Kieszak S, Nyamongo J, Backer L, Dahiye AM, Misore A, De Cock K, Rubin C, Kenya Aflatoxicosis Investigation Group (2005). Aflatoxin contamination of commercial maize products during an outbreak of acute aflatoxicosis in Eastern and Central Kenya. *Environ. Health Persp.* 11: 1763-1767.
- Mehan VK, MacDonald D, Haravu LJ, Jayanthi S (1991). The groundnut aflatoxin problem: Review and literature database. International Crops Research Institute for the Semi-Arid Tropics (ICRISAT): pp 284.
- Moeser AJ, Kim IB, van Heugten E, Kempen TATG (2002). The nutritional value of degermed, dehulled corn for pigs and its impact on the gastrointestinal tract and nutrient excretion. *J. Anim. Sci.* 80: 2629-2638.
- Muraguri N, Omukoolo LC, Kenji GM, Condiar GA (1981). Survey of mycotoxins in human and animal foods-Part 1. *E. Afr. Med. J.* 58: 484-488.
- Mutungi C, Lamuka P, Arimi S, Gathumbi J, Onyango C (2008). The fate of aflatoxins during processing of maize into muthokoi – A traditional Kenyan food. *Food Cont.* 19: 714-721.
- Mwihia JT, Straetmans M, Ibrahim A, Njau J, Muhenje O, Guracha A. (2008). Aflatoxin levels in locally grown maize from Makueni District, Kenya. *E. Afr. Med. J.* 85: 311-317.
- Nangindu A, Johnson BK, Kenya PR, Ngira JA, Ochieng DM, Nandwa H (1982). Outbreak of acute hepatitis caused by aflatoxin in Kenya. *Lancet.* 1: 1346-1348.
- Njapau H, Muzunguile EM, Changa RC (1998). Effect of village processing techniques on the content of aflatoxins in corn and peanuts in Zambia. *J. Sci. Food Agr.* 76: 450-456.
- Oyelami OA, Maxwell SM, Adeoba E (1996). Aflatoxins and ochratoxin A in the weaning food of Nigerian children. *Ann. Trop. Paediatr.* 16:137-140.
- Peraica M, Radic B, Lucic A, Pavlovic M (1999). Toxic effects of mycotoxins in humans. *Bull. World Health Org.* 77: 754-766.
- Pingali PL (2001). CIMMYT 1999-2000 world maize facts and trends. Meeting world maize needs: technological opportunities and priorities for the public sector. CIMMYT.
- Probst C, Njapau H, Cotty PJ (2007). Outbreak of an acute aflatoxicosis in Kenya in 2004: Identification of the causal agent. *Appl. Environ. Microbiol.* 73: 2762-2764.
- Rustom IYS (1997). Aflatoxin in food and feed: Occurrence, legislation and inactivation by physical methods. *Food Chem.* 59: 57-67.
- Rustom IYS, Lopez-Leiva MH, Nair BM (1993). Effect of pH and heat treatment on the mutagenic activity of peanut beverage contaminated with aflatoxin B<sub>1</sub>. *Food Chem.* 46: 37-42.
- Samarajeewa U, Sen AC, Cohen MD, Wei CI (1990). Detoxification of aflatoxins in foods and feeds by physical and chemical methods. *J. Food Prot.* 53: 489-501.
- Scott PM (1984). Effects of food processing on mycotoxins. *J. Food Prot.* 47: 489-499.
- Sennapa M, Nyagahungu KI (1982). Retention of aflatoxin in ugali and bread made from contaminated maize flour. *Food Sci. Technol.* 19: 64-65.
- Siboe GM, Muriuki GK (1995). Maize flour contaminated with toxigenic fungi and mycotoxins in Kenya. *Afr. J. Health Sci.* 2: 236-241.
- Siwela AH, Siwela M, Matindi G, Dube S, Nsiramazanga N (2005). Decontamination of maize by dehulling. *J. Sci. Food Agr.* 85: 2535-2538.
- VICAM inc. (1999). Aflatest Instruction Manual. Watertown, MA, USA: VICAM.
- Waliyar F, Siambi M, Jones R, Reedy SV, Chibonga D, Kumar PL, Denloye S (2005). Institutionalizing mycotoxin testing in Africa. In Lessie JF, Bandyopadhyay R, Visconti, A. *Mycotoxin detection methods, management and public health and agricultural trade.* Cromwell Press. Pp: 3359-368.
- Wild CP, Rasheed FN, Jawla MFB, Hall AJ, Jansen LAM, Montesano R (1991). In-utero exposure to aflatoxin in West Africa. *Lancet.* 337: 1602.
- Wild CP, Hudson GJ, Sabbioni G, Chapot B, Hall AJ, Wogan GN, Whittle H, Montesano R, Groopman JD (1992). Dietary intake of aflatoxins and the level of albuminbound aflatoxin in peripheral blood in The Gambia, West Africa. *Cancer Epidemiol. Biom. Prev.* 1: 229-234.
- Zarba A, Wild CP, Hall AJ, Montesano R, Hudson GJ, Groopman JD (1992). Aflatoxin M<sub>1</sub> in human breast milk from The Gambia, West Africa, quantified by combined monoclonal antibody immunoaffinity chromatography and HPLC. *Carcinog.* 13: 891-894