

Full Length Research Paper

Influence of Demographic and Academic Variables on Physical Fitness Level of Undergraduates in a Nigerian University

Echezona N.D. Ekechukwu¹, Chinenye D. Oluka², Esther Obidike³, Antoninus O. Ezeukwu¹

¹Department of Medical Rehabilitation, College of Medicine, University of Nigeria Enugu Campus, Enugu, Nigeria; ²Department of Physiotherapy, Bayelsa State Sports Complex, Yenagoa, Bayelsa, Nigeria; ³Department of Physiotherapy, Nnamdi Azikiwe University Teaching Hospital, Nnewi, Nigeria.

Received 15 December, 2018; Accepted 2 May, 2019

Physical inactivity and sedentariness are modifiable risk factors of cardiovascular diseases and a variety of chronic diseases; it is therefore pertinent to improve on one's physical activity and encourage regular assessment of Physical Fitness Level (PFL) especially among young adults. The level of PF and the factors that influence them among Nigerian Undergraduates appear unknown. This study assessed the PFL and the influence of selected demographic/academic factors on the PFL of undergraduates in the Faculty of Health Sciences and Technology (FHST), University of Nigeria, Enugu Campus (UNEC). Two hundred and thirty-eight undergraduates (122 females, and 116 males) recruited from four Departments in the FHST participated in this study. The Physical Activity Readiness Questionnaire (PAR Q) was used to screen the participants for safety of exercise participation, while their demographic details were obtained using data form. The Cooper's 12-Minute Walk Test (12-MWT) was used to assess their Maximal Oxygen Consumption (VO₂ max) - an index of PF. Data obtained were analysed using descriptive statistics (of mean, standard deviation, percentage, frequency) and inferential statistics of (t- test, ANOVA and Pearson's Correlation). Level of significance was set at $\alpha = 0.05$. The participants' mean age, BMI and PFL were 23.79 ± 3.11 years, 22.78 ± 2.93 Kg/m² and 17.75 ± 5.38 ml/Kg/m respectively. There was a significant difference ($t = -6.596$, $p = <0.0001$) between the PF of males and females. There was a significant difference in the PFL of the students when compared across the various departments ($F = 2.321$, $P = <0.0001$) and also when compared across the various grades of study ($F = 4.753$, $P = 0.00$). The relationship between PF and BMI was found to be linear but very weak and non-significant ($r = 0.004$, $p = 0.951$). The PFL of undergraduates in FHST is considered poor for age. Gender and grades are influencing factors of PFL for this population, whereas BMI has no significant relationship with their PFL. Regular Assessment of PFL and improved physical activities/exercise geared towards improving their PFL should be integrated into the programme of students in FHST. Replication of similar studies in other faculties is encouraged.

Key words: Physical Fitness, Physical Activity, Demographic variables, Academic Variables, Nigerian Undergraduates.

INTRODUCTION

Fitness is a broad term denoting dynamic qualities that allows an individual to study his/her needs regarding mental and emotional stability, social consciousness and adaptability, spiritual and moral fibres and physical health; consistent with his/her hereditary endowment (Pate, 1998; Sharma, 2015). Physical fitness is the ability to perform one's Activities of Daily Living (ADL), enjoy leisure and have enough vigour left to care for unforeseen emergencies without undue fatigue (Prentice, 1994; Hoeger et al, 2018). Cardiorespiratory endurance is the most important index used in the assessment of physical fitness (Charles et al, 2004; Domínguez et al, 2018). Cardiorespiratory endurance capacity is best indicated by the Maximum Capacity of the working tissue to use oxygen – V_{O2max} (Bennett et al, 2016).

Physical inactivity is a modifiable risk factor of cardiovascular diseases (Truthmann et al, 2015) and a widening variety of other chronic diseases including diabetes mellitus (Gilbert et al, 2019), colon (Richardson et al, 2016) and breast (Tamimi et al, 2016) cancer, obesity (Thasanasuwan et al, 2016), hypertension (Ibekwe, 2015), bone and joint diseases (Palazzo et al, 2016) and depression (Gallagher et al, 2016). Various studies have shown that there appears to be a linear relationship between Physical Activity (PA) and health status, such that a further increase in PA will lead to additional improvements in health status (Cooper, 1968; Sulaiman et al, 2010; Adeniyi et al, 2011; Odole et, 2019). Cooper (1968) noted this among individuals with COPD. Adeniyi et al (2011) investigated the relationship between Depression and Physical Activity among adolescents in Nigeria while Suleiman et al (2010) studied the relationship between physical fitness and duration of incarceration of inmates in some selected prisons in Nigeria.

Aside the relationship between physical activity and health status, studies (Grison, 2005; Coe et al, 2006; Castelli et al, 2007; Viaderi, 2008; Sardinha et al, 2016; Donnelly et al, 2016) have shown a linear relationship between physical activity/fitness and academic performance among students. In a study of 214 sixth-grade students, Coe et al (2006) found that students who

either performed some or met Healthy People 2010 guidelines for vigorous activity (20 minutes per session on three or more days per week) had significantly higher grades than students who performed no vigorous physical activity in both semesters. Similarly, Castelli et al (2007) found that among 259 third and fifth graders from four Illinois middle schools, field tests of physical fitness were positively related to academic achievement. Specifically, aerobic capacity was positively associated with achievement, whereas BMI was inversely related. Associations were demonstrated in total academic achievement, mathematics achievement, and reading achievement. The brain is known to benefit in many ways from exercise (Hamzat et al, 2014; Hamzat and Ekechukwu, 2015; Sleiman et al, 2016; Barha et al, 2017)

Several studies as enumerated above have shown that improved physical fitness can lead to better health status and more importantly improved academic performance among students. It is therefore pertinent to regularly assess students' level of physical fitness as well as the factors that may influence their physical fitness. This study assessed PFL, and influencing clinical/demographic factors of the PFL of undergraduate students in the Faculty of Health Sciences and Technology (FHST), University of Nigeria, Enugu Campus (UNEC).

MATERIALS AND METHODS

Participants

Two hundred and thirty-eight students participated in the study. Sample of convenience was used to recruit the participants (2nd to 5th year students) from the four undergraduate departments: Medical Rehabilitation (MRH), Medical Laboratory Sciences (MLS), Medical Radiography and Radiological Sciences (RAD) and Nursing Sciences (NUR) in FHST, UNEC.

Eligibility Criteria

Only Undergraduates from the Faculty of Health Sciences and in grade 2 to grade 5 (200 level – 500 level) were included in this study. However, those that answered 'Yes' to one or more questions on the Physical Activity Readiness Questionnaire (PAR-Q) or had any obvious cardiorespiratory disorders with emphasis on the absolute and relative contraindications to exercise testing as stipulated by the American Heart Association (Garner et al, 2017;

*Corresponding author. E-mail: talk2chv013@gmail.com; Tel: +2347065786334.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

Karnabi, 2017) and students with any known musculoskeletal disorder that limits walking were excluded from this study.

Study Design

A cross sectional exploratory research design was used in this study and purposive sampling technique was employed.

Determination of Sample Size

To estimate sample size, a power analysis was done using the mathematical relationship described by Daniel and Cross (2018) for a finite population:

$$n = \frac{N}{1 + N(e)^2}$$

where n = sample size
N = Population Size
e = Tolerance Error (0.065)

The population of students in the FHST, UNEC was 3, 800 as obtained from the faculty officer's record.

Hence, N = 3,800.

Calculated Sample Size (n) = 223.

Procedure

Ethical approval for this study was obtained from the University of Nigeria Teaching Hospital Research and Ethics Committee. The protocol for this study was explained to the prospective participants and their informed consent consequently sought and obtained. They were screened using the PAR-Q and the absolute and relative contraindications to exercise testing. Their demographic details (age, gender, height and weight) and academic details (department, grade) were obtained and recorded in a data form. Anthropometric variables of height and weight were obtained following standard protocol (Ezeukwu et al, 2015) and BMI was calculated thereafter from these two variables using the BMI formula [BMI = weight/(height)²]. The Cooper's 12Minutes' Walk Test (12-MWT) was then administered and their PFL determined using the VO₂ max estimated from the distance walked in 12 minutes (Cooper, 1968).

Data Analysis

Data was analysed using: (1) descriptive statistics of mean, standard deviation, and variance to summarise the demographic and academic variables; (2) independent t – test to compare the PFL of male and female participants; (3) One way ANOVA to compare the mean PFL across the various departments and grades; (4) Pearson correlation coefficient was used to evaluate the relationship between BMI and PFL. Level of significance was set at $\alpha = 0.05$.

RESULTS

Demographic and academic details of the participants

A total Two hundred and thirty-eight undergraduates from

the FHST, UNEC participated in this study. They comprised 122 (51.3%) females and 116 (48.7%) males. The mean age of the participants was 23.79±3.11 years and ranged between 19 and 44 years. Their mean heights and weights were 1.68±0.08m and 64.76±10.11 kg respectively. The estimated mean BMI of the participants was 22.78±2.93 kg/m² (13.42 – 37.91 kg/m²) as shown in Table 1.

Physical fitness level of the participants

The mean PFL of students in this study was 17.75 ± 5.38 ml/kg/min, it ranged between 3.17 to 31.68 ml/kg/min. The PFL of male participants (19.92 ± 5.92 ml/kg/min) was higher than those of the female participants (15.67 ± 3.80 ml/kg/min). Students in the departments of MRH (19.65±5.19 ml/kg/ml) and MLS (19.42±5.33 ml/kg/ml) had higher PFLs than those in RAD (17.01±4.54 ml/kg/ml) and NUR (14.66±4.92 ml/kg/ml). In the same vein, the participants in grades five (19.93±5.62 ml/kg/ml) and three (17.27±5.62 ml/kg/ml) had higher levels of physical fitness than those in grades four (16.99±4.56 ml/kg/ml) and two (16.85±4.88 ml/kg/ml) as shown in Table 1.

Comparison of the PFL of male and female participants using unpaired t-test

There was a significant difference in the mean PFLs between the male and female participants ($t = -6.596$, $p < 0.0001$). The level of physical fitness of males (19.92 ± 5.92 ml/kg/min) was significantly higher than those of the females (15.67 ± 3.80 ml/kg/min) as shown in Table 2

Comparison of the PFL of among the various departments and grades using one-way ANOVA

There was a significant difference in the mean PFLs of the participants when compared among the various departments ($F = 2.321$, $P < 0.0001$); Participants in MRH and MLS had significantly higher values than those in RAD and NS. Also, there was a significant difference in the mean PFLs of the participants when compared among the various grades of study ($F = 4.753$, $P < 0.0001$) as shown in Table 3

Relationship between physical fitness and body mass index (BMI) of the participants

There was a weak, linear but non- significant correlation ($r = 0.004$, $p = 0.951$) between BMI and the level of physical

Table 1: Demographic/Academic Characteristics and the Physical Fitness Level of the Participants (N = 238)

	Min.	Max.	Mean	SD
Age (yrs)	19.00	44.00	23.79	3.11
Height (m)	1.47	1.89	1.68	0.08
Weight (kg)	42.0	107.0	64.76	10.11
BMI (Kg/m ²)	13.41	37.91	22.78	2.93
PFL (ml/kg/min)	3.17	31.68	17.75	5.37
Males			19.92	5.92
Females			15.67	3.80
MRH			19.65	5.19
MLS			19.42	5.33
RAD			17.01	4.54
NUR			14.66	4.92
2 nd year			16.85	4.88
3 rd year			17.27	5.84
4 th year			16.99	4.56
5 th year			19.93	5.62

fitness of the participants as shown in Table 4.

Comparison of the PFL of male and female participants using unpaired t-test

There was a significant difference in the mean PFLs between the male and female participants ($t = -6.596$, $p < 0.0001$). The level of physical fitness of males (19.92 ± 5.92 ml/kg/min) was significantly higher than those of the females (15.67 ± 3.80 ml/kg/min) as shown in Table 2.

Comparison of the PFL of the participants among the various departments and grades using one-way ANOVA

There was a significant difference in the mean PFLs of the participants when compared among the various departments ($F = 2.321$, $P < 0.0001$); Participants in MRH and MLS had significantly higher values than those in RAD and NUR. Also, there was a significant difference in the mean PFLs of the participants when compared among the various grades of study ($F = 4.753$, $P < 0.0001$) as shown in Table 3.

Relationship between physical fitness and body mass index (BMI) of the participants

There was a weak, linear but non-significant correlation ($r = 0.004$, $p = 0.951$) between BMI and the level of physical

fitness of the participants as shown in Table 4.

DISCUSSION

The mean fitness level of students (17.75 ± 5.38 ml/kg/min) in FHST, UNEC as obtained from this study was below the normal $VO_2\max$ (38 to 46 ml/Kg/min) for most college-aged men and women in the United States as reported by Prentice (Prentice 1994). This difference may be due to anthropometric difference (height, limb length and stride lengths) between the two comparing populations (NCD Risk Factor Collaboration, 2016). Also, the average American college students engages in extracurricular sporting activities (Craft, 2012) that is unlikely observed in the study population. There appears to be a general belief in most academic institutions in Nigeria especially among the students in Health Sciences, that extra-curricular activities like exercise and sports are time consuming and so should be relegated to the background in order to have a better academic performance. Studies (Viadero, 2008; Erickson et al, 2015; Mullender-Wijnsma et al, 2015; Donnelly et al, 2016; Erwin et al, 2017;) have shown that the above belief is not only erroneous but detrimental to academic excellence. These studies reported positive linear relationship between participation in physical activities to improve physical fitness and academic performance of students.

Table 2: Independent t – test Result comparing the Physical Fitness Levels of the male and female participants

VO ₂ max	t - value	DF	P – Value	MD	SED	95 % CI	
						Lower	Upper
Equal Variances Assumes	-6.596	236	*<0.0001	-4.234	0.641	-0.642	-0.297

Table 3: One-way ANOVA Comparing the Relationship between the Mean Physical Fitness Level of Participants across the Various Departments and Grades of study.

		Sum of Squares	Df	Mean Square	F	Sig.
DEPT	Btw Groups	267.176	172	1.553	2.321	*<0.0001
	Within Groups	43.500	65	0.669		
	Total	310.676	237			
GRADE	Btw Groups	289.286	172	1.682	4.753	*<0.0001
	Within Groups	23.000	65	0.354		
	Total	312.286	237			

Table 4: Pearson Correlation between BMI and PFL of Participants

BMI	Pearson Correlation (r)	PFL
	Sig. (2-tailed)	-0.004
	N	0.951
		238

In 2004 Naperville Central High School in Illinois began a "learning readiness" physical education program (Viaderi, 2008). Students identified as underperforming in literacy were offered an early morning physical education session immediately followed by a literacy support class. By the end of the semester, it was found that the students who took part in both the early morning physical education class and the literacy intervention gained 1.34 years of improvement on a standardized reading test whereas their peers who did not participate in the physical education class prior to literacy instruction only made about 0.7 years of improvement. A more impressive result was obtained when this approach was applied for mathematics instruction. In a Quebec study, Trudeau and Shepard (2008) concluded that given competent providers, physical activity can be added to the school

curriculum by taking time from other subjects without risk of hindering student academic achievement. On the other hand, they observed that adding time to academic subjects by taking time from physical activity programs does not enhance grades, but may rather be detrimental to health.

The mean PFL of male participants in this study was significantly higher than the mean PFL of female participants. This is because males are larger, with more muscle mass thus more strength and power (Andrew, 2018); they also have larger heart volumes (Hale, 2003; Best et al, 2012) which gives them a greater oxygen transport capacity. Women's performance times are about 10% lower than men's in most track events while absolute maximal oxygen consumption is more than 40% greater in men than women (Birtch et al, 2005). Also, the amount of

and concentration of haemoglobin are higher in men; the average haemoglobin concentration for females is about 13.7 gHb/100 ml while for males, it is about 15.8gHb/100ml (WHO, 2011). This difference is attributed to the stimulating effect of androgens on haemoglobin production and the effects of menstrual blood loss as well as differences in dietary intake (Prentice, 2001; Hall, 2015).

The difference in the mean fitness level across the departments could be that the students may have had different orientation about physical activity/fitness. It is also possible that some courses (e.g. MRH with the highest value) may have exposed the students to the importance of physical activities (Adeniyi et al, 2014) coupled with the presence of a departmental gymnasium available for their utilization. It is also possible that some students may have shown greater degrees of motivation to the 12MWT, which is an important factor that affects its result (Noonan and Dean, 2000; Cazzola et al, 2008). On the other hand, these results may be reflective of the academic workloads or requirements of the students. The low values seen among students in NUR department may be explained by the fact that a good number of the students in this department are females and may have as well reflected in their sampling. Therefore, report from this study that low levels of physical fitness among the female participants may be leaned upon to explain the low value among the NUR students.

There was a significant difference in the mean fitness level across the different grades of study. This may be explained by the differences in academic workloads/expectations and sedentary lifestyle that characterizes the different grades. Students in grade five (with the highest values) may have been more physically active given that they are engaged in clinical postings and research field work (data collection for projects) unlike their colleagues in second grade who are in the preclinical class with perceived greater academic expectation of the preclinical courses and examination that may encourage greater sitting time for personal studies (i.e. increased sedentary lifestyle).

The correlation between BMI and the level of physical fitness in this study was not significant. This may imply that people with higher BMI (e.g. Obese) may not necessarily have lower PFL than those with lower BMI (e.g. Normal). However, this result should be interpreted with caution because the sample selection did not stratify the participants based on BMI. More so, some studies (Frey and Chow, 2006; Huang and Malina, 2007) found

inverse relationship between BMI and Physical fitness. Further studies to elucidate on the relationship between BMI and physical fitness is encouraged.

CONCLUSION

The fitness level of students in the Faculty of Health Science and Technology, UNEC, Nigeria is low. Academic work load in the various departments and grades appears to be a major modifiable factor influencing their level of physical fitness. It is therefore pertinent that physical activities and fitness of these students should be factored while drawing up their academic programmes.

ACKNOWLEDGEMENT

We wish to acknowledge and appreciate all the students that participated in this study.

REFERENCES

- Adeniyi, A.F., Ekechukwu, N.E., Umar, L. and Ogwumike, O.O., 2013. Research profile of physiotherapy undergraduates in Nigeria. *Education for Health*, 26(1), p.15.
- Adeniyi AF, Okafor NC and Adeniyi CY. Depression and Physical Activity in a Sample of Nigerian Adolescents: Levels, Relationships and Predictors. *Child and Adolescent Psychiatry and Mental Health*. 2011; 5:16.
- Akindede MO. Comparison of Cardiorespiratory Fitness of Nigerians with Low Back Pain and Apparently Healthy Adults. *Nig J Health Biomed Sci*. 2004; 3(2): 105 – 107.
- Andrew L. 2018. Physiological Differences Between Male and Female Athletes; Updated June 28, 201. <https://work.chron.com/physiological-differences-between-male-female-athletes-20627.html> accessed on 25/07/19 at 18:53GMT
- Barha, C.K., Galea, L.A., Nagamatsu, L.S., Erickson, K.I. and Liu-Ambrose, T., 2017. Personalising exercise recommendations for brain health: considerations and future directions. *Br J Sports Med*, 51(8), pp.636-639.
- Bennett, H., Parfitt, G., Davison, K. and Eston, R., 2016. Validity of submaximal step tests to estimate maximal oxygen uptake in healthy adults. *Sports Medicine*, 46(5), pp.737-750.
- Best, S., Okada, Y., Galbreath, M.M., Jarvis, S.S., Bivens, T.B., Levine, B.D. and Fu, Q., 2012. The effect of gender and age on hemodynamics, blood volume and cardiac size in healthy humans.
- Birch K, Maclaren D, George K. *Sports and Exercise Physiology*. New York: Garland Science / BIOS Scientific Publishers; 2005
- Castelli DM, Hillman CH, Buck SM, Erwin H. Physical fitness and academic achievement in 3rd and 5th grade students. *Journal of Sport and Exercise Psychology*. 2007; 29: 239-252.
- Cazzola, M., Biscione, G.L., Pasqua, F., Crigna, G., Appodia, M., Cardaci, V. and Ferri, L., 2008. Use of 6-min and 12-min walking test for assessing the efficacy of formoterol in COPD. *Respiratory*

- medicine, 102(10), pp.1425-1430.
- Charles B C, Ruth L, Greg Welk. Concepts of Physical Fitness, Active Lifestyle for Wellness. 10th ed. United States of America: McGraw-Hill Companies Inc; 2004
- Coe DP, Ivarnik JM, Womack C, J, Reeves MJ, Malina RM. Effects of physical education and physical activity levels on academic achievement in children. *Medicine & Science in Sports and Exercise*. 2006; 38: 1515-1519.
- Cooper KH. A Means of Assessing Maximal Oxygen Intake: Correlation Between Field and Treadmill Testing. *JAMA*. 1968; 203: 201 – 204.
- Craft, S.W., 2012. The impact of extracurricular activities on student achievement at the high school level.
- Daniel, W.W. and Cross, C.L., 2018. *Biostatistics: a foundation for analysis in the health sciences*. Wiley.
- Domínguez, R., Cuenca, E., Maté-Muñoz, J., García-Fernández, P., Serra-Paya, N., Estevan, M., Herreros, P. and Garnacho-Castaño, M., 2017. Effects of beetroot juice supplementation on cardiorespiratory endurance in athletes. A systematic review. *Nutrients*, 9(1), p.43.
- Donnelly, J.E., Hillman, C.H., Castelli, D., Etnier, J.L., Lee, S., Tomporowski, P., Lambourne, K. and Szabo-Reed, A.N., 2016. Physical activity, fitness, cognitive function, and academic achievement in children: a systematic review. *Medicine and science in sports and exercise*, 48(6), p.1197.
- Donnelly, J.E., Hillman, C.H., Castelli, D., Etnier, J.L., Lee, S., Tomporowski, P., Lambourne, K. and Szabo-Reed, A.N., 2016. Physical activity, fitness, cognitive function, and academic achievement in children: a systematic review. *Medicine and science in sports and exercise*, 48(6), p.1197.
- Erickson, K.I., Hillman, C.H. and Kramer, A.F., 2015. Physical activity, brain, and cognition. *Current opinion in behavioral sciences*, 4, pp.27-32.
- Erwin, H., Fedewa, A. and Ahn, S., 2017. Student academic performance outcomes of a classroom physical activity intervention: A pilot study. *International Electronic Journal of Elementary Education*, 4(3), pp.473-487.
- Ezeukwu, A.O., Ezeoranu, C.G., Egwuonwu, A.V., Ugwoke, U.M., Ekechukwu, N.E. and Nwankwo, M.J., 2015. Comparison of body fat percentages in Nigerian obese females using field methods. *J. Health Sci*, 5, pp.18-23.
- Frey GC, Chow B. Relationship Between BMI, Physical Fitness and Motor Skills in Youth with Mild Intellectual Disabilities. *International Journal of Obesity*. 2006; 30: 861 – 867.
- Gallagher, D., Kiss, A., Lancot, K. and Herrmann, N., 2016. Depressive symptoms and cognitive decline: a longitudinal analysis of potentially modifiable risk factors in community dwelling older adults. *Journal of affective disorders*, 190, pp.235-240.
- Garner, K.K., Pomeroy, W. and Arnold, J.J., 2017. *Exercise Stress Testing: Indications and Common Questions*. American family physician, 96(5).
- Gilbert, L., Gross, J., Lanzi, S., Quansah, D.Y., Puder, J. and Horsch, A., 2019. How diet, physical activity and psychosocial well-being interact in women with gestational diabetes mellitus: an integrative review. *BMC pregnancy and childbirth*, 19(1), p.60.
- Grissom, J. Physical fitness and academic achievement. *Journal of Exercise Physiology online*. 2005;8(1).
- Hale T. *Exercise Physiology – A Thematic Approach*. England: John Wiley and Sons Ltd; 2003
- Hall, J.E., 2015. Guyton and Hall textbook of medical physiology e-Book. Elsevier Health Sciences.
- Hamzat, T.H.K. and Ekechukwu, N.E., 2015. Aerobic exercise training in stroke rehabilitation: any gap in knowledge. *Nigerian Journal of Medical Rehabilitation*.
- Hamzat, T.K., Ekechukwu, N.E. and Olaleye, A.O., 2014. Comparison of community reintegration and selected stroke specific characteristics in Nigerian male and female stroke survivors. *African Journal of Physiotherapy and Rehabilitation Sciences*, 6(1-2), pp.27-31.
- Hoeger, W.W., Hoeger, S.A., Hoeger, C.I. and Fawson, A.L., 2018. *Lifetime physical fitness and wellness*. Cengage Learning.
- Huang YC, Malina RM. BMI and Health-Related Physical fitness in Taiwanese 9 -18 Years. *Med. Sci. Sports Exerc*. 2007; 39(4): 701 – 708.
- Ibekwe, R.U., 2015. Modifiable risk factors of hypertension and socio demographic profile in Oghara, Delta state; prevalence and correlates. *Annals of medical and health sciences research*, 5(1), pp.71-77.
- Karnabi, E., 2017. *Exercise Stress Testing*. In *Cardiology Procedures* (pp. 35-44). Springer, London.
- Mullender-Wijnsma, M.J., Hartman, E., de Greeff, J.W., Bosker, R.J., Doolaard, S. and Visscher, C., 2015. Improving academic performance of school-age children by physical activity in the classroom: 1-year program evaluation. *Journal of School Health*, 85(6), pp.365-371.
- NCD Risk Factor Collaboration, 2016. A century of trends in adult human height. *Elife*, 5, p.e13410.
- Noonan V, Dean E. Submaximal Exercise Testing: Clinical Application and Interpretation. *Phy Ther*. 2000; 80(8): 782 – 807.
- Odole, A., Ekediegwu, E., Ekechukwu, E.N.D. and Uchenwoke, C., 2019. Correlates and predictors of pain intensity and physical function among individuals with chronic knee osteoarthritis in Nigeria. *Musculoskeletal Science and Practice*, 39, pp.150-156.
- Palazzo, C., Nguyen, C., Lefevre-Colau, M.M., Rannou, F. and Poiraudou, S., 2016. Risk factors and burden of osteoarthritis. *Annals of physical and rehabilitation medicine*, 59(3), pp.134-138.
- Pate RR. A New Definition of Youth Fitness. *Phys Sports Med*. 1998; 11: 77
- Prentice W. *Fitness for College and Life*. 4th ed. United States of America: Mosby Inc; 1994
- Prentice W. Physical Fitness and Activity as Separate Heart Disease Risk Factors: a meta-analysis: *Med Sci Sports Exerc*. 2001; 33: 754 - 761.
- Richardson, A., Hayes, J., Frampton, C. and Potter, J., 2016. Modifiable lifestyle factors that could reduce the incidence of colorectal cancer in New Zealand. *Ann Richardson*, 129(1447).
- Sardinha, L.B., Marques, A., Minderico, C., Palmeira, A., Martins, S., Santos, D. and Ekelund, U., 2016. Longitudinal relationship between cardiorespiratory fitness and academic achievement. *Medicine and science in sports and exercise*, 48(5), p.839.
- Sharma, J., 2015. *HEALTH, WELLNESS, FITNESS AND HEALTHY LIFESTYLES*. Horizon Books (A Division of Ignited Minds Edutech P Ltd).
- Sleiman, S.F., Henry, J., Al-Haddad, R., El Hayek, L., Haidar, E.A., Stringer, T., Ulja, D., Karuppagounder, S.S., Holson, E.B., Ratan, R.R. and Ninan, I., 2016. Exercise promotes the expression of brain derived neurotrophic factor (BDNF) through the action of the ketone body β -hydroxybutyrate. *Elife*, 5, p.e15092.
- Sulaiman AO, Hanif S, Lamina S, Isa UL. Correlates of Selected Indices of Physical Fitness and Duration of Incarceration Among Inmates in some Selected Nigeria Prisons. *Ethiop J Health Sci*. 2010; 20(1): 65 – 69.
- Tamimi, R.M., Spiegelman, D., Smith-Warner, S.A., Wang, M., Pazaris, M., Willett, W.C., Eliassen, A.H. and Hunter, D.J., 2016. Population attributable risk of modifiable and nonmodifiable breast cancer risk

- factors in postmenopausal breast cancer. *American journal of epidemiology*, 184(12), pp.884-893.
- Thasanasuwan, W., Srichan, W., Kijboonchoo, K., Yamborisut, U., Wimonpeerapattana, W., Rojroongwasinkul, N., Khouw, I.T. and Deurenberg, P., 2016. Low Sleeping Time, High TV Viewing Time, and Physical Inactivity in School Are Risk Factors for Obesity in Pre-Adolescent Thai Children. *Journal of the Medical Association of Thailand= Chotmaihet thangphaet*, 99(3), pp.314-321.
- Thornton, J.S., Frémont, P., Khan, K., Poirier, P., Fowles, J., Wells, G.D. and Frankovich, R.J., 2016. Physical activity prescription: a critical opportunity to address a modifiable risk factor for the prevention and management of chronic disease: a position statement by the Canadian Academy of Sport and Exercise Medicine. *Br J Sports Med*, 50(18), pp.1109-1114.
- Trudeau F, Shephard RJ. Physical education, school physical activity, school sports and academic performance. *International Journal of Behavioral Nutrition and Physical Activity*. 2008; 25(5):10.
- Truthmann, J., Busch, M.A., Scheidt-Nave, C., Mensink, G.B., Gößwald, A., Endres, M. and Neuhauser, H., 2015. Modifiable cardiovascular risk factors in adults aged 40–79 years in Germany with and without prior coronary heart disease or stroke. *BMC public health*, 15(1), p.701.
- Viadero D. Exercise seen as a priming pump for students' academic strides. *Education Week* <http://www.edweek.org>; 2008.
- World Health Organization, 2011. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity (No. WHO/NMH/NHD/MNM/11.1). World Health Organization.