

Full Length Research Paper

Effect of organic and inorganic fertilizer applications on phosphate solubilizing bacteria in the rhizosphere of maize (*Zea mays* L.)

Patrick Okorie Richard^{1*} and Adeniyi Adewale Ogunjobi

Department of Microbiology, University of Ibadan, Ibadan, Oyo State, Nigeria.

Received 30 September 2016, Accepted November 24, 2016

A field experiment was conducted between August and October, 2014 at the Department of Microbiology to assess the effect of organic and inorganic fertilizers on the population and activities of Phosphate Solubilizing Bacteria (PSB) in the rhizosphere of maize variety BR9928DMRSR-Y. Three treatment groups were used in the study and these were: Groups which received applications of organic fertilizer (Poultry litter) alone, groups which received applications of inorganic fertilizer (NPK) and the third group which was the control (CON) did not receive any fertilizer application. A total of twenty-three bacteria were isolated and *in-vitro* screening was done for different phosphate solubilization activity. The study revealed that maximum population of total heterotrophic bacteria (26.8×10^9 CFU/g) was obtained with organic fertilizer (OM) treatment. Among the different PSB isolates, OMPSB had the highest bacterial count of 10.2×10^9 . The lowest bacterial populations were obtained from IFPSB with 6.0×10^9 CFU/g. Out of the 23 PSB isolates, 18 were positive for phosphate solubilization with OMPSB8 showing the highest zone with 16 mm. Results showed that application of organic fertilizers enhanced the bacterial population and also showed increase in phosphate solubilization activities in rhizosphere soil compared to NPK and control treatments. This shows that organic fertilizers would be able to sustain the soil fertility for a longer period by meeting the demand of present and future generations.

Key words: Treatment groups, population, phosphate solubilization, *in-vitro* screening.

INTRODUCTION

Soil nutrient depletion has been a major challenge in Nigeria as a result of continuous cultivation of soils without adequate addition of external inputs. This decline may occur through leaching, soil erosion and crop harvesting (Muchena et al., 2005; Mutegi et al., 2008).

Soil nutrients face the risk of continuous decline unless the nutrients are replenished through the use of organic or mineral fertilizers, partially returned through crop residues or through traditional fallow systems. The continued soil nutrient depletion has a negative impact on

*Corresponding author. E-mail: patrickrichard.ok@gmail.com. Tel: 08063853332.

sustainable agriculture as crop production especially that of maize is decreasing (Hussain et al., 2007). Maize (*Zea mays* L.), an annual crop belonging to the grass family Poaceae (OECD, 2003; USDA, 2005), together with rice and wheat are the three most important cereal crops in the world (Min et al., 2015). Maize has the following advantages compared to other cereals: High production, easy processing, easy digestibility, and they are less expensive. It is an important source of carbohydrate, protein, iron, vitamin B and minerals (IITA, 2009) which according to IITA (2001) report, contains 80% carbohydrate, 10% protein, 3.5% fiber and 2% mineral. Thus, it is very essential for food and can also be used as forage and feed for livestock because it is among the highest in net energy content and lowest in protein and fibre content (Oladejo and Adetunji, 2012). In order to maximize the yield of maize crop, improved cultural practices such as organic and inorganic (chemical) fertilizers application can be used.

Though, application of chemical fertilizers can improve the nutrient balance of soils, which may lead to increases in crop yields, its continuous use is hazardous both to human health and the environment (Glick, 2003). This may cause plant toxicity (Nazar et al., 2012) and the bioaccumulation of trace metals in plants may pose a health risk consumed (Khan et al., 2015; Roy and McDonald, 2015). However, the negative effects of chemical fertilizers can be avoided by using organic fertilizers which have a positive effect on the Plant Growth Promoting Rhizobacteria (PGPR) - a group of rhizosphere bacteria found in association with roots which can enhance the growth of plant directly or indirectly by helping plants in nutrient uptake from rhizospheric soil (Mia et al., 2010). Organic fertilizers also play key role in sustaining soil fertility and crop productivity (Soumare et al., 2003; García-Orenes et al., 2016).

Furthermore, to increase the natural fertility of the soil and develop new approaches to reduce the need for chemical fertilizers, PGPR are recognized as important factors in sustainable agricultural production as they may be important for plant nutrition by increasing the P uptake by plants, playing a significant role in the bio-fertilization of crops (Idriss et al., 2007) and serving as a natural source of fertilizers that improve the efficiency of soil and plants (Khalid et al., 2004). The rhizosphere supports large and active microbial population such as bacteria, fungi, nematodes, protozoa, algae and microarthrops (Raaijmakers and Weller, 2001) which play important roles in ecological fitness of their plant host (Kent and Triplett, 2002).

Diverse PGPR strains have been used successfully for crop inoculations and to enhance plant growth (Kumar et al., 2012); these comprise members of the bacterial genera *Azospirillum* (Cassán and García, 2008), *Bacillus* (Jacobsen et al., 2004), *Pseudomonas* (Loper and Gross, 2007), *Rhizobium* (Long, 2001), *Serratia* (De

Vleeschauer and Höfte, 2007), *Stenotrophomonas* (Ryan et al., 2009) and *Streptomyces* (Schrey and Tarkka, 2008). *Pseudomonas* and *Bacillus* genera are the most commonly investigated PGPR, and often the dominating bacterial groups in the rhizosphere (Morgan et al., 2005). The modes of action of PGPR involve complex mechanisms to promote plant growth, development and protection through phytostimulation, bio-fertilization and biocontrol (Bloemberg and Lugtenberg, 2001). Some of these bacterial species have mineralization and solubilization potential for organic and inorganic phosphorus respectively (Hilda and Fraga, 2000; Khiari and Parent, 2005). Phosphate rock minerals are often too insoluble to provide sufficient P for crop uptake; therefore, the use of phosphate solubilizing bacteria (PSB) can increase crop yields up to 70% (Verma, 1993).

The aim of this work was to determine the effect of organic and inorganic fertilizer on the phosphate solubilizing bacteria present in maize plant rhizosphere. In a previous study by Azani (1995), chicken dung showed the best fertilizer compared to cow dung and NPK, thus, chicken dung was chosen as phosphorus source in this study.

MATERIALS AND METHODS

Experimental design

The experimental design used for the field experiment was the randomized complete block design as described by Kuntal et al. (2007). The experimental field was sited behind the Department of Microbiology, University of Ibadan, Ibadan, Oyo State. The field was divided into six blocks; two were treated with organic amendments (poultry litter), two with inorganic fertilizer (N:P:K 12:12:17), and two without any treatment and was considered as the control. Grains of maize variety BR9928DMRSR-Y were obtained from IITA, Ibadan headquarters and these were sown by planting three seeds/hole in each block. A total of ten holes were bore in each block.

Sampling and analysis

Sampling was done for a period of 56 days. Rhizosphere soils were collected at different growth stages of the maize (14, 28, 42 and 56 days after planting), by uprooting four plants from each treatment and keeping the soil around root system intact. After removing the bits of plant roots and other debris, the soil particles which adhered strongly to the roots was immediately used for analysis without drying following the method described by Basul et al. (2010). During the sampling period, population of Total Heterotrophic Bacteria (THB) and Phosphate Solubilizing Bacteria (PSB) in rhizosphere soil was counted at 14 days interval (that is, 14, 28, 42 and 56).

Isolation and enumeration of rhizosphere bacteria

Three replicate samples of rhizosphere soil were taken for enumeration of total heterotrophic count. The serial dilution technique as described by Johnson and Curl (1972) was employed to enumerate the rhizosphere soil by plating on nutrient agar using \

Table 1. Physico-chemical properties of soil.

Parameter	Value
pH (H ₂ O)	7.2
Phosphorus (g/kg)	8.06
Organic carbon (%)	38.29
Moisture content (%)	9.37

Table 2. Distribution of bacterial isolates from rhizosphere soil.

Isolates type	Treatments used	No. of isolates	Isolate codes
PSB	Organic manure	9	OMPSB1, OMPSB2, OMPSB3, OMPSB4, OMPSB5, OMPSB6, OMPSB7, OMPSB8, OMPSB9
	Inorganic fertilizer	8	IFPSB1, IFPSB2, IFPSB3, IFPSB4, IFPSB5, IFPSB6, IFPSB7, IFPSB8
	Control	6	CONPSB1, CONPSB2, CONPSB3, CONPSB4, CONPSB5, CONPSB6

OM, Organic manure; IF, inorganic fertilizer; CON, control; PSB, phosphate solubilizing bacteria.

the pour plate method. The plates were incubated at 37°C for 24 h. After the incubation period, the colony forming units were counted and expressed as colony forming unit per gram (CFU/g) of soil. Phosphate solubilizing bacteria were isolated by inoculating into sterile Petri dishes of Pikovskaya medium (Pikovskaya, 1948). The plates were incubated at 37°C for 48 h. After the incubation period, the colony forming units were counted and expressed as CFU/g of soil. Colonies with a 'halo' on the different media were sub-cultured onto nutrient agar plates.

The phosphate solubilizing bacteria were isolated using Pikovskaya medium (PVK) because it acts as specific isolation medium for PSB isolation due to the presence of calcium triphosphate which is known for 'halo' zone formation according to Sharma (2005). All the isolates were maintained at 4°C in equal volumes of nutrient broth and 30% glycerol.

Determination of soil physico-chemical properties

The following physico-chemical parameters of the soil were determined: pH, organic carbon, moisture content and phosphorus content. pH of the samples was read using an electronic digital pH meter (Table 1). Moisture content was determined by drying the samples in hot air oven at 105°C for 24 h using procedures outlined in the I.I.T.A. manual (IITA, 1979). The total organic carbon was measured by the method given by according to APHA (1985). Available P was determined by molybdenum blue method (Allen et al., 1974).

Biochemical tests

All the isolates were characterized by Gram staining and biochemical tests following the methods described by Olutiola et al. (1991). The various tests performed were oxidase, methyl red voges proskauer (MR-VP), indole, citrate, urease, H₂S production and fermentation of various sugars. The morphological and biochemical characterization of the PSB isolates obtained from the rhizosphere of maize was carried out with using 24 h old bacterial cultures. The results were compared with Bergey's Manual of Systematic Bacteriology (Holt et al., 1994).

In-vitro screening of isolates for phosphate solubilization activity

The isolates were screened for phosphate solubilization ability as described by Gupta et al. (1994). On modified Pikovskaya agar with insoluble tricalcium phosphate, a loop full of each culture was placed on the center of agar plates and incubated at 30°C for 5 days. The solubilization zone was determined by subtracting the diameter of bacterial colony from the diameter of total zone.

RESULTS AND DISCUSSION

Distribution of PSB Isolates

A total of 23 isolates were obtained from the soil samples during the 56 days of sampling as shown in Table 2. Twenty-three phosphate solubilizing bacterial isolates designated as PSB were isolated from the different treatments of organic manure (OM), inorganic fertilizer (IF) and non-amended soil which served as control (CON) using Pikovskaya medium. The distribution of these isolates is presented in Table 2. Those isolates considered as PSB were those that formed a 'halo' on Pikovskaya medium. These isolates were subjected to different phosphate solubilization activity screening.

Total heterotrophic bacteria (THB) counts

The total heterotrophic bacteria (THB) within the rhizosphere of maize plants was counted using Nutrient Agar by pour plate method. THB counts in the rhizosphere of amended soil (organic manure and inorganic fertilizer) and non-amended control soil are presented in Figure 1. The total heterotrophic bacteria population in rhizosphere soil amended with organic




Figure 1. Total heterotrophic bacteria count in rhizosphere soil.




Figure 2. The count of phosphate solubilizing bacteria in the rhizosphere of maize.

fertilizer increased from 15.4×10^9 CFU/g at day 14 to 26.8×10^9 CFU/g at day 56; that amended with inorganic fertilizer increased from 16.7×10^9 CFU/g to 25.2×10^9 CFU/g while that without amendment (Control) increased from 13.5×10^9 CFU/g to 20.1×10^9 CFU/g from day 14 to day 56, respectively. All through the period of study, the sample with organic manure treatment (OM) had the highest number of bacteria with 26.8×10^9 CFU/g followed by the inorganic fertilizer treatment (IF) having 25.2×10^9 CFU/g both at day 56. The inorganic fertilizer treatment had the highest bacterial count of 16.7×10^9 CFU/g at day 14. All the additives increased the microbial counts

(population) during the period of the study more significantly compared to the control that had a lower microbial population.

Enumeration of PSB

Phosphate solubilizing bacteria (PSB) within the rhizosphere of maize plant was counted using Pikovskaya medium. The bacteria count of PSB isolates during the period of study is presented in Figure 2. The phosphate solubilizing bacteria in rhizosphere soil amended with organic manure (OM) increased from

Table 3. Morphological, biochemical characterization and sugar fermentation of PSB isolates.

Isolates	Morphology	Gram's Reaction	Catalase	Oxidase	Indole	Citrate	Motility	Methyl Red	Voges Proskauer	H ₂ S Production	Starch Hydrolysis	Urease	Glucose	Galactose	Sucrose	Fructose	Lactose	Maltose	Mannitol	Probable organism
OMPSB1	R	+	+	-	-	-	-	-	+	-	+	+	+	-	-	+	-	-	+	<i>Bacillus</i> sp.
OMPSB2	R	-	+	+	-	+	+	-	-	-	-	+	+	-	+	+	+	-	+	<i>Pseudomonas</i> sp.
OMPSB3	R	-	+	+	-	+	-	-	-	-	-	+	+	+	+	+	-	-	-	<i>Pseudomonas</i> sp.
OMPSB4	R	+	+	-	-	-	+	+	+	-	+	+	+	-	-	+	-	-	+	<i>Bacillus</i> sp.
OMPSB5	C	+	+	+	-	+	+	-	-	-	+	-	+	-	+	+	-	+	-	<i>Micrococcus</i> sp.
OMPSB6	R	-	+	+	-	+	+	-	-	-	-	+	+	-	+	+	+	-	-	<i>Pseudomonas</i> sp.
OMPSB7	R	-	+	+	-	+	+	-	-	-	-	+	+	-	+	+	+	-	-	<i>Pseudomonas</i> sp.
OMPSB8	R	-	+	+	-	+	-	-	-	-	-	+	+	+	+	+	-	-	-	<i>Pseudomonas</i> sp.
OMPSB9	R	+	+	-	-	-	-	+	+	-	-	+	+	-	-	+	-	-	+	<i>Bacillus</i> sp.
IFPSB2	R	-	+	+	-	+	+	-	-	-	-	+	+	-	+	+	+	-	-	<i>Pseudomonas</i> sp.
IFPSB3	R	+	+	-	-	-	-	-	+	-	+	+	+	-	-	+	-	-	+	<i>Bacillus</i> sp.
IFPSB4	C	+	+	+	-	+	+	-	-	-	-	-	+	-	+	+	-	+	-	<i>Micrococcus</i> sp.
IFPSB5	R	-	+	+	-	+	+	-	+	-	+	+	+	-	-	+	-	-	-	<i>Pseudomonas</i> sp.
IFPSB6	R	-	+	+	-	+	-	-	-	-	+	+	+	-	-	+	-	-	-	<i>Pseudomonas</i> sp.
IFPSB7	R	-	+	+	-	+	-	-	-	-	-	+	+	-	-	+	-	-	-	<i>Pseudomonas</i> sp.
IFPSB8	R	+	+	-	-	-	-	-	+	-	+	+	+	-	-	+	-	-	+	<i>Bacillus</i> sp.
CONPSB2	R	-	+	+	-	+	+	-	-	-	-	+	+	-	+	+	+	+	+	<i>Pseudomonas</i> sp.
CONPSB3	R	+	+	-	-	-	-	-	+	-	+	+	+	-	-	+	-	-	+	<i>Bacillus</i> sp.
CONPSB4	R	-	+	+	-	+	+	-	-	-	-	+	+	-	+	+	+	-	-	<i>Pseudomonas</i> sp.
CONPSB5	R	-	+	+	-	+	+	-	-	-	-	+	+	-	+	+	+	-	+	<i>Pseudomonas</i> sp.
CONPSB6	R	-	+	+	-	+	+	-	-	-	-	+	+	-	+	+	+	-	-	<i>Pseudomonas</i> sp.

6.3×10⁹ CFU/g to 10.2×10⁹ CFU/g; that amended with inorganic fertilizer (IF) increased from 6.0×10⁹ CFU/g to 9.8×10⁹ CFU/g while that without amendment (Control) increased from 6.1×10⁹ CFU/g to 7.0×10⁹ CFU/g from day 14 to day 56, respectively. The organic manure treatment had the highest bacterial count with 10.2×10⁹ CFU/g followed by inorganic fertilizer treatment with 10.2×10⁹ CFU/g at day 56. Here too, the amended soil treatments had higher growth potential compared to the control soil treatment.

Identification of PSB isolates

The results of the morphological and biochemical tests are presented in Table 3. The isolates belong to *Pseudomonas* sp. (13), *Bacillus* sp. (6) and *Micrococcus* sp. (2).

Screening of isolates for PSB activity

A total of twenty three (23) PSB isolates were screened for phosphate solubilization on modified Pikovskaya agar, of which eighteen (18) isolates showed the development

of phosphate solubilization zones, ranging from 1.00 to 16.00 mm. Of the 18 isolates that showed zones of solubilization, 6 isolates (OMPSB5, CONPSB4, OMPSB8, IFPSB6, CONPSB5 had CONPSB6) had zones ranging between 6.00 and 16.00 mm while the other 12 isolates showed the development of zones less than 6.00 mm. OMPSB8 and IFPSB6 showed highest phosphate solubilization, that is, 16.00 and 15.00 mm, respectively. The zones of phosphate solubilization of PSB isolates are presented in Figure 3.

Table 4 presents the highest solubilization zones of the isolates from the different treatments during each sampling period (day 14, 28, 42 and 56). For day 14, OMPSB2 had the highest solubilization of 3.00 mm. At day 28, the highest solubilization zone was by OMPSB3 with 5.00 mm. OMPSB5 and CONPSB4 both had highest solubilization zones of 6.00 mm at day 42. However, OMPSB8 and IFPSB6 showed highest phosphate solubilization zone, that is, 16.00 and 15.00 mm respectively at day 56.

Plant growth-promoting rhizobacteria (PGPR) colonize roots of plant and promote plant growth and development through a variety of mechanisms. The exact mechanism by which PGPR stimulate plant growth is not clearly known, although several mechanisms such as production




Figure 3. The zones of phosphate solubilization of PSB isolates.

Table 4. Zones of phosphate solubilization of PSB isolates at different sampling periods.

Sampling periods	Isolates	Zones of solubilization (mm)
DAY14	OMPSB2	3.00
	IFPSB1	1.00
DAY28	OMPSB3	5.00
	IFPSB2	2.00
	CONPSB2	2.00
DAY42	OMPSB5	6.00
	IFPSB4	5.00
	CONPSB4	6.00
DAY56	OMPSB8	16.00
	IFPSB6	15.00
	CONPSB6	8.00

OM, Organic manure; IF, inorganic fertilizer; CON, control; PSB, phosphate solubilizing bacteria.

of phytohormones, suppression of deleterious organisms, activation of phosphate solubilization and promotion of the mineral nutrient uptake are usually believed to be involved in plant growth promotion (Bloemberg and Lugtenberg, 2001).

In the present study, 23 beneficial bacteria were isolated from maize rhizosphere. The bacteria isolates were screened for phosphate solubilization activities and characterized by biochemical tests. Generally, the total heterotrophic counts increased with increase in the duration of maize cultivation just as observed by Liang et

al. (2012) in their 15 years study on the effect of fertilizers on soil quality. This can be attributed to the fact that the soil is a suitable medium for the growth of environmental microorganisms because it is rich in nutrients. It has also been reported that the soil receiving manure has larger bacteria pool than in the same soil receiving only chemical fertilizers (Islam and Weil, 2002).

The treatments with organic and inorganic amendments recorded higher bacteria counts as compared to the control treatment. This is similar to the report of Zhao et al. (2014). This observation may be as a result of the

additional nutrients (N, P, K and micronutrients) provided by the additives which the bacteria breakdown for plant to use. This breakdown product in turn leads to the release of more exudates and plant products for use by the rhizosphere bacteria. Hence, increase in rhizosphere bacterial biomass (Das and Dkhar, 2011).

The addition of organic amendments increased the soil bacteria count compared to the inorganic fertilizer and control. Similar observations were made in organic recycling experiments by Chakrabarti et al. (2000) where soil receiving more organic matter tends to harbor higher levels of bacteria with higher microbial activity as proposed by Mäder et al. (2002). The higher bacteria count in OM treated soils is related to the organic matter with respect to decomposition in these materials which is important for proliferation of soil microorganisms in soil. The bacteria count for the organic manure treatments increased gradually over time and became higher than the inorganic fertilizer application which had a higher bacterial count at day 14. The explanation to this phenomenon can be deduced from the statement of Adegbidi et al. (2003) that the release of nutrients from composts and processed organic manures are generally slower than nutrient release from inorganic fertilizer.

Kumar et al. (2012) isolated and identified *Pseudomonas* spp. as the major PSB in their research. They also reported that *Pseudomonas* spp. are very efficient phosphate solubilizers because they dissolve the soil P through production of low molecular weight organic acids in addition to lowering the pH of rhizosphere. This corroborates the findings in this study where more than 50% of the PSB isolated were identified as *Pseudomonas* sp.

The PSB isolated from the rhizosphere soil with organic manure treatment OMPB8 (*Pseudomonas* sp.) showed the highest phosphate solubilization zone (16.00 mm) in PVK agar just as a documented by Lazcano et al. (2013). The reports of Zhang et al. (2012) and Yang et al. (2016) have also shown the positive effect of organic manure on the soil microflora. Furthermore, Kohler et al. (2015) showed that organic manure significantly increased shoot biomass by 64%. These reports and the current study show that organic manure application supports/enhances microbial activity in the rhizosphere of plants.

Conclusion

The application of organic and inorganic treatments significantly affected the rhizosphere bacterial population. Application of fertilizers showed increased bacterial population compared to the control treatment and this is of great importance in nutrient availability in the soil.

Similarly, the plant growth promoting activities of the isolates were also enhanced especially in the treatments with organic manure. The use of organic amendments is efficient, environmentally, cost effective and can be used in place of the more expensive inorganic fertilizer as a

viable alternative in the enrichment of nutrient deficient soil. Thus, it can be concluded that the application of organic manure can improve the soil health by improving the bacterial population. This type of study is necessary as it advocates that use of organic fertilizer is an efficient approach to replace chemical fertilizers.

Conflict of interest

The authors declare that there are no conflicts of interest

REFERENCES

- Adegbidi HG, Briggs RD, Volk TA, White EH, Abrahamson LP (2003). Effect of organic amendments and slow-release nitrogen fertilizer on willow biomass production and soil chemical characteristics. *Biomass Bioenerg.* 25:389-398.
- Allen SE, Grinshaw HM, Parkinson JA, Quaramby C (1974). *Chemical analysis of ecological materials.* Blackwell Scientific Publications, Oxford.
- APHA (1985). *Standard Method for the examination of water and waste water.* American Public Health Association, Washington, DC.
- Azani S (1995). Beberapa aspek fisiologi dan agronomi yang mempengaruhi pertumbuhan pucuk manis (*Sauropus androgynus* L.). Thesis M.Sc., Jabatan Botani Universiti, Malaysia, Kuala Lumpur.
- Basul M, Bhadoria PBS, Mahapatra SC (2010). Influence of Soil Ameliorants, Manures and Fertilizers on Bacterial Populations, Enzyme Activities, N Fixation and P Solubilization in Peanut Rhizosphere under Lateritic Soil. *Br. Microbiol. Res. J.* 1(1):11-25.
- Bloemberg GV, Lugtenberg BJJ (2001). Molecular basis of plant growth promotion and biocontrol by Rhizobacteria. *Curr. Opin. Plant Biol.* 4:343-350.
- Cassán F, García SI (2008). *Azospirillum* sp.: cell physiology, plant response, agronomic and environmental research in Argentina. Buenos Aires: Asociacion Argentina de Microbiologia.
- Chakrabarti K, Sarkar B, Chakrabarty A, Banik P, Bagchi DK (2000). Organic recycling for soil quality conservation in subtropical plateau region. *J. Agron. Crop. Sci.* 184:137-142.
- Das BB, Dkhar MS (2011). Rhizosphere Microbial Populations and Physico Chemical Properties as Affected by Organic and Inorganic Farming Practices. *Am. Eurasian J. Agric. Environ. Sci.* 10(2):140-150.
- De Vleeschauwer D, Höfte M (2007). Using *Serratia plymuthica* to control fungal pathogens of plants. *CAB Review.* 2:46.
- García-Orenes F, Roldán A, Morugán-Coronado A, Linares C, Cerdà A, Caravaca F (2016). Organic Fertilization in Traditional Mediterranean Grapevine Orchards Mediates Changes in Soil Microbial Community Structure and Enhances Soil Fertility. *Land Degrad. Dev.* 27:1622-1628.
- Glick BR (2003). Plant growth promoting bacteria, molecular biology principles and applications of recombinant DNA. *Microbiol. Res.* 2:436-54.
- Gupta RS, Rekha SA, Kuhad RC (1994). A modified plate assay for screening phosphate solubilizing microorganisms. *J. Gen. Appl. Microbiol.* 40:255-260.
- Hilda R, Fraga R (2000). Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol. Adv.* 17:319-359.
- Holt JG, Kreig NR, Sneath PHA, Staley JT, Williams ST (1994). *Bergey's Manual of Determinative Bacteriology.* 9th edition, Williams and Wilkins, (Editors). Baltimore, USA, Pp.40-169.
- Hussain I, Ahsan M, Saleem M, Ahmad A (2007). Gene action studies for agronomic traits in maize under normal and water stress conditions. *Pak. J. Agr. Sci.* 46:111-117.
- Idriss EE, Makarewicz O, Farouk A, Rosner K, Greiner RH, Bochow TR, Borriss R (2007). Extracellular phytase activity of *Bacillus amyloquelaceus* FZB45 contributes to plant growth promoting effect. *Microbiology.* 148:2097-2109

- ITA (1979). Selected Methods for Soil and Plant Analysis. Manual Series No.1, 3rd Edition. International Institute of Tropical Agriculture, Ibadan, p34.
- ITA (2001). International Institute of Tropical Agriculture, Ibadan, Oyo State. Annual Report on Maize Production.
- ITA (2009). International Institute of Tropical Agriculture. www.ita.org/cms/details/maize_project_details.aspx. Accessed at October 25, 2009.
- Islam KR, Weil RR (2002). Soil quality indicator properties in mid-atlantic soils as influenced by conservation management. *J. Soil Water Conserv.* 55:69-78.
- Jacobsen N, Nielsen PS, Jeffares DC, Eriksen J, Ohlsson H, Arctander P, Kauppinen S (2004). Direct isolation of poly(A)+ RNA from 4 M guanidine thiocyanate-lysed cell extracts using locked nucleic acid-oligo(T) capture. *Nucleic Acids Res.* 32(7):64.
- Johnson LF, Curl AE (1972). Method for the research on ecology of soil borne plant pathogens. Burgess Publishing Company, Minneapolis.
- Kent AD, Triplett EW (2002). Microbial communities and their interactions in soil and rhizosphere ecosystems. *Annu. Rev. Microbiol.* 56:211-236.
- Khalid A, Arshad M, Zahir MA (2004). Screening plant growth promoting rhizobacteria for improving growth and yield of wheat. *J. Appl. Microbiol.* 96:473-80.
- Khan A, Khan S, Khan MA, Qamar Z, Waqas M (2015). The uptake and bioaccumulation of heavy metals by food plants, their effects on plants nutrients, and associated health risk: a review. *Environ. Sci. Pollut. Res.* 22:13772-13799.
- Khiari L, Parent LE (2005). Phosphorus transformations in acid light-textured soils treated with dry swine manure. *Can. J. Soil Sci.* 85:75-87.
- Kohler J, Caravaca F, Azcón R, Díaz G, Roldán A (2015). The combination of compost addition and arbuscular mycorrhizal inoculation produced positive and synergistic effects on the phytomanagement of a semiarid mine tailing. *Sci. Total Environ.* (514):42-48.
- Kumar A, Kumar A, Devi S, Patil S, Payal S, Negi S (2012). Isolation, Screening and Characterization of Bacteria from Rhizospheric Soils for Different Plant Growth Promotion (PGP) Activities: an in vitro Study. *Recent Res. Sci. Technol.* 4(1):01-05.
- Kuntal D, Raman D, Thippenahalli NS, Nazim S (2007). Influence of biofertilizers on the biomass yield and nutrient content in *Stevia rebaudiana* Bert grown in India subtropics. *J. Med. Plants Res.* 1(1):005-008.
- Lazcano C, Gómez-Brandón M, Revilla P, Domínguez J (2013). Short-term effects of organic and inorganic fertilizers on soil microbial community structure and function. *Biol. Fertil. Soils.* 49:723.
- Liang Q, Chen H, Gong, Y, Fan M, Yang H, Lal R, Kuzyakov R (2012). Effects of 15 years of manure and inorganic fertilizers on soil organic carbon fractions in a wheat-maize system in the North China Plain. *Nutr. Cycl. Agroecosyst.* 92(1):21-33.
- Long SR (2001). Genes and signals in the Rhizobium-legume symbiosis. *Plant Physiol.* 125:69-72.
- Loper JE, Gross H (2007). Genomic analysis of antifungal metabolite production by *Pseudomonas fluorescens* Pf-5. *Eur. J. Plant Pathol.* 119:265-278.
- Mäder P, Fließbach A, Dubois D, Gu L, Padrou F, Niggli U (2002). Soil fertility and biodiversity in organic farming. *Science.* 296:1694-1697.
- Mia MA, Shamsuddin Z, Mahmood M (2010). Use of Plant Growth Promoting Bacteria in Banana: A New Insight for Sustainable Banana Production. *Int. J. Agr. Biol.* 12:459-467.
- Min XJ, Powell B, Braessler J, Meinken J, Yu F, Sablok G (2015). Genome-wide cataloging and analysis of alternatively spliced genes in cereal crops. *BMC Genomics.* 16:721.
- Morgan JA, Bending GD, White PJ (2005). Biological costs and benefits to plant-microbe interactions in the rhizosphere. *J. Exp. Bot.* 56:1729-1739.
- Muchena FN, Onduru D, Gachini GN, de Jager A (2005). Turning the tides of soil degradation in Africa: capturing the reality and exploring opportunities. *Land Use Policy.* 22:23-31.
- Mutegei JK, Mugendi D, Verchot L, Kung'u J (2008). Combining napier grass with leguminous shrubs in contour hedgerows controls soil erosion without competing with crops. *Agrofor. Syst.* 74(1):37-49.
- Nazar R, Iqbal N, Masood A, Khan M, Syeed S, Khan N (2012). Cadmium Toxicity in Plants and Role of Mineral Nutrients in Its Alleviation. *Am. J. Plant Sci.* 3(10):1476-1489.
- OECD (2003). Consensus document on the biology of Zea mays subsp. mays (Maize). Report No. 27, Environment Directorate; Organisation for Economic Co-operation and Development Paris, France.
- Oladejo JA, Adetunji MO (2012). Economic Analysis of Maize (*Zea mays* L.) Production in Oyo State of Nigeria. *Agr. Sci. Res. J.* 2(2):77-83.
- Olutiola PO, Famurewa O, Sontag HG (1991). An Introduction to General Microbiology. A Practical Approach. 1st Edn., Heidelberg Verlaganstalt and Druckerei GmbH Heidelberg, Germany.
- Pikovskaya RI (1948). Mobilization of phosphorus in soil in connection with vital activity of some microbial species. *Microbiologiya.* 17:362-370.
- Raaijmakers JM, Weller DM (2001). Exploiting genotypic diversity of 2,4-diacetylphloroglucinol-producing *Pseudomonas* spp.: characterization of superior root-colonizing *P. fluorescens* strain Q8r1-96. *pp. Environ. Microbiol.* 67:2545-2554.
- Roy M, McDonald LM (2015). Metal Uptake in Plants and Health Risk Assessments in Metal-Contaminated Smelter Soils. *Land Degrad. Dev.* 26:785-792.
- Ryan RP, Fischer AM, Kudela RM, Gower JFR, King SA, Marin III R, Chavez FP (2009). Versatility and adaptation of bacteria from the genus *Stenotrophomonas*. *Nature Rev. Microbiol.* 7:514-525.
- Schrey SD, Tarkka MT (2008). Friends and foes: streptomycetes as modulators of plant disease and symbiosis. *Antonie Van Leeuwenhoek.* 94:11-19.
- Sharma K (2005). Isolation, Purification and Identification of Bacteria. In: Manual of Microbiology. Ane Books Publication, New Delhi, p41.
- Soumare M, Tack FMG, Verloo MG (2003). Effects of a municipal solid waste compost and mineral fertilization on plant growth in two tropical agricultural soils of Mali. *Bioresour. Technol.* 86:15-20.
- USDA (2005). Germplasm Resources Information Network - (GRIN) [Online Database]. United States Department of Agriculture, Agricultural Research Service, Beltsville, available online at <http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl> 14-3-2005.
- Verma LN (1993). Biofertilizer in agriculture. In: P. K. Thampam (ed.) Organics in soil health and crop production. Peekay Tree Crops Development Foundation, Cochin, India. Pp.152-183.
- Yang H, Niu J, Tao J, Gu Y, Zhang C, She S, Chen W, Yang H, Yin H (2016). The Impacts of Different Green Manure on Soil Microbial Communities and Crop Health. Preprints.
- Zhang QC, Shamsi IH, Xu DT, Wang GH, Lin XY, Jilani G, Hussain N, Chaudhry AN (2012). Chemical fertilizer and organic manure inputs in soil exhibit a vice versa pattern of microbial community structure. *Appl. Soil Ecol.* 57:1-8.
- Zhao J, Ni T, Li Y, Xiong W, Ran W, Shen B, Shen Q, Zhang R (2014). Responses of Bacterial Communities in Arable Soils in a Rice-Wheat Cropping System to Different Fertilizer Regimes and Sampling Times. *PLoS ONE.* 9(1):e85301.