

*Full Length Research Paper*

# The effects of four weeks training on leptin levels in junior female judokas

Yücel OCAK

School of Physical Education and Sports, Afyon Kocatepe University, Afyonkarahisar-Turkey. E-mail: [yocak@aku.edu.tr](mailto:yocak@aku.edu.tr).

Accepted 30 July, 2011

The purpose of this study is to investigate the effects of four weeks judo training on plasma leptin levels. Twenty-five female national team athletes with a mean age of  $18.12 \pm 1.12$  years; body weight of  $62.72 \pm 22.72$  kg; body height of  $164.28 \pm 7.28$  cm, BMI of  $23.17 \pm 6.28$  kg/m<sup>2</sup>; training experience of  $6.56 \pm 5.44$  years participated in this study. The data were collected from junior female judokas who trained six times in a week for two hours daily. Physical measurements and blood samples were taken before and after four weeks of training including technical, tactical and physical preparations. Body fat percentages were measured by using the Tanita Bioimpedance BC-418 (Tokio, Japan). Statistical Package for Social Science 15.0 (SPSS) was used to analyse the obtained results. Before and after the training, paired samples t-test was used to evaluate the differences between serum leptin and body fat percentages. Results of this study showed that pre-training serum leptin levels were significantly lower than post-training levels ( $p < 0.01$ ). There was also significant decrease in body fat percentages after four week training ( $p < 0.01$ ). Four weeks judo training results in a decrease in total body fat and serum leptin level in junior female judokas. It may be considered that lowered serum leptin levels were due to the decrease of the body fat percentage after four weeks training.

**Key words:** Plasma leptin, body fat, female judokas.

## INTRODUCTION

Sport of judo needs dynamic and intermittent workloads with a high-intensity anaerobic activity (Degoutte et al., 2000). Judokas spend significantly high amount of energy during the training and competitions which affects energy balance, hormonal profile and metabolic hemostasis (Filaire et al., 2001). There was a limited number of studies related females judokas in the literature (Prouteau et al., 2007).

Leptin hormone has been a topic in several researches as it is associated with food intake and energy use directly. Regulation of the body weight and energy balance for human organism is one of the basic functions controlled by endocrine and central nervous system.

Leptin has been recently defined as an adipocyte derived hormone and regarded as a regulator of the body weight and energy balance (Simsch et al., 2002). There are several studies showing the effects of exercise on leptin levels. Many of these studies are related to why leptin responds to exercise and shows adaptation to exercise (Robert et al., 2002). Leptin is efficient in controlling the body weight in professional athletes and

exercise density and the amount of the energy spent are important in alternation in serum leptin level (Gomez-Merino et al., 2002). Exercise results in a decrease in fat mass and thereby effecting leptin levels. However, the exercise types requiring more calory and energy expenditure may result in more alterations in leptin levels (Perusse et al., 1997; Loucks et al., 1998). Decrease in plasma leptin levels associated with exercise may be balanced by nutrition (Koistinen et al., 1998). Low leptin levels in athletes occur together with the energy deficit caused by the exercise and associated decreased fat percentage (Casimiro-Lopes et al., 2009). But, low fat proportion may not always indicate low leptin level. This result can be seen in some studies (Koury et al., 2007). They showed that unaltered fat proportion values occurred in spite of low leptin levels after the training (Koury et al., 2007). It is thought that exercise makes its effects on leptin through sympathetic adrenergic system (Torjman, 2001). It was suggested that alterations occurring in the levels of growth hormone, cortisol, insulin, testosterone, epinefrin and norepinefrin during the

**Table 1.** Plasma leptin levels pre-training and post-training.

Plasma leptin levels		N	X	S	t	Sig
Leptin	Pre-training	25	4.70	3.48	5.307	0.000**
	Post-training	25	2.96	1.96		

\*\*There was significant difference at 0.01 level.

**Table 2.** Body fat levels pre-training and post-training.

Body fat levels		N	X	S	t	Sig
Body fat (%)	Pre-training	25	17.31	1.52	11.290	0.000**
	Post-training	25	15.48	1.60		

\*\*There was significant difference at 0.01 level.

exercise result in delayed decrease in leptin (Essing et al., 2000). In this respect, there are also studies in the literature showing that short-term exercise (<60 min.) has no effect on serum leptin levels (Weltman et al., 2000; Essing et al., 2000; Houmard et al., 2000; Olive and Milller, 2001; Sütken et al., 2006; Yamaner et al., 2010).

The aim of this study is to investigate the effects of four weeks training on plasma leptin levels and body fat percentages in junior female judokas.

## MATERIALS AND METHODS

### Subjects

The mean values of the subjects were 18.12±1.12 years for age, 62.72±22.72 kg for body weight, 164.28±7.28 cm for body height, 23.17±6.28 kg/m<sup>2</sup> for body mass index, and 6.56±5.44 years for training experience, respectively. Body mass index was calculated using following formula, BMI=body weight (kg)/body height<sup>2</sup> (m) (Fahey et al., 2007).

### Sample collections

The data were collected from junior female judokas who trained six times in a week for two hours daily for world championship. Physical measurements and blood samples were taken before and after four week training including technical, tactical and physical preparations. Body fat percentages were measured by the Tanita Bioimpedance BC-418 (Tokio Japan).

### Statistical analysis

Statistical Package for Social Science 15.0 (SPSS) was used to analyse the obtained results. Before and after four weeks training, paired samples t-test was used to evaluate the differences between serum leptin and body fat percentages.

## RESULTS

Serum leptin levels obtained after the training was

statistically significantly lower than plasma leptin levels before the training (p=0.000) (Table 1).

Body fat levels obtained after the training was statistically significantly lower than body fat levels before the training (p=0.000) (Table 2).

## DISCUSSION

Results of this study showed that plasma leptin and body fat percentages of judokas after judo training were significantly decreased. There are similarities with other studies reported in the literature (Landt et al., 1997; Leal-Cerro et al., 1998; Essig et al., 2000; Olive and Milller, 2001; Zaccaria et al., 2002; Yamaner et al., 2010). Burning the fat tissues through training and energy deficit associated with this case result in the same percentage of decrease in serum leptin levels (Witek et al., 2003). It was reported that long-term medium level exercises suppress the plasma leptin levels related to the decrease of body fat percentage presented by BMI. But, it was noted that serum leptin levels did not alter in short-term exercise (Sütken et al., 2006).

Some researchers reported that these results were also in disagreement with the some studies (Essing et al., 2000; Olive and Milller, 2001). They found that exercise had no acute effect on leptin levels and they also reported that serum leptin levels were decreased 30 to 40% after the exercise (Essing et al., 2000; Olive and Milller, 2001). In another study on acute effects of training, there was a decrease in fat mass, while there was not any alteration in serum leptin levels after the training (Perusse et al., 1997).

In a study made by Türkmen, judo and cycling exercises were found to have no influence on leptin, LDL, triglyceride, total cholesterol level (Türkmen, 2011). In a similar study, Kishali found out that 8 weeks of exercise does not change serum leptin levels and body fat percentages of male university students (Kishali, 2011).

Some studies were carried out on animals in order to

determine the effects of exercise on leptin levels (Pagano et al., 1999). They reported that no alteration was found in leptin levels of obese animals after the exercise while leptin levels of animals with lower body weight showed decrease (Pagano et al., 1999). Conversely there were also other studies in which serum leptin levels of different subjects having different fat mass show difference in terms of their responses to exercise (Hickey et al., 1996; Moller et al., 1998).

## Conclusions

It was found that judo training had a short term effect on plasma leptin level and body fat percentages. Following the four weeks training, there was a decrease in serum leptin levels and body fat percentages. The decrease in plasma leptin levels after training programme was associated with the decrease in body fat percentage. In other words plasma leptin levels may be altered depending on the body fat percentage.

## REFERENCES

- Casimiro-Lopes G, Oliveira AV, Portella ES, Lisboa PC., Donangelo CM, Moura EG, Koury JC (2009). Plasma leptin, plasma zinc, and plasma copper are associated in elite female and male judo athletes. *Biol. Trace Elem. Res.*, 127: 109- 115.
- Degoutte F, Jouanel EAN., Pandian MR, Elias AN, Pandian MR, Wang L, Suarez E, James N, Wilson AF (2000). Leptin and IGF-I levels in unconditioned male volunteers after short term exercise. *Psychoneuroendocrinol*, 25: 453-461.
- Essing DA, Alderson NL, Ferguson MA, Bartoli WP, Durstine JL (2000). Delayed effects of exercise on the plasma leptin concentration. *Metabolism*, 49(3): 395-399.
- Fahey DT, Insel MP, Roth TW (2007). *Fit and Well*, Seventh Edition, McGraw Hill, New York, USA.
- Filaire E, Maso F, Degoutte F, Jouanel P, Lac G. (2001). Food restriction, performance, psychological state and lipid values in judo athletes. *Int. J. Sports Med.*, 22: 454-459.
- Gomez-Merino D., Chennaoui M., Drogou C., Bonneau D., Guezzenec C.Y. (2002). Decrease in serum leptin after prolonged physical activity in men. *Med. Sci. Sports Exerc.*, 34(10): 1594-1599.
- Hickey MS, Considine RV, Israel RG, Mahar TL, McCammon MR, Tyndall GL, Houmard JA, Caro JF (1996). Leptin is related to body fat content in male distance runners. *Am. J. Phy.-Endoc. Met.*, 271(5): E938-E940.
- Hickey MS, Israel RG, Gardiner SN, Considine RV, McCammon MR, Tyndall GL, Houmard JA, Marks RH, Caro JF(1996). Gender differences in serum leptin levels in humans. *Biochem. Mol. Med.*, 59: 1-6.
- Houmard JA, Cox JH, MacLean PS, Barakat HA (2000). Effect of short-term exercise training on leptin and insulin action. *Metabolism*, 49(7): 858-861.
- Kishali NF (2011). Serum leptin level in healthy sedentary young men after a short-term exercise. *Afr. J. Pharm. Pharmacol.*, 5 (4): 522-526.
- Koistinen HA, Tuominen JA, Ebeling P, Heiman ML, Stephens TW, Koivisto VA (1998). The effect of exercise on leptin concentration in healthy men and in type 1 diabetic patients. *Med. Sci. Sports Exerc.*, 30: 805-810.
- Koury JC, Oliveira KJF, Casimiro-Lopes G, Oliveira AV, Moura EG, Donangelo CM (2007). Plasma zinc, copper, leptin and body composition are associated in elite female judo athletes. *Biol. Trace Elem. Res.*, 115: 23-30.
- Landt M, Lawson GM, Helgeson JM, Davila-Roman VG, Landenson JH, Jaffe AS, Hickner RC (1997). Prolonged exercise decreases serum leptin concentrations. *Metabolism*, 46: 1109-1112.
- Leal-Cerro A, Garcia-Luna PP, Astorga R (1998). Serum leptin levels in male marathon athletes before and after the marathon run. *J. Clin. Endocrinol. Metab.*, 83: 2376- 2379.
- Loucks AB, Verdun M, Heath EM (1998). Low energy availability, not stress of exercise, alters LH pulsatility in exercising women. *J. Appl. Phys.*, 84: 37- 46.
- Moller N, O'Brien P, Sreekumaran NK (1998). Disruption of the relationship between fat content and leptin levels with aging in humans. *J. Clin. Endocrinol. Metab.*, 83: 931-934.
- Olive JL, Miller GD (2001). Differential effects of maximal and moderate-intensity runs on plasma leptin in healthy trained subjects. *Nutrition*, 17(5): 365-369.
- Pagano C, Marzolo M, Granzotto M, Ricquier D, Federspil G, Vettor R (1999). Acute effects of exercise on circulating leptin in lean and genetically obese fa/fa rats. *Biochem. Biophys. Res. Com.*, 255: 698-702.
- Perusse L, Collier G, Gagnon J, Leon AS, Rao DC, Skinner JS, Wilmore JH, Nadeau A, Zimmet PZ, Bouchard C (1997). Acute and chronic effects of exercise on leptin levels in humans. *J. Appl. Phys.*, 83(1): 5-10.
- Prouteau S, Ducher G, Serbescu C, Benhamou L, (2007). Courteix. Gender differences in response to weight cycling in elite judoists. *Biol. Sport.*, 24(2): 91-104.
- Robert RK, Hongnan C, Daniel C (2002). Leptin and exercise. *Exp. Biol. Med.*, 227: 701-708.
- Simsch C, Lormes W, Petersen KG, Baur S, Liu Y, Hackney AC, Lehmann M, Steinacker JM (2002). Training intensity influences leptin and thyroid hormones in highly trained rowers. *Int. J. Sports Med.*, 23: 422-427.
- Sütken E, Balköse N, Özdemir F, Alataş Ö, Tunalı N, Çolak Ö, Uslu S, Öner S (2006). Effects of long term exercise on leptin levels in professional athletes. *J. Turk. Clin. Biochem.*, 4(3): 115-120.
- Torjman MC (2001). On the delayed effects of exercise on leptin: More questions than answers. *Nutrition*, 17(5): 420-422.
- Türkmen M (2011). A study on serum leptin, lipoproteins and glucose levels of judoists and cyclists of Turkey. *Afr. J. Pharm. Pharmacol.*, 5(4): 532-537.
- Yamaner F, Bayraktaroğlu T, Atmaca H, Ziyagil MA, Tamer K (2010). Serum leptin, lipoprotein levels, and glucose homeostasis, between national wrestlers and sedentary males. *Turk. J. Med. Sci.*, 40(3): 471-477.
- Weltman A, Pritzlaff CJ, Wideman L, Considine RV, Fryburg DA, Gutgesell ME, Hartman ML, Veldhuis JD (2000). Intensity of acute exercise does not affect serum leptin concentrations in young men. *Med. Sci. Sports Exerc.*, 32(9): 1556-1561.
- Witek K, Wit B, Lerezak K, Sempolska K, Glinkowski W (2003). Influence of a monitored aerobic training and a low energy diet on serum leptin concentration in obese women. *Biol. Sport.*, 20(1): 33-41.
- Zaccaria M, Ermolao A, Roi GS, Englaro P, Tegon G, Varnier M. (2002). Leptin reduction after endurance races differing in duration and energy expenditure. *Eur. J. Appl. Physiol.*, 87: 108-111.