

## Review

# Diversity of endophytic fungi in some tropical medicinal plants – A report

Kalyanaraman Rajagopal<sup>1\*</sup>, Maheswari S.<sup>2</sup> and Kathiravan G.<sup>1</sup>

<sup>1</sup>Department of Biotechnology, School of Life Sciences, Vels University, Pallavaram, Chennai 600117, India.

<sup>2</sup>Department of Botany, Vinayaka Missions University, Salem 603 308, Tamil Nadu, India.

Accepted 23 January, 2012

**Endophytic fungi are a taxonomically and ecologically heterogenous group of organisms; mainly belong to ascomycetes, coelomycetes and hypomycetes. The abundance of endophytic fungi in tropical regions has been studied. However, scant attentions have been given to medicinal plants of tropics. Synthesis of Indole acetic acid and metabolites showing antiviral, volatile metabolites, anticancer agents, anti-oxidants, insecticidal activities, antidiabetic and immunosuppressive compounds are widespread among the endophytic fungi. Now, more concentration has been given to the chemistry and bioactive compounds of endophytic fungi, mycodiversity and their ecological role. This review deals with the brief survey of the endophytic fungi of some medicinal hosts and their bioactive compounds characterized.**

**Key words:** Ascomycetes, coelomycetes, ecology, hypomycetes, secondary metabolites, and tropical.

## INTRODUCTION

The term endophyte was first coined by De Bary in 1866 (Wilson, 1995). Since then it has undergone many subtle changes.

### Endophyte: Circumscription

The term endophyte itself evolved to accommodate our increasing knowledge regarding this group of fungi. Ainsworth (1971) defined endophyte as “a plant living inside another organism”. Carroll (1986) confined the use of the term to organisms that cause asymptomatic infections within plant tissue; this definition excluded pathogenic fungi. Petrini (1991) widened the scope of this definition to include all organisms inhabiting plant organs that, at some time in their life, can colonize internal plant tissues without causing apparent harm to the host. This circumscription would include latent pathogens and epiphytes that are found as endophytes during some stages of their life. Thus, some endophytes residing in plant tissues become pathogenic when their hosts are

weakened (Rajagopal, 1998). Wilson (1995) proposed that endophytes are fungi or bacteria that, for all or part of their life cycle, invade the tissues of living plants and cause unapparent and asymptomatic infections entirely within plant tissues but cause no symptoms of disease. The fossil record indicates that plants have been associated with endophytic (Krings et al., 2007) and mycorrhizal (Redecker et al., 2000) fungi for > 400 million years. Almost all the plant species (~400,000) harbor one or more endophytic organisms (Tan and Zou, 2001). Dreyfuss and Chapela (1994) estimated that endophytic fungi from the 270,000 species of plants existing on this planet could account for  $1.38 \times 10^6$  unique fungal sp. The enormity of this estimation is because endophytic fungi have been isolated from every plant species examined to date and endophytes are important components of fungal diversity (Arnold et al., 2000; Kumaresan and Suryanarayanan, 2002; Suryanarayanan and Rajagopal, 1998).

The presence of endophytic fungi in plant tissues was discovered more than 75 years ago when Sampson (1935) reported such fungi from *Lolium* grass. The contemporary resurgence of research on endophytic fungi began when Bernstein and Carroll (1977) reported the presence of endophytes in needles of *Pseudotsuga*

\*Corresponding author. E-mail: ammaaappa@gmail.com.

*menziesii*. The study of endophytic fungi apart from shedding light on diversity of fungal kingdom, offers a promising digression since some endophytes produce novel metabolites of pharmaceutical and agricultural value (Rajagopal et al., 2010).

### **Biodiversity and ecology of endophytic fungi of medicinal plants**

Fungal endophytes are ubiquitous. Almost all vascular plant species examined to date were found to harbor endophytic bacteria and or fungi (Sturz et al., 2000; Arnold et al., 2000). They have been isolated from such diverse groups of plants as marine algae (Cubit, 1974), mosses and ferns (Petrini, 1986), coniferous trees (Carroll and Carroll, 1978; Johnson and Whitney, 1992), angiosperms such as *Dalbergia latifolia*, *Azadirachta indica*, Grasses (Clay, 1990; Suryanarayanan and Rajagopal, 2000).

Medicinal plants are reported to harbor endophytes (Strobel, 2002), which in turn provide protection to their host from infectious agents and they are unexpected producers of metabolites useful to pharmaceutical and agricultural industries (Petrini et al., 1992b). Research on species composition of endophyte communities of many hosts has shown that a large number of fungal taxa could be isolated from a single host species (Petrini et al., 1989). However, usually one or a few fungal species dominate in a host plant (Rajagopal et al., 2010; Suryanarayanan et al., 1998). The distribution of endophytes in a host is influenced by environmental factors and age of the host species or tissue. The number of endophytic species that can be recovered from host decreases with extreme climates (Suryanarayanan et al., 1998). Several studies have shown that the frequency of colonization increases with age of the tissue colonized (Espinosa-Gracia and Langenheim, 1990; Rajagopal, 1998). Similarly, the endophyte community of a plant is affected by the quality of air (Petrini, 1991) and leaf chemistry (Rajagopal et al., 2010). The endophytes occupy a unique niche. Though they may not be exposed to the vagaries of environment as the phylloplane fungi are, they have to encounter the defense reactions of the host. Hence, their life strategies are likely to be different from those of other fungi. Some of the medicinal host for endophytic fungi, host plant and chemical compound reported tabulated in (Tables 1, 2 and 3).

### **Physiology of endophytes**

The production of extracellular enzymes such as cellulases, pectinases, esterases, amylases, proteases etc., by some endophytic fungi has been studied by substrate utilization test and isozyme analysis (Sieber et al., 1991; Rajagopal, 1998). Production of the growth

hormone indole acetic acid has been demonstrated for fungal endophytes such as *Aureobasidium pullulans* and *Epicoccum purpurascens* (Pugh et al., 1971, 1972) *A. indica* (Rajagopal, 1998). Carroll and Petrini (1983) investigated the ability of endophytes to tolerate or metabolize phenolics and other defense chemicals of the host tissues. Suryanarayanan and Rajagopal (1998) reported that endophytic fungi are known to produce compounds that interfere with plant cell division.

Endophytes have recently been shown to be key elements in plant symbiosis, affecting host tolerance in stressful conditions (Redman et al., 2002; Rodriguez et al., 2004; Marquez et al., 2007) plant defense (Omacini et al., 2001; Bailey et al., 2006) plant growth (Ernst et al., 2003) plant community biodiversity (Clay and Holah, 1999) melanin synthesis (Rajagopal et al., 2011). However the diversity, geographic distribution and host specificity of endophytes remain largely unknown (Arnold et al., 2001, Ganley et al., 2006, Otero et al., 2002, 2007, Arnold and Lutzoni, 2007, Higgins et al., 2007). Recent estimation of fungal diversity (Hawksworth, 1991, 2001; Schmit and Mueller, 2007) suggest that more than 90% of fungal species are not described and there are no specific estimates of the number of existing endophytes (Hyde and Soyong, 2007, Hyde et al., 2007).

### **Biotechnological prospective of endophytic fungi of medicinal plants**

Endophytic fungi are one of the major potential sources for new, useful metabolites (Dreyfuss and Chapela, 1994). There has been a great interest in endophytic fungi as potential producers of novel, biologically active products (Schulz et al., 2002; Strobel and Daisy, 2003; Tomita, 2003; Urairaj et al., 2003; Wildman, 2003).

Tan and Zou (2001) reported that endophytic fungi are rich source of functional metabolites which include alkaloids, amines, amides, indole derivatives, pyrrolizidines, steroids, terpenoids, sesquiterpenes, diterpenes, isocoumarin derivatives, quinines, flavonoids, phenyl propanoids, lignans, peptides, phenol, phenolicacids, aliphatic compounds, chlorinated metabolites etc.

Endophytic fungi colonize very special and often very hostile habitats and were increasingly recognized as a group of organisms that are likely to be sources of new metabolites useful in biotechnology and agriculture (Bills and Polishook, 1992). Many endophytic fungi elaborate antibiotic compounds in culture that are active against human and plant pathogens. Fisher et al. (1984a) showed that more than 30% of the endophytic fungi that they tested possess antifungal and antibacterial activity. Fisher et al. (1984b) have isolated a broad-spectrum antibiotic from an endophyte of *Vaccinium* sp. Endophytes such as *A. pullulans* and *E. purpurascens* influenced seed germination by producing phyto-

**Table 1.** Host plants surveyed.

S/N	Host plants	S. No	Host plants	S. No	Host plants
1.	<i>Abies spectabilis</i>	13.	<i>Collicarpa tomentosa</i>	25.	<i>Podocarpus netifolius</i>
2.	<i>Adhatoda zeylanica</i>	14.	<i>Crataeva magna</i>	26.	<i>Saussurea involucrate</i>
3.	<i>Aegle marmelos</i>	15.	<i>Eucalyptus globules</i>	27.	<i>Stipa grandis</i>
4.	<i>Azadirachta indica</i>	16.	<i>Garcinia mongostana</i>	28.	<i>Taxus cuspidate</i>
5.	<i>Bauhinia phoenicea</i>	17.	<i>Garcinia pravifolia</i>	29.	<i>Taxus muirei</i>
6.	<i>Butea monosperma</i>	18.	<i>Holarrhena antidysenterica</i>	30.	<i>Taxus wallichiana</i>
7.	<i>Calotropis procera</i>	19.	<i>Leucas aspera</i>	31.	<i>Terminalia arjuna</i>
8.	<i>Camptotheca acuminata</i>	20.	<i>Lobelia nicolinifolia</i>	32.	<i>Terminalia chebuba</i>
9.	<i>Catharanthus roseus</i>	21.	<i>Madhuka nerifolia</i>	33.	<i>Tridax procumbens</i>
10.	<i>Cedrus deodara</i>	22.	<i>Naregamia alata</i>	34.	<i>Tsuga dumosa</i>
11.	<i>Clerodendron serrature</i>	23.	<i>Ocimum basilicum</i>	35.	<i>Vitex negundo</i>
12.	<i>Coleus aromatics</i>	24.	<i>Ocimum sanctum</i>	36.	<i>Withania somnifera</i>

**Table 2.** Endophytic fungi genera reported.

S/N	Endophytic fungi	S/N	Endophytic fungi	S/N	Endophytic fungi
1.	<i>Chaetomium</i>	13.	<i>Phomopsis</i>	25.	<i>Fusarium</i>
2.	<i>Glomerella</i>	14.	<i>Phyllosticta</i>	26.	<i>Myrothecium</i>
3.	<i>Guignardia</i>	15.	<i>Sphaeropsis</i>	27.	<i>Nigrospora</i>
4.	<i>Pyrenospora</i>	16.	<i>Acremonium</i>	28.	<i>Paecilomyces</i>
5.	<i>Xylaria</i>	17.	<i>Alternaria</i>	29.	<i>Penicillium</i>
6.	<i>Botryosphaeria</i>	18.	<i>Aspergillus</i>	30.	<i>Pithomyces</i>
7.	<i>Colletotrichum</i>	19.	<i>Aurobasidium</i>	31.	<i>Rhizoctonia</i>
8.	<i>Cytospora</i>	20.	<i>Bipolaris</i>	32.	<i>Sporidesmium</i>
9.	<i>Monochaetia</i>	21.	<i>Cladosporium</i>	33.	<i>Stenella</i>
10.	<i>Pestalotia</i>	22.	<i>Curvularia</i>	34.	<i>Trichoderma</i>
11.	<i>Pestalotiopsis</i>	23.	<i>Cylindrocarpon</i>	35.	<i>Verticillium</i>
12.	<i>Phoma</i>	24.	<i>Drechslera</i>	36.	<i>Mucor</i>

hormones (Petrini et al., 1992a). In developed countries, fungal endophytes are viewed, as biocontrol agents for fungal plant diseases (Clay, 1989; Petrini, 1991). They are readily selectable component of the fungus-plant system and are ideal for experimental manipulations. Endophyte represents natural genomes vested with useful attributes that could be selected and introduced into host plants for desirable traits.

Alternatively, endophytes could be used as vectors of genes to be introduced artificially into a population of host plants (Petrini et al., 1992a).

Endophyte association offer greatest potential for biocontrol programmes because these fungi are integrated into host systems (Clay, 1989). Dewan and Sivasithamparam (1989) reported that a fungal endophyte isolated from wheat provides significant protection to the host from infection by "take all" fungus. Endophytic fungi of conifer needles produce chemicals that ward off insect pests. The best example for endophyte-mediated antagonism towards an insect pest is that of Douglas fir. The needles of this plant harbour

*Rhabdocline parkeri* as an endophyte, and this fungus controls the gall midge, *Contarina* sp. (Carroll and Carroll, 1978). Thus, some endophytes are potential candidates for biocontrol programme (Dorworth and Callan, 1996). Host specific endophytic fungi could be used as a vector for introducing foreign genes into the host. Endophytic fungi could be genetically manipulated to produce novel compounds in a host- a kind of indirect genetic engineering of plants (Clay, 1989).

Endophytes are being explored by both pharmaceutical and agricultural industries (Monaghan et al., 1995) as they represent an untapped pool of secondary metabolites (Dreyfuss and Chapela, 1994). Merck Research Laboratories, USA, is collaborating with Costa Rica in INBIO project to screen fungal endophyte from forest trees for novel antibiotics (Bills and Polishook, 1994; Sanders et al., 1995). Recently Li et al. (1998) reported that an endophyte occurring in the bark of *Taxus wallichiana* growing in Nepal Himalayas produces taxol, an anticancer drug. Souvik et al. (2008) reported that an endophyte from *Hypericum perforatum* produced

**Table 3.** Compounds extracted from endophytic fungi.

S/N	Chemical compounds reported	S/N	Chemical compounds reported
1.	Taxol	38.	Cyclohexane, 4, methyl
2.	Asperfumin	39.	Decane 3, 3, 6-trimethyl
3.	Cryptocin,	40.	Undecane, 4, 4-demethyl
4.	Cryptocandin,	41.	Alkanes including those of pentyl, hexyl, heptyl, octyl, sec-octyl and decyl alcohols
5.	Jesterone	42.	Decane, 3, 3, 5-trimethyl
6.	Oocydin	43.	Chaetomugilin A(2)
7.	Isopestacin	44.	Chaetoglobosins A(3) Chaetoglobosins C (4) Polyhydroxylated C <sub>29</sub>
8.	Pseudomycins	45.	Sterol, 25 $\epsilon$ - methyl 22-homo-5 $\alpha$ -cholest- 7
9.	Ambuic acid	46.	22-diene-3 $\beta$ , 6 $\beta$ , 9 $\alpha$ - triol
10.	Ergovaline	47.	Designated globosterol (1) Tetrahydroxylated ergosterol 22E, 24R-ergosta-7,
11.	Peramine	48.	22-diene-3 $\beta$ , 5 $\alpha$ , 6 $\beta$ - 9 $\alpha$ -tetrol (2)
12.	2-methoxy-4-hydroxy-6-methoxymethyl-benzaldehyde	49.	7-hydroxyphthalide [4]
13.	Cryptosporiopsin and 4-epi-ethiosolide	50.	4- hydroxyphthalide 5
14.	loline (1-aminopyrrolizidines)	51.	5-methoxy-7- hydroxyphthalide[6]
15.	Fumonisia	52.	5,7, dihydroxyphthalide [7] (3R,4R)-cis-4-hydroxymellein [8]
16.	6-isoprenylindole-3-carboxylic acid	53.	(3R,4R)-cis-4-hydroxy-5-methylmellein [9]
17.	3 $\beta$ , 5 $\alpha$ -dihydroxy-6 $\beta$ -acetoxo-ergosta 7, 22-diene	54.	Ergosterol [10]
18.	Cytosporones D and E	55.	5 $\alpha$ , 8 $\alpha$ - epidioxyergosterol [11]propanoic acid
19.	Trihydroxybenzene laotones	56.	2-methyl-methyl ester
20.	Phomoxanthone A and B	57.	2-methyl- 1-butanol
21.	(-)-(2S,5Z)-2-(1-methylethyl)-4-oxo-5-(phenylmethylene)-1	58.	3-methyl 1-butanol
22.	3-dioxalane-2 carboxylic acid	59.	3-methyl, acetate
23.	Munumbicins A, B, C and D,	60.	2-methyl- 2-methylbutyl ester Ethanol
24.	Bicyclic fusidilactones A (1) and B(2)	61.	Diethylphthalate
25.	Fusidilactone C (4)	62.	Ethyl alcohol
26.	Taxoids	63.	Ethyl ether
27.	Naphthalene 1, 1'- oxybis	64.	Tricyclic sesquiterpenoids Brasilamides A-D
28.	(2'S)-2-(Propan 2'-01)-5-hydroxy-benzopyran-4-one(2)	65.	Pinthunamide
29.	2, 3-dihydro-2 methyl-benzopyran-4,5-diol (4):	66.	Brasilamide (A-D)
30.	Benzopyrans; 2-methyl-5-methoxy-benzopyrone-4-one(1)	67.	Diketopiperazine (DKP) Colletopiperazine,
31.	(2R) -2, 3-dihydro-2-methyl 5-methoxy-benzopyran-4-one (3)	68.	Fusapepazine C and E
32.	Hypericin	69.	Sesquiterpenoids
33.	Naphthoquinone spiroketals	70.	Polyketide
34.	Palmarumycin	71.	Eremophilane
35.	Preussomerin	72.	Mairetolide F (6)
36.	Undecane-2,6-dimethyl	73.	Eremophilanolides $\alpha$ -methylene- $\gamma$ -lactone (1-3)

Table 3. Contd.

37.	Decane-3,3,5-trimethyl	74.	Butanedioic acid
-----	------------------------	-----	------------------

Hypericin (Tables 1, 2 and 3).

Endophyte represents natural genomes vested with useful attributes that could be selected and introduced into host plants for desirable traits. Alternatively, endophytes could be used as vectors of genes to be introduced artificially into a population of host plants (Petrini et al., 1992a).

### Conclusion

Endophytic fungi of medicinally important host are poorly investigated group of microorganisms that represent untapped pool of bioactive and novel chemical compounds which can be exploited in a variety of agriculture, pharmaceutical and chemical industries in near future. Some species of endophytic fungi have been identified as source of anticancer, antidiabetic, insecticidal and immunosuppressive compounds are widespread among endophytic fungi. All aspects of the biology and interrelation of endophytic with the respective hosts are largely under investigation. More information about host plant and their internal microbiology would be exceedingly useful in developing the search for novel bioactive compounds. A great deal of uncertainty also exists between what the endophyte produces in culture and what it may produce in nature. This may be especially true if the bioactive product of the endophyte is unique to it and the host does not produce it. Even though studies on isolation, characterization and extraction bioactive compounds from endophytic fungi from medicinal plants have just started and use of this enormous mycoresource is poorly understood. More discoveries of endophytes and their products from

this field hold exciting promise. This is amply supported by the identification of wide variety of endophytic fungi and products reported so far. To get more benefit from these groups of organisms natural forests must be protected in countries like India, that have different forest types. Hence, more studies on these groups of organisms are required. Joint effort is needed between chemists and mycologists to understand fungal biology, ecology and the mycologist will have the opportunity to gain more insight into the multifarious diversity of the fungal kingdom. The chemists could discover new chemical compounds including proteins or enzymes.

### REFERENCES

- Ainsworth GC (1971). Ainsworth and Bisby's Dictionary of the fungi. Commonwealth Mycol. Inst. Kew: CAB International, pp. 663.
- Arnold AE, Maynard Z, Gilbert GS, Coley PD, Kursar TA (2000). Are tropical fungal endophytes hyperdiverse? *Ecol. Lett.*, 3: 276-274.
- Arnold AE, Maynard Z, Gilbert GS (2001). Fungal endophyte in dicotyledonous neotropical trees: Patterns of abundance and diversity. *Mycol. Res.*, 105: 1502-1507.
- Arnold AE, Lutzoni E (2007). Diversity and host range of foliar fungal endophytes. Are tropical leaves biodiversity hotspots? *Ecology*, 88: 541-549.
- Bailey BA, Bae H, Strem MS, Roberts DP, Thomas SE, Crozier J, Samuels GJ, Choi I, Holmes KA (2006). Fungal and gene expression during the colonization of *Cacao* seedlings by endophytic isolates of four *Trichoderma* species. *Planta*, 224: 1449-1464.
- Bernstein ME, Carroll GC (1977). Internal fungi in old-growth Douglas fir foliage. *Can. J. Bot.*, 55: 644-653.
- Bills GF, Polishook JD (1992). A new species of *Mycocleptodiscus* from living foliage of *Chamaecyparis thyoides*. *Mycotaxon*, 43: 453-460.
- Bills GF, Polishook JD (1994). Abundance and diversity of microfungi in leaf litter of a lowland rain forest in Costa Rica. *Mycologia*, 86: 187-198.
- Carroll GE (1986). The biology of the endophytism in plants with particular reference to woody perennials. In Fokkema NJ, Van den Heuvel L (eds) *The Microbiology of the Phyllosphere*. Cambridge University Press, Cambridge, pp. 205-222.
- Carroll GC, Carroll FE (1978). Studies on incidence of coniferous needle endophyte in the Pacific Northwest. *Can. J. Bot.*, 56: 3034-3043.
- Carroll GC, Petrini O (1983). Patterns of substrate utilization by some endophytes from coniferous foliage. *Mycologia*, 75: 53-63.
- Clay K (1989). Clavicipitaceous endophytes of grasses: their potential as biocontrol agents. *Mycol. Res.*, 92(1): 1-12.
- Clay K (1990). Fungal endophytes of grasses. *Annu. Rev. Ecol. Syst.*, 21: 275-297.8
- Clay K, Holah J (1999). Fungal endophyte symbiosis and plant diversity in successional field. *Science*, 285: 1742-1744.
- Cubit (1974). Interaction of seasonally changing physical factors and grazing affecting high intertidal communities on a rocky stone. PhD dissertation, Univ Oregon, Eugene, Oregon.
- De Bary A (1866). *Morphologie und Physiologie der Pilze, Flechten und Myxomyceten*. Hofmeister's Handbook *Physiol. Bot. Leipzig*, p. 2.
- Dewan MM, Sivasithamparam K (1989). Behaviour of a plant growth promoting sterile fungus on agar and roots of ryegrass and wheat. *Mycol. Res.*, 93: 161-166.
- Dorworth CE, Callan BE (1996). Manipulation of endophytic fungi to promote their utility as vegetation biocontrol agents. In Redlin SC, Carris LM (eds) *Endophytic Fungi in Grasses and Woody Plants-Systematics, Ecology and Evolution*, APS Press, St Paul, MN, USA, pp. 209-218.
- Dreyfuss MM, Chapela IH (1994). Potential of fungi in discovery of novel low molecular weight pharmaceuticals. In Gullo VP (eds) *The discovery of Natural Products with Therapeutic Potential*, Butterworth-Heinemann, London, UK, pp. 49-80.
- Ernst M, Mendgen KW, Wirsal SGR (2003). Endophytic fungal mutualists: Seed borne *Stagnospora* sp. enhances seed biomass production in axenic microcosms. *Mol. Plant. Microbe Interact.* 16: 580-587.
- Espinosa-Gracia FL, Langenheim JH (1990). The endophytic fungal community in leaves of a coastal redwood population-diversity and spatial patterns. *New Phytol.*, 116:

- 89-97.
- Fisher, Anson AE, Petrini O (1984a). Antibiotic activity of some endophytic fungi from ericaceous plants. *Bot. Helv.*, 94: 249-253.
- Fisher, Anson AE, Petrini O (1984b). Novel antibiotic activity of some endophytic *Cryptosporiopsis* sp. isolated from *Vaccinium myrtillus*. *Trans. Brit. Mycol. Soc.*, 83: 145-148.
- Ganley RJ, Newcombe G (2006). Fungal endophytes in seeds and needles of *Pinus monticola*. *Mycol. Res.*, 100: 318-332.
- Hawksworth DL (1991). The fungal dimension of biodiversity. Magnitude, Significance and Conservation. *Mycol. Res.* 95: 641-655.
- Hawksworth DL (2001). The magnitude of fungal diversity. The 1.5 million species estimate revisited. *Mycol. Res.*, 105: 1422-1432.
- Higgins KL, Arnold AE, Miadlikowska J, Sarvate SD, Lutzoni F (2007). Phylogenetic relationship, host affinity and geographic structure of boreal and arctic endophytes from three major plant lineages. *Mol. Phylogenet. Evol.*, 42: 543-555.
- Hyde KD, Bussaban B, Paulus PW, Crous S, Lee McKenzie EHC, Photita W, Lumyong S (2007). Diversity of saprobic microfungi. *Biodivers. Conserv.*, 16: 7-35.
- Hyde KD, Soyong K (2007). Understanding microfungi diversity—a critique. *Cryptogamie Mycol.*, 28: 281-289.
- Johnson JA, Whitney NJ (1992). Isolation of fungal endophytes from black spruce (*Picea mariana*) dormant buds and needles from New Brunswick, Canada. *Can. J. Bot.*, 70: 1754-1757.-1
- Krings M, Taylor TN, Hass H, Kerp H, Dotzler N, Hermsen EJ (2007). Fungal endophytes in a 400 million year old land plants: Infection pathways, spatial distribution, and host responses. *New Phytol.*, 174: 648-657.
- Kumaresan V, Suryanarayanan TS (2002). Endophytes assemblages in young mature and senescent leaves of *Rhizophora apiculata* evidence for the role of endophytes in mangrove litter degradation. *Fungal Divers.* 9: 81-91.
- Li JY, Sidhu RS, Ford EJ, Long DM, Hess WM, Strobel GA (1998). The induction of taxol production in the endophytic fungus *Periconia* sp. from *Torreya grandifolia*. *J. Ind. Microbiol. Biotechnol.*, 20(5): 259-264.
- Marquez LM, Redman RS, Rodriguez RJ, Roossinck MJ (2007). A virus in a fungus in a plant: Three way symbioses required for thermal tolerance. *Science*, 315: 513-517.
- Monaghan RL, Polishook JD, Pecore VJ, Bills GF, Nallin-Omstead M, Streicher SL (1995). Discovery of novel secondary metabolites from fungi—is it really a random walk through a random forest? *Can. J. Bot.*, 73: 925-931.
- Omacini M, Chaneton EJ, Ghersa CM, Mueller CB (2001). Symbiotic fungal endophytes control insect host-parasite interaction webs: *Nature*, 409: 78-81.
- Otero JT, Ackerman JD, Bayman P (2002). Diversity and host specificity of endophytic *Rhizoctonia*-like fungi from tropical orchids. *Am. J. Bot.*, 89: 1852-1858.
- Otero JT, Flanagan NS, Herre EA, Ackerman JD, Bayman P (2007). Widespread mycorrhizal specificity correlates to mycorrhizal function in the neotropical, epiphytic orchid *Lonopsis utricularioides* (Orchidaceae). *Am. J. Bot.*, 94: 1944-1950.
- Petrini P (1986). Taxonomy of endophytic fungi of aerial plant tissues. In: Fokkema NJ, Van Den Heuvel J (eds). *Microbiology of the phytosphere*. Cambridge University Press, Cambridge, pp. 175-187.
- Petrini O (1991). Fungal endophytes of tree leaves. In Andrew IA, Hirano SS (eds) *Microbial Ecology of Leaves*, Springer-Verlag, New York, pp 179-197.
- Petrini LE, Petrini O, Laflamme G (1989). Recovery of endophytes of *Abies balsamea* from needles and galls of *Paradiplosis tumifex*. *Phytoprotection*, 70: 97-103.
- Petrini O, Fisher PJ, Petrini LE (1992a). Fungal Endophytes of bracken (*Pteridium aquilinum*) with some reflection on their use in biological control. *Sydowia*, 44: 282-293.
- Petrini O, Sieber TN, Toti L, Viret O (1992b). Ecology, Metabolite production and Substrate utilization in endophytic fungi. *Natl. Toxin.*, 1: 185-196.
- Pugh GJF, Buckley NG (1971). *Aureobasidium pullulans*: an endophyte in sycamore and other trees. *Trans. Br. Mycol. Soc.*, 57: 227-231.
- Pugh GJF, Buckley NG, Mulder JL (1972). The role of phylloplane fungi in the early colonization of leaves symposium. *Acta. Biol. Hung.*, 11: 329-333.
- Redecker D, Kodner R, Graham LE (2000). Glomalean Fungi from the Ordovician. *Science*, 289: 1920-1921.
- Redman RS, Seehan KB, Stout RG, Rodriguez RJ, Henson JM (2002). Thermo tolerance generated by plant/fungal symbiosis. *Science*, 298: 581.
- Rajagopal K (1998). Biology and ecology of endophytic fungi from tropical trees with special reference to neem (*Azadirachta indica* A. Juss). PhD dissertation, University of Madras, Chennai.
- Rajagopal K, Kalavathy S, Kokila S, Kathiravan G, Prasad R, Balasubramanian P (2010). Diversity of fungal endophytes in few medicinal herbs of South India. *Asian J. Exp. Biol.*, 1(2): 415-418.
- Rajagopal K, Kathiravan G, Karthikeyan S (2011). Extraction and characterization of melanin from *Phomopsis*: A phellogenic fungi isolated from *Azadirachta indica* A. Juss. *Afr. J. Microbiol. Res.*, 5 (7): 762-766.
- Rodriguez RJ, Redman RS, Henson JM (2004). The role of fungal symbiosis in the adaptation of plants to high stress environments. *Mitig. Adapt. Strategies glob. Chang.* 9: 261-272.
- Sampson K (1935). The presence and absence of endophytic fungi in *Lolium temulentum* and *L. perenne*. *Trans. Br. Mycol. Soc.*, 19: 337-343.
- Sanders IR, Koide RT, Shumway DL (1995). Community-level interactions between plants and vesicular-arbuscular mycorrhizal fungi. In Varma A, Hock B (eds.), *Mycorrhiza*, Springer-Verlag, Berlin.
- Schmidt JP, Mueller GM (2007). An estimate of the lower limit of global fungal diversity. *Biodivers. Conserv.*, 16: 99-111.
- Schutz B, Boyle C, Draeger S, Rommert AK, Krohn K (2002). Endophytic fungi; a source of novel biologically active secondary metabolites. *Mycol. Res.*, 106: 996-1004.
- Sieber-Canavesi F, Petrini O, Sieber TN (1991). Endophytic *Leptostroma* species on *Picea abies*, *Abies alba*, and *Abies balsamea*: a cultural, biochemical, and numerical study. *Mycologia*, 83: 89-96.
- Souvik K, Marc L, Sebastian Z, Michael S (2008). An endophytic fungus from *Hypericum perforatum* that produces Hypericin. *J. Nat. Prod.*, 71(2): 159-162.
- Strobel GA (2002). Microbial gifts from rain forests. *Can. J. Plant Pathol.*, 24:14-20.
- Strobel GA, Daisy B (2003). Bioprospecting for microbial endophytes and their natural products. *Microbiol. Mol. Biol. Rev.*, 67: 491-502.
- Sturz AV, Christie BR, Nowak J (2000). Bacterial endophytes: Potential role in developing sustainable systems of crop production. *Crit. Rev. Plant Sci.*, 19: 1-30.
- Suryanarayanan TS, Rajagopal K (1998). Fungal endophytes in leaves of some south Indian tree species. *Proceedings of the Asia – Pacific Mycological Conference on Biodiversity and Biotechnology*, Hua-Hin, Thailand. pp. 252-256.
- Suryanarayanan TS, Kumaresan V, Johnson A (1998). Foliar fungal endophytes from two species of the mangrove *Rhizophora*. *Can. J. Microbiol.*, 44: 1003-1006.
- Suryanarayanan TS, Rajagopal K (2000). Fungal endophytes (Phellogenic) of some tropical forest trees. *Indian For.*, 126(2): 165-170.
- Tan RX, Zou WX (2001). Endophytes: a rich source of functional metabolites. *Nat. Prod. Rep.*, 18: 448-459.
- Tomita F (2003). Endophytes in South East Asia and Japan: their taxonomic diversity and potential applications. *Fungal Divers.* 11: 169-175.
- Urairaj C, Khanongnuch C, Lumyong S (2003). Ligninolytic enzymes from tropical endophytic xylariaceae. *Fungal Divers.* 13: 209-219.
- Wildman HG (2003). The rise and fall of natural products screening for drug discovery. *Fungal Divers.* 13: 221-231.
- Wilson D (1995). Endophyte - the evolution of a term, and clarification of its use and definition. *Oikos*. 73: 274-276.