

Full Length Research Paper

***In vivo* and *in vitro* protein profiling in *Acacia nilotica* (L.): A nitrogen fixing tree**

Kshipra Dhabhai* and Amla Batra

Plant Tissue Culture Laboratory, Department of Botany, University of Rajasthan, Jaipur, Rajasthan 302012, India.

Accepted 22 August, 2011

Present investigation was the first attempt which deals with the *in vivo* and *in vitro* comparative study of protein level in *Acacia nilotica* L., a nitrogen fixing tree. Protein was investigated in callus, seed, leaf and stem by means of SDS-PAGE. For obtaining the *in vitro* explant, the seeds were cultured on ½ MS medium under the *in vitro* condition. The cotyledonary nodal explants were taken from the *in vitro* seedlings and cultured in the MS medium supplemented in combination of 2, 4-D (0.4 mg/l) with BAP (0.25 mg/l). Though some differences were observed in the protein contents of *in vivo* and *in vitro* samples, the data proved that protein content in callus was higher than the seed following leaf and stem. In this study it was found that *A. nilotica* contained several protein bands of molecular weight 54.3, 44.1, 42.7, 40.1, 35.6, 31.2, 28.6, 24.7 and 19.5 kDa. These results indicate that the intensity of protein bands was high in *in vitro* sample compared to *in vivo* samples.

Key words: SDS-PAGE, *in vitro*, *in vivo*

INTRODUCTION

Acacia nilotica L. is native to the drylands of tropic Africa. It is found in India, Australia and Africa. *Acacia nilotica* L. belongs to the family leguminosae, commonly called as Babool or Kikar. Due to their ability of restore soil fertility, it is known as nitrogen fixing tree. *Acacia* species are a possible source of protein for human use (Bukhari, 2002). *Acacias* obtain their nitrogen from ground water rather than from atmosphere, but can produce more crude protein per ha than many grain crops (Prakash et al., 2001). At the molecular level, from the last century, many researches have been done on *Acacia* species. While, Protein and enzyme analysis of seeds in *Acacia* species (Ali, 1994). Seed protein analysis in *Acacia* species (Akkad, 2004), RAPD analysis in *Acacia* species (Rashmi et al., 2004), Genetic variability Analysis in *Acacia nilotica* L. (Ndir et al., 2008) and Extraction of DNA in *Acacia nilotica* L. (Sablok et al., 2009). However, no research has been done in *Acacia* species for comparison of protein profile *in vitro* and *in vivo*. Although, other medicinal plant species were examined for their protein *in vivo* and *in*

vitro viz. Bacopa monnieri (Mohapatra and Rath, 2005); *Boerhaavia diffusa* (Sharma, 2006).

MATERIALS AND METHODS

Establishment of aseptic seedlings

Mature seeds of *Acacia nilotica* L. were collected from Sirsi (Haatoj) District, Jaipur, Rajasthan. Prior to surface sterilization, seeds were treated in boiled water at 60°C for about 30 min and then soaked in distilled water for about 24 h. Then they were kept under running tap water for about 10 to 15 min followed by washing with 1% (v/v) Rankleen (Ranklem-India) for 2 min and rinsed with double distilled water for three times. Prior to inoculation, sterilized seeds were again sterilized with 0.1% (w/v) aqueous HgCl₂ for about 2 min followed by 2 to 3 rinsing with double distilled water in Laminar Air flow cabinet. These sterilized seeds were inoculated on half strength (Murashige and Skoog) MS salts medium in cultured bottles. After 7 to 10 days, seeds germinated and gave rise seedlings. These *in vitro* seedlings were used as source of explants.

Callus induction and maintenance

The Murashige and Skoog (MS) medium was prepared by adding 3% sucrose as a carbon source and 0.8% (w/v) agar as a solidifying agent. *In vitro* cotyledonary node (1.0 cm) of 20 day old

*Corresponding author. E-mail: kshipramicrobiology@gmail.com.

Table 1. Protein profiling with intensities in *Acacia nilotica* L.

Marker molecular weight (KDa)	Leaf	Stem	Callus	Seed
61.5	-	-	+	+
60.9	+	-	++	++
58.1	-	+	+	+
43.9	-	-	+	+
42.3	+	+	++	+
40.7	+	-	+++	+++
35.6	+++	+++	+++	-
31.2	-	+	+	+
28.5	+++	+++	+++	+++
24.7	+	-	+	+
19.5	+	++	++	++

Strong intensity =+++ , moderate intensity =++ , weak intensity =+ , Absent= - .

seedlings inoculated as explants for callus induction on MS medium supplemented with a series of 2, 4-D (0.4 mg/l) combination with BAP (0.2 mg/l).

The pH of medium was adjusted to 5.8 ± 0.2 before autoclaving at 121°C for 15 min at 15 lb/in². 20 ml of molten agar medium was poured into a culture bottle and plugged with nonabsorbent cotton. All cultures were incubated in 16 h /8 h photoperiod under light intensity of 50 $\mu\text{E}/\text{m}^2/\text{s}$ provided by cool, white and fluorescent light at $25 \pm 2^\circ\text{C}$ with 55% relative humidity. Each treatment performed using eight replicates and the experiment was repeated at least thrice.

Protein content determination

Leaves and stem collected from disease free and healthy plant of *Acacia nilotica* L. The specimen was authenticated by the department of Botany, University of Rajasthan and the voucher specimen (Voucher No. RUBL 20432) was deposited for future reference in the Botany Department Herbarium. Leaves and stem were used as an *in vivo* sample for comparing protein content with green, friable calli from cotyledon obtained after 28 days.

For protein estimation these samples were lyophilized, macerated in 80% ethanol and elucidated by the method of Lowry et al., (1951). Protein in the unknown sample was estimated at 660 nm using bovine serum albumin as standard and expressed per gm fresh weight basis.

Analysis of protein profile by SDS-PAGE

Electrophoresis has become a useful tool for the characterization of plant proteins. Protein profiles were studied by sodiumdodecylsulphate polyacrylamide gel electrophoresis (SDS-PAGE) (Laemmli, 1970). A vertical slab gel apparatus as described by Studier (1973), Desatron 3000/200 power supply and Frigostat, West Germany, were used during the electrophoretic work. In SDS-PAGE, proteins are treated with sodium dodecyl sulfate (SDS) before electrophoresis so that the charge density of all proteins is made roughly equal. When these samples are electrophoresed, proteins are separated according to mass. The protein bands were visualized by transilluminator and photographs were taken for comparison of results.

RESULTS AND DISCUSSION

During the present set of experiment, the total protein

estimated in callus, *in vivo* leaves, stem and seed (Figure 1) after that to find out the molecular weight of the total protein these were subjected to SDS-PAGE analysis. The proteins were found to be composed of a total of 33 bands ranging from 14.4 to 66.2 KDa were recognized (Figure 2). Protein profiles further showed variability on the basis of presence or absence and intensities of protein bands with banding pattern (Table 1).

SDS-PAGE is considered as a reliable method of genetic characterization because electrophoretic patterns of the protein fractions are directly related to the genetic background of the proteins and can be used to certify the genetic make-up (Rehana et al., 2004). In order to estimate the variability at genetic level, SDS-PAGE banding pattern of the gel using total protein was investigated.

Overall out of 33 protein bands, molecular weights 19.5, 28.5 and 42.3 KDa shown same protein banding pattern in callus, *in vivo* leaves, stem and seed samples but with variation in intensities. Left behind protein banding pattern exhibited a considerable range of variability with regard to their mobilities and intensities. Callus exposed at protein molecular weight ranging from 14.4 to 66.2 KDa. Seed only lacking at the molecular weight 35.6 KDa but leaves absent in 31.2, 58.1 and 61.1 KDa. Whereas, stem missing in the molecular weight of 24.7, 40.7, 43.9 and 60.9 KDa. From this we concluded that the highest numbers of protein bands were observed in callus followed by seed, leaves and stem. Regarding this experiment, no similar and contrary results were available in *Acacia nilotica* L. and *Acacia* species, but in other plants, analogous results were reported in *Artemisia vulgaris* (Kumar and Ranjitha, 2009), *Glycine max* (L.) Merr (Radhakrishnan and Ranjitha, 2009), and *Plumbago zeylanica* L. (Rout et al., 2010). No research has been done contrary to these results. This paper presents and sheds light on the *Acacia nilotica* L. proteins which grow on the Nile banks of Delta region and in arid regions. Therefore, very much attention should be given to arid species since they are under severing threats




Figure 1. Protein estimation in *Acacia nilotica* L.




Figure 2. Protein analysis via. SDS-PAGE in *Acacia nilotica* L.

these valuable plants which showed some signs as an important resource for desert ecosystem and arid environment.

ACKNOWLEDGEMENTS

The authors are thankful to the Head of the Department of Botany, University of Rajasthan, Jaipur, for providing the basic research facilities. The authors are also sincerely grateful to Dr. M. M. Sharma, R. N. Verma, Govind Vyas, Sakshi Singh and Antara Sen, from the Department of Botany, University of Rajasthan, Jaipur for their critical review and helpful discussions.

REFERENCES

- Ali A, Al- Helal (1994). Electrophoretic analysis of protein and selected enzymes of *Acacia* seeds from Saudia Arabia. *JKAU:Sci.*, 6:17-28.
- Akkad SS (2004). Phylogenetic Relationship and Similarity Indices of Some *Acacia* Species Using Seed Protein Analysis. *Intr. J. Agr. Biol.*, 6(3).
- Bukhari YM (2002). Seed weight and protein concentration of whole seed, cotyledons and seed coat of 57 accessions of *Acacia* and *Prosopis* (Mimosaceae): Phylogenetic and nutritional implications. *Indian For.*, 128: 686–699.
- Kumar PS, Kumari Ranjitha BD (2009). In vitro and In vivo Identification of variation in protein expression in *Artemisia vulgaris* L. *Adv. Biol. Res.*, 3(5-6): 63-70.
- Laemmli UK (1970). Cleavage of structural proteins during the

- assembly of the head of bacteriophage T4. *Nature*, 227: 680-685.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951). Protein measurement with the Folin Phenol reagent. *J. Biol. Chem.*, 193: 265-275.
- Mohapatra HP, Rath SP (2005). *In vitro* studies of *Bacopa monnieri* – an important medicinal plant with reference to its biochemical variations. *Indian J. Exp. Biol.*, 43(4): 373-376.
- Ndir KN, Samb PI, Chevallier MH (2008). Genetic Variability Analysis of the Polyploid Complex of *Acacia nilotica* (L.) Willd. Using RAPD Markers. *Tropiculture*, 26(3): 135-140.
- Prakash D, Niranjana A, Tewari SK, Pushpangadan P (2001). Underlisted legumes: Potential sources for low-coast protein. *Int. J. Food Sci. Nutr.*, 52: 337–341.
- Radhakrishnan R, Ranjitha Kumari BD (2009). Changes in Protein Content in Micropropagated and Conventional Soybean Plants (*Glycine max* (L.) Merr.). *World J. Agric. Sci.*, 5(2): 186-189.
- Rashmi M, Nanda Nayak S, Rout GR (2004). Studies on genetic relatedness of *Acacia* tree species using RAPD markers. *Biologia, Bratislava*, 59(1): 115-120.
- Rehana A, Rabia S, Afzal M, Akthar S (2004). Inter and Intraspecific variation in SDS-PAGE of total seed protein in rice (*Oryza sativa* L.) germplasm. *Pak. J. Biol. Sci.*, 7: 139-143.
- Rout JR, Kanungo S, Das R, Sahoo SL (2010). *In vivo* protein profiling and Catalase Activity of *Plumbago zeylanica* L. *Nat. Sci.*, 8(1): 87-90.
- Sablok G, Gahlot P, Gupta AK, Pareek K, Shekhawat NS (2009). Extraction of PCR-usable DNA from trees adapted to arid environment. *Plant Omics. J.*, 2(3): 103-109.
- Sharma A (2006). *In vitro* morphogenetic studies of *Boerhaavia diffusa* L. Ph.D. Thesis University of Rajasthan.
- Studier FW (1973). Analysis of bacteriophage T7 early RNAs and proteins on slab gels. *J. Mol. Biol.*, 79: 237-248.