

Full Length Research Paper

# General steps in the production and establishment of transgenic chicken lines carrying a bacterial lacZ gene expressing $\beta$ -galactosidase

Esmailzadeh Mahdi\*, Nejadali Abolfazl and Kazemzadeh Fariba

Basic Sciences Department, Nikshahr Branch, Islamic Azad University, Nikshahr, Iran.

Accepted 27 December, 2011

Replication-defective retroviral vectors are efficient vehicles for the delivery of exogenous genes, and they may be used in the generation of transgenic animals. The replication-defective retroviral SNTZ vector carrying the lacZ gene with a nuclear localized signal was injected into the subgerminal cavity of freshly laid eggs. Subsequently, the eggs were allowed to hatch, and the chickens were screened for the lacZ gene by using the polymerase chain reaction. Eight of 15 male chickens that survived to sexual maturity contained the lacZ gene in their semen. Subsequently, these males were mated with wild-type female chickens. In this study, performance of the SNTZ vector carrying the *Escherichia coli* lacZ gene to express a nuclear-localized  $\beta$ -galactosidase was injected into the subgerminal cavity of embryos collected from newly laid eggs was reviewed.

**Key words:** Transgenic chickens, lacZ,  $\beta$ -galactosidase and retroviral vector.

## INTRODUCTION

Of the current transgenesis methods for birds, retroviruses present the most effective gene delivery systems (Shuman and Shoffner, 1986; Shuman, 1991; Bosselmann et al., 1989, 1990, Petite, 2002; Sang, 2004). Two types of avian retroviral vectors have been used: the replication competent virus and replication-defective virus.

The replication-competent retroviruses retain all genomic sequences necessary for producing infectious viral particles, integration, and self-replication from the host genome whereas the replication-defective viruses retain the packaging signal, sequences for the viral integration, and transcription initiation but most parts of the retroviral genes removed and inserted with foreign deoxyribonucleic acid (DNA). Thus, the replication-defective retroviral vectors require the packaging cell lines to supply all viral proteins needed to assemble infectious particles. Once the retroviral RNA genome

enters the host, the viral ribonucleic acid (RNA) is reverse transcribed into double-stranded DNA; migrates to the host genome and is integrated as the provirus. At that point, proviral DNA is only replicated along with the host genome (Shuman, 1991). The utilization of the replication-competent viruses to produce transgenic chickens is not practical because of the limited size of the inserted genes and the pathogenesis (Shuman, 1991; Wentworth and Wentworth, 2000).

The replication-defective retroviral vectors derived from the avian leukosis and sarcoma viruses (ALSV) and the reticulo endothelosis viruses (REV) were used to produce transgenic chickens without the evidence of viremia and horizontal infection of helper viruses. Vick et al. (1993) produced transgenic birds with Avian Leukosis Virus-based vectors, but did not report the expression of the transgenes carrying neomycin resistance (Neor) and the lacZ gene. Thoroval et al. (1995) used the ALV-based vectors to produce the transgenic chickens with Neor and the lacZ gene, but the evaluation of the expression was limited to chicken embryonic fibroblasts (CEFs). A successful production of transgenic chickens expressing a reporter gene that could be useful for cell lineage has

\*Corresponding author. E-mail: mehdi\_dna@yahoo.com. Tel: +98 935 979 3491.

not been reported.

Mikawa et al. (1991, 1992a, b) developed a SNV-based replication defective retroviral vector carrying a lacZ gene, called SNTZ, for cell lineage analysis of cardiac development. This retroviral vector can infect a wide variety of cell types in early chick embryos. Beta-galactosidase, the product of the lacZ gene, was functional in various cell types in various embryonic tissues and cell cultures. Furthermore, the lacZ gene contained a nuclear location sequence to concentrate activity in the nucleus. The lacZ gene was stably inherited and expressed by the progeny of the infected cells. Moreover, expression of the gene could be detected at hatch.

Thus, it appears that the SNTZ vector would be an appropriate retroviral system potentially for the generation of transgenic chickens expressing  $\beta$ -galactosidase. The main objective of the study was to use SNV-based replication defective SNTZ retroviral vector developed by Mikawa et al. (1992b) to produce the transgenic chickens carrying the lacZ gene encoding nuclear-localized  $\beta$ -galactosidase. The goal of this study was to reviewed general steps in the production transgenic chicken lines to stably inherit the lacZ gene and express in a variety of tissues throughout developmental stages which can be used as an ideal cell marker for cell lineage studies.

## MATERIALS AND METHODS

In the production of transgenic chickens, first the retroviral vectors are harvested and injected into a prospective G0 embryo before incubation. The injected embryos were cultured in a surrogate eggshell and hatch at approximately 21 days of incubation. After hatching, it takes another 17 to 20 weeks for the birds to reach sexual maturity. Semen can be collected and screened for the transgene. In general, the positive G0 roosters are preferably used in mating program for the production of transgenic G1 progeny carrying the transgene from many females. This step is inefficient due to the mosaic status of the germline of G0 birds. Subsequently, the transgenic G1 offspring will take another 21 weeks to reach sexual maturity and at least another 3 to 4 weeks until there is sufficient G2 progeny to demonstrate Mendelian inheritance.

The replication-defective retroviral vector was derived from SNV, except for the splice acceptor sequences, which were derived from avian reticuloendotheliosis virus strain A (REV-A), a virus which has 98% nucleotide sequence homology to SNV (Dougherty and Temin, 1986) (Figure 2). Unintegrated linear viral SNV DNA is approximately 8.3 kbp long (Chen et al., 1981) (Figure 1).

The plasmid pJD214 was constructed to lack the 5' end of the 5'LTR and the 3' end of the 3' LTR (Dougherty and Temin, 1986). The gag, pol, and env sequences were removed, replaced with a pUC12 polylinker and ligated into the pBR322 at EcoRI/BamHI sites resulting in only 1.45 kbp in length.

The pJD214 was modified for convenience with future cloning by removing one PstI site located within the ampicillin-resistance gene with PvuI and BglI and inserted the PvuI-BglI of pGEM4 (Figure 2). The new plasmid construct was called pJDp- (Mikawa et al., 1992a). A fragment containing the SV40 early region promoter and enhancer sequences were isolated from pMSG (Pharmacia, Piscataway, NJ) and inserted into the BamHI/HindIII sites of pGEM4 (Promega, Madison, WI). Then the insert was isolated with EcoRI/HindIII. A HindIII/EcoRI fragment containing the neomycin

resistance gene with a splice donor and polyA signal was isolated from pSV2neo. These two fragments were then co-ligated into the EcoRI site of pJDp- (Figure 4). The new plasmid DNA construct, called pSN, encodes the SV40 promoter and neo-resistance gene in the same orientation as the SNV long terminal repeat (Mikawa et al., 1992b) (Figure 3).

The lacZ sequences fused to the nuclear leading or localization sequence (NLS) of SV40 large T antigen was isolated from the plasmid LZ12. The construction of the plasmid LZ12 was described by Galileo et al. (1990) (Figure 4). The 5' end of 2-kb fusion sequence of pLZ12 for the lacZ gene was isolated by Bgl II/Sst I digestion and ligated into the Bam HI/Sst I sites of pGEM4. The insert was then removed from the vector with Xba I/Sst I. The 1-kbp fragment of the 3' end of lacZ was cut from pMC1987 (Pharmacia) by Sst I/Sal I digestion. These two fragments were ligated into pSN at Xba I/Sal I site to form pSNTZ (Mikawa et al., 1992b) (Figure 5). The pSNTZ is about 7.9 kbp.

## Production of retroviral virions containing SNTZ

The created pSNTZ vector encoding lacZ was transfected into the SNV packaging cell lines, D17.2G, which provides the products of gag, pol, and env gene in trans without producing infectious virus as a result of the deletion of E (Dougherty and Temin, 1988). The D17.2G SNTZ-producing cells were grown in Dulbecco's Modified Eagle's Medium (DMEM), supplemented with 7% fetal bovine serum, and 1% penicillin-streptomycin.

All the embryos used for retroviral transfection were obtained from freshly laid unincubated fertile White Leghorn eggs which were at about stage X (Eyal-Giladi and Kochav, 1976) with approximately 50,000 cells per embryo.

Embryo cultures were performed based on the procedures of Perry (1988) and modified by Borwornpinyo (2000) and Borwornpinyo et al. (2005). The embryos from the newly laid eggs (donor eggs) were transferred into recipient chicken eggshells through the window cut at the sharp end.


After 3 d of incubation, the embryos and whole culture contents were transferred into new recipient turkey eggshells through the window cut at the blunt end. The birds generated from viral injection formed the G0 chimeric founder chickens.

## DNA isolation

Genomic DNA isolation from chicken blood was modified from Petite et al. (1994). First, 5  $\mu$ l of whole blood was diluted with 45  $\mu$ l PBS (1:10 dilution) in a 0.5 ml microtube, and then mixed with 100  $\mu$ l lysis buffer (10 mM Tris HCl pH 7.5, 5 mM MgCl<sub>2</sub>, 0.32 M sucrose, 1% Triton X-100) by vortexing. The mixture was microcentrifuged at 1,200  $\times$  g for 20 sec to pellet the nuclei of erythrocytes and the supernatant was discarded. After removing the supernatant, the pellets of nuclei were incubated in 100  $\mu$ l TEN buffer (10 mM Tris HCl pH 8.0, 2 mM EDTA, 400 mM NaCl, 1% SDS, and 12  $\mu$ l Proteinase-K (20 mg/ml) at 50°C for overnight. After incubation, 50  $\mu$ l saturated NaCl was added to precipitate proteins and microcentrifuged at 1,200  $\times$  g for 15 min. The supernatant was transferred into a 1.5 ml microtube. Two volumes of ethanol were added to the tube and microcentrifuged at 1,600  $\times$  g for 15 min to precipitate DNA. The recovered DNA was air dried and resuspended in 50  $\mu$ l Tris-EDTA buffer.

## PCR screening

The presence of the lacZ gene in the offspring was determined using Polymerase chain reaction (PCR). Genomic DNA isolated



**Figure 1.** Vector construction and deduced restriction map of pJD214. The sequences required in cis for viral replication are the primer binding site (PBS), polypurine tract (PPT), and encapsidation sequence (E). Other abbreviations: SS, splice site and LTR, long terminal repeat. The multiple cloning sites were derived from the polylinker of pUC12. The virus RNA derived from pJD214 is 950 bp long. The unintegrated SNV DNA is approximately 8.3 kbp in length.


from various sources as described above was subjected to 35 cycles of PCR for the presence of the *E. coli* lacZ gene. The 3-steps pattern involved denaturation at 95°C for 30 s, primer annealing at 54°C for 1 min and extension at 72°C for 1 min using a thermocycler 8.

A 25 µl reaction volume containing 200 ng of DNA samples, 1.5 mM MgCl<sub>2</sub>, 0.8 U of Taq polymerase, 1 µM of dNTPs and 1 µM of each primer, the forward primer 5' TTCTGTATGAACGGTCTGGTC-3', and the reverse primer 5'-ACTTACGCCAATGTCGTTATC-3' was used to amplify a 588-bp fragment specific to the lacZ gene.

Subsequently, the 588-bp amplified products were fractionated through a 1.5% agarose gel and visualized with ethidium bromide.

#### β -Galactosidase expression

Experiment demonstrate β -galactosidase express in G2 chickens was to isolate myoblasts from the pectoralis thoracicus muscle to show that *E. coli* β -galactosidase is expressed in the myogenic satellite cell population after *in vitro* culture. Myoblasts from three of



**Figure 2.** Vector construction and deduced restriction map of pJDp-. The cis sequences of pJD214 illustrated in Figure 2. The pGEM4 was obtained from Promega Corporation.


six G2 chickens express  $\beta$ -galactosidase *in vitro* (Figure 6).  $\beta$ -galactosidase expression in other G2 embryos, and the embryos also express  $\beta$ -galactosidase (Figure 7).

### Staining embryos

Embryos was fix at 4°C with 2% formaldehyde and 0.2% glutaraldehyde in PBS pH 7.4 for 30 min, rinsed in PBS, and incubated in X-Gal solution (1 mg/ml X-Gal PBS pH 7.4, 5 mM potassium ferrocyanide, 5 mM potassium ferricyanide, 2 mM MgCl<sub>2</sub>, 0.2% Triton X-100) overnight in the dark at 37°C. Subsequently, the embryos were washed with PBS and stored in 70% ethanol before microscopic evaluation.

### RESULTS

Table 1 summarizes the effect of injection strategies on hatchability and the percentage of chimeric G0 chickens carrying the lacZ gene in blood and semen. When injected with the concentrated retroviral stocks into the subgerminal cavity, 24 embryos out of 66 injected embryos survived to hatch. Ten embryos out of 54 embryos injected with the concentrated retroviral-containing medium and cells survived to hatch. Fourteen of 40 embryos injected with the concentrated retroviral containing media into subgerminal cavity and again one



**Figure 3.** Vector construction and deduced restriction map of pSN.



day after incubation at the germinal crescent survived to hatch. Overall, thirty percent hatchability was obtained in the experiment (48 of the total 168 injected embryos survived to hatch).

It has been shown that embryo injection without retrovirus-producing cells resulted in the highest hatchability which accounted for thirty-six percent hatchability (24 of 66 injected embryos). Forty G0 chicks, 25 females and 15 males, survived to sexual maturity. Two G0 females carried lacZ gene in their blood DNA when using PCR screening. These two females resulted from a single injection of retroviral containing media. One-hundred-and-one progeny produced from one G0 female and 90 progeny produced from the second G0 female were screened for the lacZ gene by means of PCR in their blood (Table 2). Neither G0 females generated lacZ-positive offspring. In addition, all blood lacZ-negative G0 females did not produce lacZ-positive


progeny in their blood. Eight of 15 G0 males (53%) carried the lacZ gene in their semen DNA, but none of them were blood lacZ-positive (Figure 8). A single injection of retroviral stocks resulted in a greater number of G0 males carrying the lacZ gene in their semen (Table 1). This suggests that a single injection of concentrated retrovirus into the subgerminal cavity of the stage X embryos collected from the newly-laid eggs are the most efficient procedure for retroviral infection and gene transfer. Only one semen lacZ-positive G0 male produced 2 blood lacZ-positive chickens from a total 224 progeny which accounted for the 0.89%.

## DISCUSSION


This study attempted to show the efficiency of retroviral gene delivery into germline precursors by varying



**Figure 4.** Vector construction and deduced restriction map of pLZ12.




**Figure 5.** Vector construction and deduced restriction map of pSNTZ. The cis elements of the retroviral sequences were illustrated in Figure 1.



**Figure 6.** Myoblast cultures from day old chickens. A: Bright field photomicrograph of primary myoblast cultures from lacZ-positive chickens. Cells with stained nuclei represent  $\beta$ -galactosidase-positive cells. B: Phase-contrast photomicrograph of primary myoblast cultures from lacZ-negative chickens. Scale bar = 150  $\mu$ m in B (applies to A, B).

different methods of retroviral injection. Regardless of injection procedures, no overt differences were observed in gene transfer between injecting the concentrated retrovirus and delivering the concentrated retrovirus along with the virus-producing cell or performing multiple injections with retrovirus. Thus, a single injection of the concentrated SNTZ retrovirus into the subgerminal cavity of stage-X embryos (Eyal-Giladi and Kochav, 1976) containing approximately 50,000 cells appears to be the most efficient method for producing chimeric germline transgenic chicken founders.

Overall hatchability of the injected embryos in many experiments was 30% which was approximately the same reports (38, 36, and 27% hatchability obtained from Bosselman et al. (1989), Harvey et al. (2002), and McGrew et al. (2004), respectively). Nevertheless, the percentage hatchability in Mozdziak et al., (2003) study was relatively higher than those obtained from Thoraval et al. (1995) and Chapman et al. (2005) who reported 2.3 and 4% hatchability of viral injected embryos, respectively. The acceptable hatchability obtained in Mozdziak et al. (2003) study was mainly due to the surrogate eggshell system (Perry, 1988) with modifications (Borwornpinyo, 2000, Borwornpinyo et al., 2005) for improving survivability and hatchability.



**Figure 7.** Embryos (Stage 8; Hamburger and Hamilton, 1951). From (A)  $\beta$ -galactosidase-positive transgenic chickens and from (B) wild-type  $\beta$ -galactosidase-negative chickens. Inset in A shows staining in the somites of  $\beta$ -galactosidase-positive transgenic chickens. Scale bar = 50  $\mu$ m in inset, 300  $\mu$ m in B (applies to A, B).

In many studies, no correlation between the presence of the lacZ gene detected in the blood (somatic cells) and the sperm or ova (germ cells) were observed (Thoraval et al., 1995; McGrew et al., 2004). In the Mozdziak et al., (2003) experiment, two blood lacZ-positive hens were generated from the retroviral injection and together produced 191 G1 offspring when mated with wild-type roosters. None of the progeny carried the lacZ in the genomic DNA based on PCR screening. On the other hand, eight roosters were found lacZ-positive in their semen DNA, but their blood DNA samples were lacZ-negative.

Gene transfer into the germline using retroviral infection into embryos obtained from newly laid eggs is not efficient. Mozdziak et al. (2003) showed, a total of 1,639 offspring generated from mating 8 semen lacZ-positive roosters and wild-type hens had to be screened in order to detect 2 lacZ positive G1 chicks. Notably, only one rooster was shown to transmit the lacZ gene to the two G2 transgenic chicks which accounted for 0.89% (2/224). The low germline transmission rates were consistent with previous reports (Bosselman et al., 1989, 1990; Harvey et al., 2002; Chapman et al., 2005).

However, the efficiency of gene transfer into germline precursors based replication defective vectors can be improved by infecting the unincubated embryos with retrovirus pseudotyped with vesicular stomatitis virus

**Table 1.** Effect of injection procedures on hatchability (% hatch) and effect of injection procedure on the percentages of G0 chickens carrying lacZ gene in blood and semen.

Injection procedure	N	% Hatch (#)	% carrying lacZ in the blood (# positive/ total # screened)	% carrying lacZ in the semen (# positive/ total # screened)
Virus only <sup>1</sup>	66	36(24)	10 (2/20) <sup>4</sup>	80 (4/5)
Virus and cells <sup>2</sup>	54	19(10)	0 (0/7)	40 (2/5)
Virus double injection <sup>3</sup>	40	35(14)	0 (0/13)	40 (2/5)

N, number of eggs injected; % Hatch is the percentage of injected eggs that hatched; #, number. 1) Virus only, embryos collected from newly-laid eggs were injected with 5 µl of the concentrated retroviral stocks into the subgerminal cavity. 2) Virus and cells, embryos collected from newly-laid eggs were injected with 5 µl of the concentrated retroviral stocks containing 500 SNTZ-producing cells into the subgerminal cavity. 3) Virus double injection, embryos collected from newly-laid eggs were with 5 µl of the concentrated retroviral stocks into the subgerminal cavity, and the same embryos were again injected 1 d after incubation with 4 injections of 3 µl per injection of the retroviral stocks into the germinal crescent. 4) These 2 lacZ-positive birds in blood were females.

**Table 2.** PCR screening of the progeny from chimeric G0 females generated from retroviral injection<sup>1</sup>.

Chimeric G0 females	Number of G1 chicks PCR screened for lacZ	G1 chicks carrying lacZ in their blood
1	101	0
2	90	0
3	66	0
4	88	0
5	32	0
6	22	0
7	24	0
8	49	0
9	55	0
10	22	0
11	40	0
12	65	0

1) PCR, polymerase chain reaction. 2) These chimeric G0 hens were positive for the lacZ gene in their blood, and produced from the injection at stage X embryos collected from newly-laid eggs with concentrated virus.


envelop protein (VSV-G).

Mizuarai et al. (2001) reported that 80% germline transmission rate was obtained from the G0 quail injected with Moloney murine leukemia

virus (MoMLV) pseudotyped with VSV-G protein. Similarly, McGrew et al. (2004) generated G0 roosters that transmitted the transgene to their offspring with rates ranging from 4 to 45% based

using equine infectious anaemia virus (EIAV) vectors pseudotyped with vesicular stomatitis virus glycoprotein (VSV-G). In addition, relatively high retroviral titers can be produced when using





**Figure 8.** Polymerase chain reaction analysis of the semen from G0 chickens. Water is a negative control using a water blank. NC is a negative control using DNA isolated from semen of wild-type chickens. PC is a positive control reaction containing the lacZ gene from pmwZ plasmid (Kadokawa et al., 1990) spiked into semen DNA from wild-type chickens. Lanes 1, 2, and 6 represent negative birds; lanes 3, 4, and 5 represent positive birds. All positive lanes contain a 588-bp fragment.

**Table 3.** PCR screening of the progeny from G0 male chickens carrying the lacZ gene in their semen<sup>1</sup>.

G0 males carrying lacZ in their semen	Number of G1 chicks PCR screened for lacZ	G1 chicks carrying lacZ in their blood
12	2243	24
2	252	0
3	143	0
4	241	0
5	365	0
6	195	0
7	152	0
8	196	0

1) PCR, polymerase chain reaction. 2) This germline chimeric sires were produced from the injection at a stage X embryo collected from newly laid eggs with concentrated virus. 3) Screening of the progeny from these two sires was terminated because the birds died. 4) Germline transmission rate of this G0 sire was 0.89% (2/224)

the VSV-G-pseudotyped retroviral vectors.

The stable integration and inheritance of transgenes is one of the most important aspects for the production of transgenic chicken lines. In Mozdziak et al. (2003) study, it is clear that the germline transmission rates of G1, G2, and G3 transgenic birds are consistent with the expected Mendelian ratios indicating that the lacZ transgene acts as a hemizygous dominant allele and is stably inherited for both transgenic lines (P1 and P2) which were generated from the same G0 sire. Recently, Mozdziak research group (Mozdziak et al., 2006) studied the integration site and copy number of the lacZ provirus in both transgenic G1 male founders and their G2 offspring using Southern analysis. They found that these two G1 males contain a single copy and an identical integration site of the provirus suggesting that they are from the same germline precursor. The results confirmed that the germline transmission of transgenic chicken lines for the

lacZ gene were consistent with the predicted Mendelian ratios.

Six criteria for an ideal cell tracer for studies of cell lineage were mentioned (McLaren, 1976; Oster-Granite and Gearhart, 1981). First, the expression of  $\beta$ -galactosidase was localized only in the nucleus of transgenic cells. Moreover, this nuclear-localized  $\beta$ -galactosidase alleviates the problem of endogenous activity when X-gal staining is applied. Second, the expression was cell autonomous in that only cell containing the lacZ exhibit staining and the  $\beta$ -galactosidase is not transferred among cells. Third, the lacZ gene is stably inherited and expresses  $\beta$ -galactosidase in subsequent cell division. Fourth, the expression of  $\beta$ -galactosidase was ubiquitous among both the internal and external tissues of the body. The  $\beta$ -galactosidase activity can be detected in entire embryos from at least stages 8, 20, and 28 (Hamburger and Hamilton, 1951).

G3 birds expressed the lacZ in all tissues examined including brain, bursa, gizzard, intestine, liver, lung, kidney, skeletal muscle, spleen, and testes. Fifth, the expression was easy to detect, both grossly and in histological sections using X-gal staining. However, the expression was not present in every cell based on the X-gal staining. This might be due to the low sensitivity of detection using X-gal staining (Couffinhal et al., 1997).

Immunohistochemistry staining using anti- $\beta$ -galactosidase antibody drastically improved the detection of the positive cells carrying the lacZ gene (Couffinhal et al., 1997). In addition, *insitu* polymerase chain reaction (*insitu* PCR) could be used to localize all cells carrying a single copy of the lacZ gene in tissue sections (Komminoth and Long, 1993). Finally, the transgenic birds were normal throughout development indicating that they were developmentally neutral, not causing cell selection or influencing developmental processes. Recently, satellite cells isolated from skeletal muscle of transgenic adult chickens can be used to trace their fate after implantation into stage 14 chicken embryos (Mozdziak et al., 2006).

Furthermore, Zhang et al. (2006) have shown that hematopoietic stem cells (HSCs) have potential to differentiate into multiple non-blood cell lineages using parabiosis between quail and transgenic chick embryos. Thus, production transgenic birds as an alternative cell marker can be used in experiments in homospecific grafts for cell lineage studies and to confirm and validate the results obtained from the chick-quail cell tagging method. In addition, this stable genetic tagging is ideally suitable for long-term cell lineage analysis because the marker expression does not decline over time and the chimeras made in embryos between the same species and strains would survive throughout their entire lives without immunological rejection.

## Conclusion

In this study, the stages of production transgenic chickens carrying the lacZ gene and expressing  $\beta$ -galactosidase have been reviewed. The expression of  $\beta$ -galactosidase can be observed in all examined tissues from embryos and adults at different stages of development. The produced transgenic chickens could meet the six criteria for an ideal cell marker which would be (1) cell localized, not secreted extracellular; (2) cell autonomous, not transferred between cells; (3) stable in all cells; (4) ubiquitous throughout development in all tissue types; (5) easy to detect, both grossly and in tissue sections, and (6) developmentally neutral.

## REFERENCES

Borwornpinyo S (2000). Optimal hatchability of cultured chicken embryos from freshly laid eggs in surrogate eggshell (master's thesis). Raleigh NC: North Carolina State University.  
Borwornpinyo S, Brake J, Mozdziak PE, Petite JN (2005). Culture of

chicken embryos in surrogate eggshells. *Poult. Sci.*, 84:1477-1482.  
Bosselman RA, Hsu RY, Boggs T, Hu S, Bruszewski J, Ou S, Kozar L, Martin F, Green C, Jacobsen F, Nicolson M, Schultz JA, Semon KM, Rishell M, Steward RG (1989). Germline transmission of exogenous genes in the chicken. *Science*, 243:533-535.  
Bosselman RA, Hsu RY, Brinskin MJ, Boggs T, Hu S, Nicolson M, Souza LM, Schultz JA, Rishell W, Steward RG (1990). Transmission of exogenous genes into the chicken. *J. Reprod. Fert. Suppl.*, 41: 183-195.  
Chapman SC, Lawson A, MacArthur WC, Wiese RJ, Loechel RH, Burgos-Trinidad M, Wakefield JK, Ramabhadran R, Mauch TJ, Schoenwolf GC (2005). Ubiquitous GFP expression in transgenic chickens using a lentiviral vector. *Development*, 132:935-940.  
Chen ISY, Mak TW, O'Rear JJ, Temin HM (1981). Characterization of reticuloendotheliosis virus strain T DNA and isolation of a novel variant of reticuloendotheliosis virus strain T by molecular cloning. *J. Virol.*, 40:800-811.  
Couffinhal T, Kearney M, Sullivan A, Silver M, Tsurumi Y, Isner JM (1997). Histochemical staining following lacZ gene transfer underestimates transfection efficiency. *Hum. Gene Ther.*, 8:929-934.  
Dougherty JP, Temin HM (1986). High mutation rate of a spleen necrosis virus-based retrovirus vector. *Mol. Cell Biol.*, 168:4387-4395.  
Dougherty JP, Temin HM (1988). Determination of the rate of base-pair substitution and insertion mutations in retrovirus replication. *J. Virol.*, 62:2817-2822.  
Eyal-Giladi H, Kochav S (1976). From cleavage to primitive streak formation: a complementary normal table and a new look at the first stages of the development of the chick. *Dev. Biol.*, 49: 321-337.  
Galileo DS, Gray GE, Owens GC, Majors J, Sanes JR (1990). Neurons and glia arise from a common progenitor in chicken optic tectum: Demonstration with two retroviruses and cell type-specific antibodies. *Proc. Natl. Acad. Sci. USA*, 87:458-462.  
Hamburger V, Hamilton HL (1951). A series of normal stages in the development of the chick embryo. *J. Morphol.*, 88:49-92.  
Harvey AJ, Speksnijder G, Baugh LR, Morris JA, Ivarie R (2002). Consistent production of transgenic chickens using replication-deficient retroviral vectors and high throughput screening procedures. *Poult. Sci.*, 81:202-212.  
Komminoth P, Long AA (1993). *In-situ* polymerase chain reaction. An overview of methods, applications, and limitations of a new molecular technique. *Virchows. Arch. B. Cell Pathol. Incl. Mol. Pathol.*, 64:67-73.  
McGrew MJ, Sherman A, Ellard FM, Lillico SG, Gilhooley HJ, Kingsman AJ, Mitrophanous KA, Sang H (2004). Efficient production of germline transgenic chickens using lentiviral vectors. *EMBO Rep.*, 5:728-733.  
McLaren A (1976). *Mammalian Chimaeras*. Cambridge Univ. Press, London/New York.  
Mikawa T, Borisov A, Brown AMC, Fischman DA (1992a). Clonal analysis of cardiac morphogenesis in the chicken embryo using a replication-defective retrovirus: I. Formation of the ventricular myocardium. *Dev. Dyn.*, 193:11-23.  
Mikawa T, Cohen-Gould L, Fischman DA (1992b). Clonal analysis of cardiac morphogenesis in the chicken embryo using a replication-defective retrovirus. III: Polyclonal origin of adjacent ventricular myocytes. *Dev. Dyn.*, 195:133-141.  
Mikawa T, Fischman DA, Dougherty JP, Brown AMC (1991). In vivo analysis of new lacZ retroviral vector suitable for cell lineage lineage marking in avian and other species. *Exp. Cell Res.*, 195:516-523.  
Mizuarai S, Ono KI, Yamaguchi K, Nishijima KI, Kamihira M, Iijima S (2001). Production of transgenic quails with high frequency of germline transmission using VSV-G pseudotyped retroviral vectors. *Biochem. Biophys. Res. Com.*, 286:456-463.  
Mozdziak PE, Wu Q, Bradford JM, Pardue SL, Borwornpinyo S, Giamario C, Petite JN (2006). Identification of the lacZ insertion site and betagalactosidase expression in transgenic chickens. *Cell Tissue Res.*, 324:41-53.  
Mozdziak PE, Borwornpinyo S, McCoy DW, Petite JN (2003). Development of Transgenic Chickens Expressing Bacterial  $\beta$  - Galactosidase. *Dev. Dyn.*, 226:439-445.  
Oster-Granite ML, Gearhart J (1981). Cell lineage analysis of cerebellar

- purkinje cells in mouse chimeras. *Dev. Biol.*, 85:199-208.
- Perry MM (1988). A complete culture system for the chick embryo. *Nature*, 331:70-72.
- Petitje JN (2002). The avian germline and strategies for the production of transgenic chickens. *Poult. Sci.*, 39:205-228.
- Petitje JN, Kegelmeyer AE, Kulic MJ (1994). Isolation of genomic DNA from avian whole blood. *Biotechniques*, 17:664-666.
- Sang H (2004). Prospects for transgenesis in the chick. *Mech. Dev.*, 121:1179-1186
- Shuman RM (1991). Production of transgenic birds. *Experientia*, 47: 897-905.
- Shuman RM, Shoffner RN (1986). Gene transfer by avian retroviruses. *Poult. Sci.*, 65:1437-1444.
- Thoraval P, Afanassieff M, Cosset FL, Lasserre F, Verdier G, Coudert F, Dambrine G (1995). Germline transmission of exogenous genes in chickens using helper-free ecotropic avian leucosis virus-based vectors. *Transgenic Res.*, 4:369-377.
- Vick L, Li Y, Simkiss K (1993). Transgenic birds from transformed primordial germ cells. *Proc. R. Soc. Lond. B Biol. Sci.*, 251:179-182.
- Wentworth AL, Wentworth BC (2000). The avian primordial germ cell: An enigma of germline evolution. *Development and immortality. Avian Poult. Biol. Rev.*, 11: 173-282.
- Zhang N, Mustin D, Reardon W, DeAlmeida A, Mozdziaik P, Mrug M, Eisenberg LM, Sedmera D (2006). Blood-borne stem cells differentiate into vascular and cardiac lineages during normal development. *Stem Cells Dev.*, 15:17-28.

RETRACTED