

Short Communication

Coxiella burnetii in goat bulk milk samples in Iran

Ebrahim Rahimi

Department of Food Hygiene, Faculty of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran. E-mail: rahimi@iaushk.ac.ir

Accepted 19 July, 2010

Q fever is a zoonotic disease caused by the rickettsial organism *Coxiella burnetii*, the objective of this study was to determine the prevalence rate of *C. burnetii* in bulk milk samples from dairy goat herds in Fars, Ghom, Kerman, Khuzestan and Yazd provinces, Iran. In the present study, 296 bulk milk samples from 89 dairy goat herds were tested for *C. burnetii* using a nested PCR assay. The animals which their milk samples collected for this study were clinically healthy. In total, 6 of 296 (2.0%) goat milk samples were positive; the positive samples originated from 4 of 22 (18.2%) dairy herds in Fars, 1 of 24 (4.2%) dairy herds in Khuzestan and 1 of 18 (5.5%) dairy herds in Yazd. All 76 goat bulk milk samples from 25 goat breeding farms collected in Ghom and Kerman provinces were negative, although no extensive prevalence study was undertaken, the results of this study indicate those clinically healthy dairy goats are important sources of *C. burnetii* infection in Iran.

Key words: *Coxiella burnetii*, PCR, milk, goat, Iran.

INTRODUCTION

Coxiella burnetii is the causative agent of Q fever in human and animals and ticks are considered to be the natural primary reservoirs of *C. burnetii* responsible for the spread of the infection in wild animals and for transmission to domestic animals (Norlander, 2000; Pluta et al., 2010). Cattle, sheep and goats are the main sources of human infection (Lang, 1990). Infected animals shed highly stable bacteria in urine, faces, milk and through placental and birth fluids. Infection via inhalation of aerosolized organisms or ingestion of raw milk or fresh dairy products has been reported in humans and animals (Tissot-Dupont and Raoult, 1993).

Infection in animals is mainly sub clinical but has been associated with late abortions, stillbirth, delivery of weak offspring and also infertility (Aitken, 1989). Abortions during coxiellosis epizootics have been described in goats and sheep but abortion in dairy cows is rare, although reproductive disorders and mastitis can occur (To et al., 1998).

In human beings, symptoms are highly variable and about 60% of infections are asymptomatic sero-conversion patients. However, Q fever may lead to serious complications and even death in patients with acute disease. Predominant clinical manifestations are

fever, pneumonia and granulomatous hepatitis for acute cases and endocarditis for chronic cases (Arricau-Bouvery and Rodolakis, 2005; Maurin and Raoult, 1999; Zhang et al., 1998).

Serological methods have been used to detect antibodies to *C. burnetii* (Addo, 1980; Soliman et al., 1992; Thomas et al., 1995; Rodolakis et al., 2007). These assays may not be useful for the diagnosis of acute infection due to the delay in antibody development. Furthermore, it is difficult to discriminate between current and past infection because antibodies often persist after the organisms disappear from the blood (Zhang et al., 1998). PCR assay has become a useful tool for the detection of *C. burnetii* in clinical samples because of the low detection limit and high sensitivity (Zhang et al., 1998; Berri et al., 2003; Öngör et al., 2004; Fretz et al., 2007; Guatteo et al., 2007). This assay has been described as the most sensitive and rapid means to identify shedder animals (Arricau-Bouvery and Rodolakis, 2005).

The objective of the present study was to determine the prevalence rate of *C. burnetii* in bulk milk samples from dairy goat herds in 5 different provinces, Iran using a nested PCR assay.

Table 1. Prevalence of *C. burnetii* in bulk milk samples from dairy goat herds in Fars, Ghom, Kerman, Khuzestan and Yazd provinces, Iran.

Province	No. of herds studied	No. of samples per herd	No. of milk samples	No. (%) of <i>C. burnetii</i> positive samples
Fars	22	2 - 4	60	4 (6.7)
Ghom	10	2 - 3	36	-
Kerman	15	2 - 4	50	-
Khuzestan	24	2 - 4	90	1 (1.1)
Yazd	99	2 - 4	296	6 (2.0)

MATERIALS AND METHODS

Collection of samples

From January to May 2010, a total of 296 goat bulk milk samples were collected from 89 goat breeding farms in Fars, Ghom, Kerman, Khuzestan and Yazd Provinces, Iran (Table 1). The animals which their milk samples collected for this study were clinically healthy and the milk samples showed normal physical characteristics. The samples were immediately transported to the laboratory in a cooler with ice packs and were processed within an hour of collection.

PCR detection of *C. burnetii*

C. burnetii was isolated from milk samples by centrifuging and removing cream and milk layers as described previously by Berri et al. (2003). Purification of DNA was achieved using a genomic DNA purification kit (Fermentas, GmbH, Germany) according to the manufacturer's instruction and the total DNA was measured at 260 nm optical density according to the method described by Sambrook and Russell (2001).

All oligonucleotide primers were obtained from a commercial source (Cinna Gen, Iran). The nested PCR assay used to screen for *C. burnetii* was designed from the nucleotide sequence of the com1 gene encoding a 27KD outer membrane protein (OMP) as previously described (Zhang et al., 1998) and the amplification was carried according to the method described elsewhere (Fretz et al., 2007). For the nested PCR assay with primers OMP1-OMP2 and OMP3-OMP4, the first amplification was performed in a total volume of 25 µl containing 5 µl of DNA sample, 0.5 mM MgCl₂, 0.2 mM (each) dNTPs, 1 µM primer OMP1, 1 µM primer OMP2, and 0.5 U/reaction of Taq DNA polymerase (Roche Applied Science, Germany).

The PCR assay was performed at 94°C for 4 min and then for 30 cycles of 94°C for 1 min, 56°C for 1 min and 72°C for 1 min in a DNA thermal cycler (Master Cycler Gradient, Eppendorf, Germany). In the second amplification, the reaction was performed in a total volume of 25 µl containing 2 µl of DNA sample, 0.5 mM MgCl₂, 0.2 mM (each) dNTPs, 0.8 µM primer OMP3, 0.8 µM primer OMP4 and 0.5 U/reaction of Taq DNA polymerase. The PCR assay was performed at 95°C for 4 min and then for 30 cycles of 94°C for 1 min, 57°C for 1 min, and 72°C for 1 min. The PCR-amplified products (OMP1-OMP2: 501 bp; OMP3-OMP4: 438 bp) were examined by electrophoresis in a 1.5% agarose gel, stained with a 1% solution of ethidium bromide, and examined under UV illumination. In the present study, *C. burnetii* DNA (Serial Number: 3154; Genekam Biotechnology AG, Germany) and DNase free water were used as the positive and negative controls, respectively.

RESULTS AND DISCUSSION

In the present study, a total of 296 bulk milk samples from 89 dairy goat herds in Fars, Ghom, Kerman, Khuzestan and Yazd provinces of Iran were tested for *C. burnetii* using a nested PCR assay. In total, 6 of 296 (2.0%) goat milk samples were positive (Table 1). The positive samples were from 4 of 22 (18.2%) dairy herds in Fars, 1 of 24 (4.2%) dairy herds in Khuzestan and 1 of 18 (5.5%) dairy herds in Yazd. All 76 caprine bulk milk samples from 25 goat breeding farms collected in Ghom and Kerman were negative.

In a recent study in Switzerland, all 81 ovine and 39 goat bulk milk samples were negative for *C. burnetii* using a nested PCR assay (Fretz et al., 2007). In another study conducted in Chaharmahal va Bakhtiari province of Iran, 1.8% of goat bulk milk samples was positive for *C. burnetii* (Rahimi et al., 2009).

Testing animal based on only bulk milk sample can be lead to misclassify the status of the animal because *C. burnetii* may be shed by other routes such as vaginal mucus, feces, urine, placenta, or birth fluids, (Guatteo et al., 2006). It seems goat excrete *C. burnetii* in their vaginal discharges, feces and milk (Rodolakis et al., 2007). Shedding of *C. burnetii* by infected animals occurs mainly during parturition and lactation. Therefore, detection of *C. burnetii* in bulk tank milk greatly depends on the sampling time. The use of repeated sampling can reduce the likelihood of falsely classifying a herd as *C. burnetii* negative (Guatteo et al., 2007).

Our data indicate, although prevalence *C. burnetii* was 2.0% in goat bulk milk samples, clinically healthy dairy goats are important sources of *C. burnetii* infection in Iran. Therefore, in order to prevent the spread of infection in animal and human populations, control goat coxiellosis should be instituted. Although, governmental regulation of milk pasteurization and sanitation in dairy processing plants has been established in Iran for many years, direct sale of unpasteurized milk and dairy products from producers to the consumer is not uncommon in many regions including Fars, Ghom, Kerman, Khuzestan and Yazd provinces. In fact, the consumption of fresh, unpasteurized milk from goat is a traditional practice in some rural areas. The present results also suggest that

testing bulk tank milk as an easy and inexpensive method could be used to assess the efficiency of control schemes aimed at controlling and/or preventing *C. burnetii* infection in dairy herds. Further work is now required to characterize the epidemiology of the infection more thoroughly.

ACKNOWLEDGEMENTS

The authors would like to thank Majed Reahi and Manouchehr Momeni for the sincere help in performing technical parts of the project. We are also grateful to Morteza Moosavian, Hamid Reza Kazemeini, Mazear Rafei, Zahra Emami, Ziba Babadi and Mohamad Derakhshesh for assistance with sampling.

REFERENCES

- Addo PB (1980). A serological survey for evidence of Q fever in camels in Nigeria. *Br. Vet. J.*, 136: 519-521.
- Aitken ID (1998). Clinical aspects and prevention of Q fever in animals. *Eur. J. Epidemiol.*, 5: 420-424.
- Arricau-Bouvery N, Rodolakis A (2005). Is Q fever an emerging or re-emerging zoonosis? *Vet. Res.*, 36: 327-349.
- Berri M, Arricau-Bouvery N, Rodolakis A (2003). PCR-based detection of *Coxiella burnetii* from clinical samples. In: Sachse K., Frey J. (Eds.), *Methods in molecular biology*, Humana Press Inc., Totowa, NJ, pp. 153-161.
- Fretz R, Schaeren W, Tanner M, Baumgartner A (2007). Screening of various foodstuffs for occurrence of *Coxiella burnetii* in Switzerland. *Int. J. Food Microbiol.*, 116: 414-418.
- Guatteo R, Beaudou F, Joly A, Seegers H (2006). Shedding routes of *Coxiella burnetii* in dairy cows: implications for detection and control. *Vet. Res.*, 37: 827-833.
- Guatteo R, Beaudou F, Joly A, Seegers H (2007). Assessing the within-herd prevalence of *Coxiella burnetii* milk-shedder cows using a real-time PCR applied to bulk tank milk. *Zoonoses and Public Health*, 54: 191-194.
- Maurin M, Raoult D (1999). Q fever. *Clin. Microbiol. Rev.*, 12: 518-553.
- Norlander L (2000). Q fever epidemiology and pathogenesis. *Microbes Infect.*, 2: 417-424.
- Öngör H, Cetinkaya B, Karahan M, Nuri Acik M, Bulut H, Muz A (2004). Detection of *Coxiella burnetii* by immunomagnetic separation-PCR in the milk of sheep in Turkey. *Vet. Rec.*, 154: 570-572.
- Pluta S, Hartelt K, Oehme R, Mackenstedt U, Kimmig P (2010). Prevalence of *Coxiella burnetii* and *Rickettsia* spp. in ticks and rodents in southern Germany. *Ticks and Tick-borne Dis.* (In press) doi:10.1016/j.ttbdis.2010.04.001.
- Rahimi E, Doosti A, Ameri M, Kabiri E, Sharifian B (2009). Detection of *Coxiella burnetii* by nested PCR in bulk milk samples from dairy bovine, ovine, and caprine herds in Iran. *Zoonoses and Public Health* (In press) doi: 10.1111/j.1863-2378.2009.01289.
- Rodolakis A, Berri M, Hécharde C, Caudron C, Souriau A, Bodier CC, Blanchard B, Camuset P, Devillechaise P, Natorp JC, Vadet JP, Arricau-Bouvery N (2007). Comparison of *Coxiella burnetii* shedding in milk of dairy bovine, caprine, and ovine herds. *J. Dairy Sci.*, 90: 5352-5360.
- Sambrook J, Russell DW (2001). *Molecular Cloning: A Laboratory Manual*. 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Soliman AK, Botros BA, Watts DM (1992). Evaluation of a competitive enzyme immunoassay for detection of *Coxiella burnetii* antibody in animal sera. *J. Clin. Microbiol.*, 30: 1595-1597.
- Tissot Dupont H, Raoult D (1993). Epidemiologie de la fièvre Q. *Bull. Epidem. Hebd.*, 5: 17-18.
- To H, Htwe K, Kako N, Kim HJ, Yamaguchi T, Fukushi H, Hirai K (1998). Prevalence of *Coxiella burnetii* infection in dairy cattle with reproductive disorders. *J. Vet. Med. Sci.*, 60: 859-861.
- Zhang GQ, Nguyen SV, To H, Ogawa M, Hotta A, Yamaguchi T, Kim HJ, Fukushi H, Hirai K (1998). Clinical evaluation of a new PCR assay for detection of *Coxiella burnetii* in human serum samples. *J. Clin. Microbiol.*, 36: 77-80.