

Full Length Research Paper

The effect of a continuous mercury stress on mercury reducing community of some characterized bacterial strains

Ashraf M. M. Essa

Botany Department, Faculty of Science, Fayoum University, El Fayoum, Egypt. E-mail: ashraf.essa@yahoo.com.

Accepted 29 December, 2011

Active resistance to the mercuric ion is widely distributed in environmental microbes and results from the action of mercuric reductase. Five mercury resistant bacteria: *Escherichia coli* Z1, *Escherichia coli* Z3, *Pseudomonas putida* Z2, *Serratia marcescens* Z4 and *Xanthomonas* sp. Z5 were isolated and identified from sludge sample. The presence of mercury resistance determinants was screened by PCR using *merA*-specific primers. Based on the analysis of *merA* amplicons, high similarity was recorded between the *merA* region of the strains *P. putida* Z2 and *Xanthomonas* sp. Z5 with those of Tn5053; while the *merA* of *E. coli* Z1 was analogous to those of Tn21. In case of the bacterial strains *E. coli* Z3 and *S. marcescens* Z4 a great matching was obtained between their *merA* and those of Tn5036. The effect of mercury stress upon the structure of mercury reducing biofilm at the species level and the type of mercury resistance determinants was studied in a continuous bioreactor. Community analysis suggested that the bacterial strain *E. coli* Z3 containing Tn5036-like determinant is the well adapted strain that tolerated elevated levels of mercury whereas the other strains showed a less fitness under these extreme conditions.

Key words: Mercury resistant bacteria, mercuric reductase gene, PCR-RFLP, mercury stress.

INTRODUCTION

The removal of widespread industrial and agricultural heavy metal contamination is considered a challenge for environment management. Microorganisms in contaminated environment have developed resistance to mercury and are playing a major role in natural decontamination (Cursino et al., 1999). The detoxification of mercury by mercury resistant bacteria offers a potential cheaper and safer alternative to conventional methods; moreover, some mercury resistant bacteria can not only detoxify mercury but also remove other metals such as cadmium and lead (De et al., 2008).

Resistance against mercury has been identified in a wide range of Gram-negative and Gram-positive bacteria in natural and mercury contaminated environments and it is often found on plasmids or other mobile genetic elements such as transposons (Osborn et al., 1997; Narita et al., 2004). Mercury resistance mechanism is based on a group of genes located in a mercury resistant operons which allows bacteria to reduce the toxic Hg(II)

into volatile metallic mercury Hg(0) through its enzymatic reduction (Summers, 1986; Brown et al., 1991; Misra, 1992; Barkay et al., 2003); these operons contain genes encoding the functional proteins for regulation (*merR*), transport (*merT*, *merP*) and reduction (*merA*) in addition to some accessory genes (*merC*, *merF* and *merB*) (Ji and Silver, 1995; Nies, 1999). The bioremediation of mercury from synthetic solution or wastewater via volatilization using natural or immobilized mercury resistant bacterial cells has been described by several investigators (Brunke et al., 1993; von Canstein et al., 1999, 2002; Dzair et al., 2004; Wagner-Dobler et al., 2000).

Microbial biodiversity has become a research subject for understanding engineered ecosystems. Several studies have reported the importance of measuring the microbial diversity in laboratory bioreactors in order to understand the relationship between the composition of the microbial community and operational parameters (Liu et al., 1997; Boon et al., 2002). It is well established that

Table 1. Synthetic oligonucleotide primers used in this study.

Primer	Sequence 5'-3'	Amplified region	Size	Reference
PA 530r	AGAGTTTGCCTGGCTCAG GTATTACCGCGGCTGCTG	Conserved region of 16S rDNA gene	500 bp	Lane et al., 1985
F3 R4	GGGGGCACCTCAGAAAACGGA GGAATCGCGCAGACCTCACCT	IR - <i>merT</i> region of Tn21-like operon	730 bp	Essa et al., 2003
KI KII	GGGGTCGTCTCAGAATTCGG GACAAGCCCTATGGCAGCAT	IR - <i>merR</i> region of Tn5036-like operon	350 bp	Essa et al., 2003
MI3 MI2	GGAGTCGCCTCAGAAAACG TACGGAGTCAAGCGATATGGA	IR - <i>merR</i> region of Tn5053-like operon	500 bp	Essa et al., 2003
MRS1 MRS2	ACCATCGGCGGCACCTGCG AAGGTCTGS*GCCGR*AGCTTC	<i>merA</i> region of Hg ^r operons	1300 bp	Glendinning, 2000

S* = C+G, R* = A+G.

toxic effects of heavy metals are highly selective in microbes; such selective targeting of specific enzymatic systems and pathways suggests that certain members of the microbial community would be more sensitive to heavy metal exposure than others, depending on the sensitivity of their critical metabolic pathways (Fulladosa et al., 2005; Sobolev and Begonia, 2008).

The aim of this study is the use of PCR-based techniques targeting the *merA* gene that codes for mercuric reductase in order to explore the functional diversity of a mercury reducing community under continuous mercury stress.

MATERIALS AND METHODS

Isolation and purification of mercury resistance bacterial strains

Luria Bertani (LB) broth supplemented with 10 µg/ml HgCl₂ was inoculated with sludge sample obtained from the Zenein Waste Water Treatment Plant (ZWWTP) localized in the Giza Governorate, Egypt and incubated at 30°C on shaking incubator (200 rpm) for 48 h followed by pour plate method on LB agar medium. A single bacterial colony was aseptically picked up and transferred onto a fresh medium with a streaking technique and incubated for 24 h at 30°C. Transferring was repeated until obtaining a pure bacteria culture and the isolated colonies were plated on LB agar plates supplemented with 20 µg/ml HgCl₂. Bacterial colonies which showed better growth on HgCl₂ plates were taken and streaked in the LB agar slants and stored.

Total DNA and plasmid preparation

The total bacterial DNA was prepared according to the method of Goldberg and Ohman (1984), the small scale purification of plasmid DNA was performed by the modified alkaline method of Le Gouill et al. (1994).

PCR amplification of DNA encoding the 16S rRNA gene

Amplification of the 16S rDNA gene was carried out by using primer pair pA/530r (Table 1). The PCR mixture was prepared as the following; 10 µl (10x) PCR buffer, 3 µl (50 mM) MgCl₂, 1 µl (20 pmole/µl) of each primer, 1 µl (10 mM) dNTPs mixture, 0.5 µl (2.5U) Taq DNA polymerase, 2 µl total DNA extract, and the volume is completed to 100 µl by SDH₂O. PCR were carried out for 35 cycles under the following conditions: denaturation step at 94°C for 40 s, annealing step at 55°C for 1 min, extension step at 72°C for 2 min and final extension at 72°C for 10 min. An aliquot of the PCR products (10 µl) was mixed with 2 µl of DNA loading buffer and analyzed by electrophoresis (15 V/cm, 60 min) on 0.7% horizontal agarose gel in TBE buffer containing 0.5 µg/ml ethidium bromide, then visualized on an UV transilluminator.

PCR for amplification of *merA* region


The purified plasmid DNA of the mercury resistant strains was used as a template in PCR by using MRS1/MRS2 primers (Table 1) to amplify the *merA* region. The PCR mixture was prepared as described above and PCR were carried out for 35 cycles under the following conditions: denaturation step at 94°C for 40 s, annealing step at 57°C for 1 min, extension step at 72°C for 2 min and final extension at 72°C for 10 min. The PCR products were analyzed as mentioned above.

Purification of the PCR products and nucleotide sequence analysis

Aqueous PCR products were purified by using a QIAquick PCR purification kit as described by the manufacturer's instructions. The purified PCR products were sequenced using ABI PRISM Big Dye Terminator Cycle Sequencing Ready Reaction Kits with Ampli Taq DNA polymerase (CliniLab, Egypt). The sequence data were analysed by BLASTN search at the National Centre for Biotechnology Information (<http://www.ncbi.nlm.nih.gov>) to identify the most similar sequences.

Table 2. The restriction enzymes: *PshAI*, *Accl* and *Eco01091* were used for the RFLP analysis of *merA* amplicons based on their DNA sequence.

<i>merA</i> amplicons	<i>PshAI</i>	<i>Accl</i>	<i>Eco01091</i>
Tn21-like operon	Single cut	Do not cut	Do not cut
Tn5036-like operon	Do not cut	Do not cut	Single cut
Tn5053-like operon	Do not cut	Single cut	Do not cut

**Figure 1.** Gel electrophoresis of PCR products of: A) the partial 16S rDNA gene of the bacterial isolates Z1 (lane 2), Z2 (lane 3), Z3 (lane 4), Z4 (lane 5) and Z5 (lane 6), B) the *merA* gene from plasmid DNA of *Escherichia coli* Z1 (lane 2), *Escherichia coli* Z3 (lane 3), *Pseudomonas putida* Z2 (lane 4), *Xanthomonas* sp. Z5 (lane 5), *Serratia marcescens* Z4 (lane 6). Lane 1 in both figures contains Hyperladder I marker.

PCR-RFLP pattern

According to the DNA sequence and the restriction map of the *merA* regions that were amplified from the bacterial isolates (Restriction Site Analyzer and Map Generator, www.algosome.com), the restriction enzymes: *PshAI*, *Accl* and *Eco01091* (GibcoBRL, Life Technologies) were chosen to digest the *merA* amplicons (Table 2). The reaction was set up as follows; 1.5 µg PCR product, 5 µl restriction enzyme, 10 µl (10x) restriction enzyme buffer, and the volume was completed up to 100µL by sterile distilled water. The reaction was incubated at 37°C for 1 h. After inactivation (65°C for 20 min), the reaction mixture was mixed with 0.2 volume of DNA loading buffer and analyzed by electrophoresis.

PCR with primers specific for the different *mer* determinants

According to the obtained DNA sequence of *merA* region, some primers were used to discriminate between the different *mer* operons (Table 1). The purified plasmid DNA of the mercury resistant strains was used as a template in PCR by using the following primers E3/E4 to identify the Tn5075 operon, MI3/MI2 to identify the Tn5053 operon and KI/KII to identify the Tn5036 operon. The PCR mixture was prepared as described above. PCR were carried out for 35 cycles under the following conditions; denaturation step at 94°C for 40 s, annealing step at 56°C for 1 min, extension step at 72°C for 2 min and final extension at 72°C for 10 min. The PCR products were analyzed as mentioned above.

Bioreactor setup and operation

The mercury resistance bacterial isolates from the sludge sample, which contains different mercury resistant determinants, were

grown individually in LB broth supplemented with 10 µg/ml HgCl₂ at 37°C on a shaking incubator at 200 rpm for 24 h. A 25 ml of each culture were mixed together and used as an inoculum for the bioreactor which contains about 1.5 L LB broth. The bioreactor was maintained under aerobic conditions by pumping in filter-sterilized air, 37°C for 30 days. The LB broth supplemented with HgCl₂ (10 to 60 µg/ml) was flowed through the bioreactor at 100 ml/h and the bacterial growth was monitored by measuring the protein content. Moreover, the DNA extracted from the effluent of the bioreactor during the operating period was subjected to PCR by using specific primers for the different determinants (Table 1). At the end of the experiment (30 days), the community composition at the strain level was analyzed by 16S ribosomal DNA analysis. At the same time, the RFLP technique was used to profile the type of the mercury resistant determinants based on their *merA* genes. The protein content was used to follow the bacterial growth under different HgCl₂ concentrations. Samples from the effluent of the bioreactor were centrifuged for 10 min (10,000 rpm), and cell pellets were re-suspended in 500 µl of NaOH (0.5 M) and lysed for 1 h. The protein content was estimated according to the method of Lowry et al. (1951).

RESULTS AND DISCUSSION

Isolation and characterization of mercury resistant bacteria

The mercury resistant bacterial strains designated Z1, Z2, Z3, Z4 and Z were isolated from the sludge sample (ZWWTP) and were identified by partial 16S ribosomal DNA technique (Figure 1A). The purified PCR products were sequenced and databank compared. The isolates




Figure 2. Gel electrophoresis of RFLP pattern of *merA* amplicons digested with *Eco01091*, *Accl* and *PshAI*. *Xanthomonas* sp. Z5 containing Tn5053-like determinant is represented in lanes (2 to 4), *S.marcescens* Z4 containing Tn5036-like determinant is represented in lanes (5 to 7) and *E. coli* Z1 containing Tn21-like determinant is represented in lanes (8 to 10). Digestion by *Eco01091* is represented in lanes (4, 7 and 10), *Accl* in lanes (3, 6 and 9) and *PshAI* in lanes (2, 5 and 8). Lane 1 contains Hyperladder I marker.

Z1 and Z3 were identified as *Escherichia coli* (99.6 and 99.9% identity, respectively), isolate Z2 was identified as *Pseudomonas putida* (96.7% identity), isolate Z4 was identified as *Serratia marcescens* (99.7% identity), and isolate Z5 was identified as *Xanthomonas* sp. (97.8% identity).

The purified plasmid DNA of each mercury resistant strain was screened by PCR for *merA* genes. Results in Figure 1B demonstrated the presence of the *merA* gene (approximately 1300 bp) in the mercury resistant isolates. The purified PCR products were sequenced and databank compared; the *merA* region of *E. coli* Z1 strain showed a high similarity to those of Tn21 (99%, Liebert et al., 1999). In case of *P. putida* Z2 and *Xanthomonas* sp. Z5 the amplified *merA* region recorded the highest identity to those from Tn5053 (96 and 98%, Kholodii et al., 1995) while, the amplified *merA* region of *E. coli* Z3 and *S. marcescens* Z4, showed the highest identity to those from Tn5036 (96 and 99%, Yurieva et al., 1997).

According to the DNA sequence and the restriction map of the *merA* region of the different determinants (Restriction Site Analyzer and Map Generator, www.algosome.com), some restriction enzymes were used to digest the *merA* amplicons (Table 2) resulting in a specific RFLP pattern (Figure 2). *PshAI* digested the *merA* of Tn21-like determinant into two fragments (760 to 480 bp), *Accl* digested the *merA* of Tn5053-like determinant into two fragments (880 to 360 bp) whereas *Eco01091* digested the *merA* of Tn5036-like determinant into two fragments (790 to 450 bp); moreover, PCR with specific primers (Table 1) was used to confirm the type of these determinants in the bacterial strains. Data in Figure 3 showed that PCR amplicons were obtained by using

primers E3/E4 with Tn21-like determinant (730 bp), MI3/MI2 with Tn5053-like determinant (500 bp), and KI/KII with Tn5036-like determinant (350 bp).

Effect of mercury stress on a mercury reducing biofilm

The isolated mercury resistant strains: *E. coli* Z1 and Z3, *P. putida* Z2, *Xanthomonas* sp. Z5 and *S. marcescens* Z4, were grown together inside a continuous bioreactor for 30 days under selective continuous mercury stress and the total protein content was monitored as a parameter for the bacterial growth (Figure 4A). At the same time, the DNA extracted from the effluent of the bioreactor upon the pilot plant operation was subjected to PCR by using specific primers for the different mercury resistance determinants. The obtained data (Figure 4B) demonstrated that the Tn5053-like determinant disappeared after 18 days (at 40 µg/ml of HgCl₂), the Tn21-like determinant vanished after 24 days (at 60 µg/ml of HgCl₂) whereas the Tn5036-like determinant recorded a high tolerance capability under these extreme conditions.

At the end of the experiment (after 30 days), the composition of the mercury resistant community at the strain level was analyzed on the basis of the 16S ribosomal DNA gene showing the presence of only *E. coli* (Z3), meanwhile the other strains were completely gone. The use of 16S rRNA gene as a marker to study the composition and the dynamics of some bacterial communities has been reported in previous studies (Wagner-Dobler et al., 2000; Saikaly et al., 2005).




Figure 3. Gel electrophoresis of PCR products from plasmid DNA of *E. coli* Z1 containing Tn21-like determinant by using primers E3/E4 (lane 2), *Xanthomonas* sp. Z5 containing Tn5053-like determinant by using primers MI3/MI2 (lane 3) and *S. marcescens* Z4 containing Tn5036-like determinant by using primers KI/KII (lane 4). Lane 1 contains the hyperladder I marker.





Figure 4. (A) The growth of a mercury resistance bacterial population (expressed as mg/mL protein) consisting of *E. coli* Z1, *E. Z3*, *P. putida* Z2, *Xanthomonas* sp. Z5 and *S. marcescens* Z4 in a continuous aerobic bioreactor for 30 days on LB broth supplemented with different HgCl_2 concentrations: 20 $\mu\text{g}/\text{mL}$ up to the 3rd day, 30 $\mu\text{g}/\text{mL}$ up to the 9th day, 40 $\mu\text{g}/\text{mL}$ up to the 15th day, 50 $\mu\text{g}/\text{mL}$ up to the 21st day, 60 $\mu\text{g}/\text{mL}$ up to the end of the operation. (B) The gel electrophoresis of PCR products from plasmid DNA of the bioreactor effluent obtained at the start of the experiment (lane 1), at 6 days (lane 2), at 12 days (lane 3), at 18 days (lane 4), at 24 days (lane 5), at 30 days (lane 6) by using primers KI/KII for Tn5036-like determinant, MI3/MI2 for Tn5053-like determinant and E3/E4 for Tn21-like determinant. Lane 1 contains Hyperladder I marker.




Figure 5. The gel electrophoresis of: A) RFLP pattern of *merA* amplicons obtained from plasmid DNA of the bioreactor effluent at the end of the experiment (30 days) digested with *Eco*01091 (lane 2), *Acc*I (lane 3) and *Psh*AI (lane 4), B) the PCR products from plasmid DNA of the bioreactor effluent obtained after 30 days by using primers KI/KII (lane 2), MI3/MI2 (lane 3) and E3/E4. Lane 1 contains Hyperladder I marker.

Actually, the alteration of a bacterial community in laboratory bioreactors under the influence of the heavy metals stress by targeting of some marker genes especially those responsible for the resistance mechanism will produce more accurate image for the biodiversity of the bacterial population; so the mercury resistant community was analysed to profile the type of their determinant through using RFLP analysis of the obtained *merA* amplicons which showed the presence of Tn5036-determinant while the other determinants were not found (Figure 5A). These results were confirmed via subjecting the DNA extracted from the effluent of the bioreactor after 30 days to PCR by using some specific primers for the different *mer* determinants (Figure 5B). A clear DNA band (350 bp) was obtained with primers MR3/MR2 that are specific to Tn5036-like determinant whereas no PCR products were obtained by using the primers of the other determinants. These results are compatible with those who used the *merA* gene as a molecular marker to follow the assortment of mercury resistant bacterial population under the pressure of mercury toxicity in aerobic and anaerobic environments (Felske et al., 2003; Simbahan et al., 2005; Sotero-Martins et al., 2008).

This study clarified the presence of a strong selective pressure on the microbial community inside the bioreactor due to the mercury toxicity which led to the predominance of *E. coli* (Z3) containing Tn5036-like determinant. These results are in accordance with other studies which showed that the continuous exposure to elevated levels of mercury altered the microbial community and exclusively select bacteria that can cope with such levels (Osborn et al., 1993; Muller et al., 2001; Ramaiah and De, 2003).

The domination of *E. coli* Z3 containing Tn5036-like determinant over the other strains under the strong selective pressure exerted by mercury toxicity was attributed to the well adaptation of this strain which might be linked with the type of mercury resistance determinant. This assumption is consistent with the finding of previous studies that correlated the functional diversity and the adaptation ability of some bacterial communities

under the influence of mercury with the frequency and the type of mercury resistance operon (Smalla et al., 2006; Chien et al., 2010); moreover, the capability of *E. coli* Z3 to tolerate an elevated level of mercury could be based on the presence of some additional mechanisms beside mercury volatilization that give this organism an extra-tolerance capability to cope with such mercury stress. Agreeing with this hypothesis, Haferburg and Kothe (2007) reported that the adaptation to heavy metal rich environments resulted in microorganisms which show activities for biosorption, bioprecipitation, extracellular sequestration and chelation. Such resistance mechanisms may play a role in transforming the toxic metals into other forms that are not biologically available to the cells. One of these mechanisms is the precipitation of the soluble metal ions away from the cells via its complexation into insoluble metal precipitates via the production of some metabolites (Essa et al., 2006).

Finally, an environment with a raised concentration of heavy metals constitutes a prospective stimulus for toxic metal tolerant bacteria; such polluted environments encourage adaptation for heavy metal resistance and markedly affect on the composition of a bacterial biofilm. The use of *merA* gene, the key enzyme of mercury volatilization, as molecular markers in order to follow the diversity of mercury resistant bacterial population under the pressure of mercury toxicity can provide significant information about the functional alterations of these communities especially in contaminated environments. Despite of the intrinsic role of the different *mer* operons in mercury detoxification process, work is still necessary to illustrate the distribution and diversity of these genetic determinants in the bacterial communities under heavy metals stress in order to employ them for the bioremediation of these toxic pollutants.

REFERENCES

- Barkay T, Miller SM, Summers AO (2003). Bacterial mercury resistance from atoms to ecosystems. *FEMS Microbiol. Rev.*, 27: 355–384.
 Boon N, Windt WD, Verstraete W, Top EM (2002). Evaluation of nested

- PCR-DGGE (denaturing gradient gel electrophoresis) with group specific 16S rRNA primers for the analysis of bacterial communities from different wastewater treatment plants. *FEMS Microbiol. Ecol.*, 39: 101-112.
- Brown NL, Camakaris J, Lee BT, Williams T, Morby AP, Parkhill J, Rouch DA (1991). Bacterial resistances to mercury and copper. *J. Cell Biochem.*, 46: 106-114.
- Brunke M, Deckwer WD, Frischmuth A, Horn JM, Lunsdorf H, Rhode M, Rohricht M, Timmis KN, Weppen P (1993). Microbial retention of mercury from waste streams in a laboratory column containing *merA* gene bacteria. *FEMS Microbiol. Rev.*, 11: 145-152.
- Chien M, Lin MK, Chang J, Huang C, Endo G, Suzuki S (2010). Distribution of mercury resistance determinants in a highly mercury polluted area in Taiwan. *Interdisciplinary Studies on Environmental Chemistry - Biological Responses to Contaminants*, Eds., pp. 31-36.
- Cursino L, Oberda SM, Cecilio RV, Moreira RM, Chartone-Souza E, Nascimento AMA (1999). Mercury concentration in the sediment at different gold prospecting sites along the Carmo stream, Minas Gerais, Brazil & frequency of resistant bacteria in the respective aquatic communities. *Hydrobiologia*, 394: 5-12.
- De J, Ramaiah N, Vardanyan L (2008). Detoxification of toxic heavy metals by marine bacteria highly resistant to mercury. *Mar. Biotechnol.*, 10(4): 471-477.
- Dzair FZ, Zeroual Y, Moutaouakkil A, Taoufik J, Talbi M, Loutfi M, Lee K, Blaghen M (2004). Bacterial volatilization of mercury by immobilized bacteria in fixed and fluidized bed bioreactor. *Ann. Microbiol.*, 54: 353-364.
- Essa AM, Julian DJ, Kidd SP, Brown NL, Hobman JL (2003). Mercury resistance determinants related to Tn21, Tn1696, and Tn5053 in enterobacteria from the preantibiotic era. *Antimicrob. Agents Chemother.*, 47: 1115-1119.
- Essa AM, Creamer NJ, Brown NL, Macaskie LE (2006). A new approach to the remediation of heavy metal liquid wastes via off-gases produced by *Klebsiella pneumoniae* M426. *Biotechnol. Bioeng.*, 95(6): 576-583.
- Felske ADM, Fehr W, Pauling BV, von Canstein H, Wagner-Döbler I (2003). Functional profiling of mercuric reductase (*merA*) genes in biofilm communities of a technical scale biocatalyzer. *BMC Microbiol.*, 3: 22-27.
- Fulladosa E, Murat JC, Villaescusa I (2005). Study on the toxicity of binary equitoxic mixtures of metals using the luminescent bacteria *Vibrio fischeri* as a biological target. *Chemosphere*, 58: 551-557.
- Glendinning KG (2000). Studies on mercuric reductase and haemophilic mercury resistance. Ph.D. thesis, the University of Birmingham, UK.
- Goldberg JB, Ohman DE (1984). Cloning and expression in *Pseudomonas aeruginosa* of a gene involved in production of alginate. *J. Bacteriol.*, 158: 1115-1121.
- Haferburg G, Kothe E (2007). Microbes and metals: interactions in the environment. *J. Basic Microbiol.*, 47: 453-467.
- Ji G, Silver S (1995). Bacterial resistance mechanisms for heavy metals of environmental concern. *J. Ind. Microbiol.*, 14: 61-75.
- Kholodii GY, Mindlin SZ, Bass IA, Yurieva OV, Minakhina SV, Nikiforov VG (1995). Four genes, two ends, and a *res* region are involved in the transposition of Tn5053: a paradigm for a novel family of transposons carrying either a *mer* operon or an integron. *Mol. Microbiol.*, 17: 1189-1200.
- Lane DJ, Pace B, Olsen GL, Stahl DA, Pace NR (1985). Rapid determination of 16S rRNA sequence for phylogenetic analysis. *Proc. Natl. Acad. Sci. USA*, 82: 6955-6959.
- Le Gouill C, Parent JL, Rola-Pleszczynski M, Stankova J (1994). Analysis of recombinant plasmids by a modified alkaline lysis method. *Anal. Biochem.*, 219: 164-169.
- Liu WT, Marsh TL, Cheng H, Forney LJ (1997). Characterization of microbial diversity by determining terminal restriction fragment length polymorphisms of genes encoding 16S rRNA. *Appl. Environ. Microbiol.*, 63: 4516-4522.
- Liebert CA, Hall RM, Summers AO (1999). Transposon Tn21, flagship of the floating genome. *Microbiol. Mol. Biol. Rev.*, 63: 507-522.
- Lowry OH, Rosebrough AL, Farr AL, Randall RJ (1951). Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, 193: 265-275.
- Misra TK (1992). Bacterial resistances to inorganic mercury salts and organo-mercurials. *Plasmid*, 27: 4-16.
- Muller AK, Rasmussen LD, Sorensen SJ (2001). Adaptation of the bacterial community to mercury contamination. *FEMS Microbiol. Lett.*, 16: 49-53.
- Narita M, Matsui K, Huang CC, Kawabata Z, Endo G (2004). Dissemination of TnMER11-like mercury resistance transposons among *Bacillus* isolated from worldwide environmental samples. *FEMS Microbiol. Ecol.*, 48: 47-55.
- Nies NH (1999). Microbial heavy-metal resistance. *Appl. Microbiol. Biotechnol.*, 51: 730-750.
- Osborn AM, Bruce KD, Strike P, Ritchie DA (1993). Polymerase Chain Reaction-restriction fragment length polymorphism analysis shows divergence among *mer* determinants from Gram-negative soil bacteria indistinguishable by DNA-DNA hybridization. *Appl. Environ. Microbiol.*, 59: 4024-4030.
- Osborn AM, Bruce KD, Strike P, Ritchie DA (1997). Distribution, diversity and evolution of the bacterial mercury resistance (*mer*) operon. *FEMS Microbiol. Rev.*, 19: 239-262.
- Ramaiah N, De J (2003). Unusual rise in mercury-resistant bacteria in coastal environs. *Microbiol. Ecol.*, 45: 444-454.
- Saikaly PE, Stroot PG, Oerther DB (2005). Use of 16S rRNA gene terminal restriction fragment analysis to assess the impact of solids retention time on the bacterial diversity of activated sludge. *Appl. Environ. Microbiol.*, 71(10): 5814-5822.
- Simbahan J, Kurth E, Schelert J, Dillman A, Moriyama E, Jovanovich S, Blum P (2005). Community analysis of a mercury hot spring supports occurrence of domain-specific forms of mercuric reductase. *Appl. Environ. Microbiol.*, 71(12): 8836-8845.
- Smalla K, Haines AS, Jones K, Krogerrecklenfort E, Heuer H, Schloter M, Thomas CM (2006). Increased abundance of IncP-1 plasmids and mercury resistance genes in mercury-polluted river sediments: First discovery of IncP-1 plasmids with a complex *mer* transposon as the sole accessory element. *Appl. Environ. Microbiol.*, 72(11): 7253-7259.
- Sobolev D, Begonia MFT (2008). Effects of heavy metal contamination upon soil microbes: lead-induced changes in general and denitrifying microbial communities as evidenced by molecular markers. *Int. J. Environ. Res. Pub. Heal.*, 5: 450-456.
- Sotero-Martins A, de Jesus MS, Lacerda M, Moreira JC, Lauria Filgueiras AL, Guimarães Barrocas PR (2008). A conservative region of the mercury reductase gene (*merA*) as a molecular marker of bacterial mercury resistance. *Braz. J. Microbiol.*, 39: 307-310.
- Summers AO (1986). Organization, expression and evolution of genes for mercury resistance. *Annu. Rev. Microbiol.*, 40: 607-634.
- von Canstein H, Kelly S, Li Y, Wagner-Döbler I (2002). Species diversity improves the efficiency of mercury-reducing biofilms under changing environmental conditions. *Appl. Environ. Microbiol.*, 68(6): 2829-2837.
- von Canstein H, Li Y, Timmis KN, Deckwer WD, Wagner-Döbler I (1999). Removal of mercury from chloralkali electrolysis wastewater by a mercury-resistant *Pseudomonas putida* strain. *Appl. Environ. Microbiol.*, 65(12): 5279-5284.
- Wagner-Döbler I, von Canstein H, Li Y, Timmis KN, Deckwer WD (2000). Removal of mercury from chemical wastewater by microorganisms in technical scale. *Environ. Sci. Technol.*, 34: 4628-4634.
- Yurieva O, Kholodii G, Minakhin L, Gorlenko Z, Kalyaeva E, Mindlin S, Nikiforov V (1997). Intercontinental spread of promiscuous mercury resistance transposons in environmental bacteria. *Mol. Microbiol.*, 24: 321-329.