

Short Communication

Endophytic *Fusarium* spp. from wild banana (*Musa acuminata*) roots

Latiffah Zakaria^{1,2*} and Nur Hidayah Abdul Rahman¹

¹School of Biological Sciences, Universiti Sains Malaysia, 11800 USM, Pulau Pinang, Malaysia.

²Centre of Marine and Coastal Studies, Universiti Sains Malaysia (CEMACS-Mukahead), 11800 USM, Pulau Pinang, Malaysia.

Accepted 23 August, 2011

Endophytic *Fusarium* species were isolated from roots of wild banana (*Musa acuminata*) collected randomly in several locations in Penang Island, Malaysia. A total of 54 isolates of *Fusarium* were recovered from 100 root fragments. Based on morphological features of macroconidia, microconidia and conidiogenous cells, three *Fusarium* species were identified in which the most common species was *F. oxysporum* (41.5%) followed by *F. solani* (32.1%) and *F. semitectum* (24.5%). The present study showed that there are diverse endophytic *Fusarium* species in the roots of wild banana and has the potential to be used as biological control agents.

Key words: Endophyte, *Fusarium*, root, wild banana.

INTRODUCTION

Endophytic fungi live inside healthy plant tissues without causing any diseases or apparent damages to the host which indicated that many endophytic fungi form mutually beneficial relationships with their host plant (Petrini et al., 1992; Lodge et al., 1996). In the mutualistic association, the host plants are protected from pest and pathogenic fungi and in return, the host plant provided nutrition to the fungi (Azevedo et al., 2000; Saikkonen et al., 2004). Other definition of endophytic fungi includes symptomless latent pathogen which inhabiting plant tissues at some time in their life cycle without causing apparent harm to the host (Petrini, 1991).

Endophytic *Fusarium* species have been recovered from many types of plants such as Araceae, Bromeliaceae, Orchidaceae (Petrini and Dreyfuss, 1981), Palmae (Rodrigues, 1994) and Compositae (Fisher et al., 1995) including Musaceae. Therefore, endophytic *Fusarium* species are abundant and distributed widely in different types of plant.

The present study was conducted to isolate and to identify endophytic *Fusarium* species from roots of wild banana (*Musa acuminata*). We wanted to gather information on the occurrence and diversity of *Fusarium*

species as endophytic species of *Fusarium* is less studied.

MATERIALS AND METHODS

Isolation and identification of Endophytic *Fusarium*

Symptomless wild banana (*Musa acuminata*) roots were randomly sampled along Paya Terubong and Balik Pulau roads, Penang Island, Malaysia. The roots were collected from nine young and healthy wild banana trees, kept in plastic bags and brought to the laboratory for further analysis. The roots were then washed under running tap water for 24 h to thoroughly clean the root samples.

The roots were cut about 1 to 1.5 cm fragments and surface sterilized in 75% ethanol for 1 min followed by 1% sodium hypochlorite for 3 min and 95% ethanol for 5 min. The samples were then dried using sterilized filter paper.

For isolation of endophytic *Fusarium* isolates, 100 root fragments were plated onto peptone chloro-nitro benzene (PCNB) medium, a semi-selective medium for isolation of *Fusarium* species. The plates were incubated at room temperature (25±1°C) until visible growth of mycelium from the root fragments was observed. The mycelium was subsequently sub-cultured onto fresh PDA plates to obtain pure culture.

For identification, mycelium was grown on carnation leaf-piece agar (CLA) and microscopic and macroscopic structures such as size and shape of microconidia, shape, size, number of septa, shape of apical and basal cells of macroconidia, conidiogenous cells (monophialide and polyphialide) and presence of chlamydospores were observed (Summerell et al., 2003). Species descriptions in The *Fusarium* Laboratory Manual (Leslie and

*Corresponding author. E-mail: Lfah@usm.my. Tel: 04 – 6533506. Fax: 04 – 6565125.

Summerell, 2006) were used for identification of the endophytic *Fusarium* spp.

The efficiency of surface sterilization method for isolation of endophytic fungi was determined for each tissues using imprint method as described by Schulz et al. (1993).

RESULTS AND DISCUSSION

Fifty-four isolates of endophytic *Fusarium* isolates were recovered from the root fragments. No fungal growth was observed on the imprint plates which showed the effectiveness of the surface-sterilization method. Based on morphological characteristics, three *Fusarium* species were identified namely, *F. oxysporum* (41.5%) followed by *F. solani* (32.1%) and *F. semitectum* (24.5%).

Isolation of endophytic *Fusarium* species from wild banana was reported by Photita et al. (2001) and Cao et al. (2002). Three *Fusarium* species designated as *Fusarium* sp. 1, *Fusarium* sp. 2 and *Fusarium* sp. 4 were isolated from young and old leaf tissues of wild banana in Thailand (Photita et al., 2001). *Fusarium* sp. was among the fungal flora isolated from roots of wild banana in South China, comprising 10% of total endophytic fungi recovered from healthy root tissues (Cao et al., 2002). In addition to root tissues, *Fusarium* sp. has been isolated from banana leaves of mature wild banana from different localities in Sao Paulo, Brazil (Pereira et al., 1993).

In the present study, the most common species recovered from the root tissues of wild banana was *F. oxysporum* followed by *F. solani*. Both species have been isolated from healthy roots of Cavendish banana with *F. oxysporum* as the most dominant species recovered from the roots (Athman, 2006). *F. solani* was also recovered from healthy leaf pieces of *Musa* spp. in Hong Kong and south east Queensland (Brown et al., 1998).

In addition to *F. oxysporum* and *F. solani*, *F. semitectum* was also isolated from healthy roots of wild banana. Isolates of *F. semitectum* has been reported to be the most dominant *Fusarium* species recovered from healthy rhizomes of banana CV Pisang Awak (Niere, 2001).

Fusarium oxysporum and *F. solani* are well-known plant pathogens, infecting wide range of crops including banana plants. *Fusarium oxysporum* f.sp. *cubense* is the causal agent of Fusarium wilt or Panama disease and *F. solani* (teleomorph *Nectria haematococca*) caused root rot of banana. *Fusarium semitectum* is associated with causing crown rot of banana (Marin et al., 1996; Knight et al., 2008). In the present study, it is not possible to suggest that the endophytic *Fusarium* species from the wild banana roots are pathogenic or non-pathogenic. Pathogenicity tests are required to confirm pathogenicity of the species on banana plants.

Fungal endophytes have been proposed as potential biological control agents (Brown et al., 1998). Endophytic *F. oxysporum* isolates have been tested to suppress growth of *F. oxysporum* f.sp. *cubense* (Nel et al., 2006). In another study, endophytic *F. oxysporum* reduced

nematode numbers and weevil damage on banana and enhance banana plant growth in the presence of nematode infestation (Paparú et al., 2009). Endophytic *F. semitectum* has been used as biocontrol of ergot on pearl millet by reducing sclerotia formation and development (Rao and Thakur 1988). McFarlane and Rutherford (2005) reported that based on *in-vitro* study, endophytic *F. sacchari* from sugarcane inhibit the development of sugarcane borer moth. Thus, the potential of using endophytic *Fusarium* species as biological control agents of pest and diseases is promising and requires further studies.

The present study demonstrated that similar with other studies, endophytic *Fusarium* species were recovered from roots of wild banana and three *Fusarium* species were identified namely, *F. oxysporum* (41.5%) followed by *F. solani* (32.1%) and *F. semitectum* (24.5%).

REFERENCES

- Athman SY (2006). Genetic diversity of endophytic *Fusarium* species associated with Cavendish banana in South Africa. PhD thesis. University of Pretoria, South Africa.
- Azevedo (Jr.) JL, Walter M, Pereira JV, Araujo WL (2000). Endophytic microorganisms : a review on insect control and recent advances on tropical plants. *Electron J. Biotechnol.*, 3: 40 – 65.
- Brown KB, Hyde KD, Guest DI (1998). Preliminary studies on endophytic fungal communities of *Musa acuminata* species complex in Hong Kong and Australia. *Fungal Divers*, 1: 27 – 25.
- Cao LX, Yon JL, Zhao SN (2002). Endophyte fungi from *Musa acuminata* leaves and roots in South China. *World J. Microbiol. Biotechnol.*, 18: 169 – 171.
- Fisher PJ, Petrini LE, Sutton BC, Petrini O (1995). A study of fungal endophytes in leaves, stems and root of *Gynoxix oleifolia* Muchler (Compositae) from Ecuador. *Nova Hedwigia*, 60: 589 – 594.
- Knight C, Cutts DF, Colhoun J (2008). The role of *Fusarium semitectum* in causing crown rot of banana. *J. Phytopath.*, 89: 170 – 176.
- Leslie JF, Summerell BA (2006). *The Fusarium Laboratory Manual*. Blackwell Publishing Ltd, Iowa.
- Lodge DJ, Fisher PJ, Sutton BC (1996). Endophytic fungi of *Manilkara bidentata* leaves in Puerto Rico. *Mycologia*, 88: 733 – 738.
- Marin DH, Sutton TB, Blankenship M, Swallow WH (1996). Pathogenicity of fungi associated with crown rot of banana in Latin America on Grande naine and disease-resistant hybrid banana. *Plant Dis.*, 80: 525-528.
- McFarlane SA, Rutherford SR (2005). *Fusarium* species isolated from sugarcane in KwaZulu-Natal, and their effect on *Eldana saccharina* (Lepidoptera: Pyralidae) development *in vitro*. *Proc. S. Afr. Sugarcane Technol. Ass.*, 79: 120-123.
- Nel B, Steinberg C, Labuschagne N, Viljoen A (2006). The potential of non-pathogenic *Fusarium oxysporum* and other biological control organisms for suppressing Fusarium wilt of banana. *Plant Pathol.*, 55: 217-223.
- Niere B (2001). Significance of non-pathogenic isolates of *Fusarium oxysporum* Schlecht: Fries for the biological control of the burrowing nematode *Radopholus similis* (Cobb) Thorne on tissue cultured banana. PhD Thesis, University of Bonn, Bonn, Germany.
- Paparú P, Dubois T, Coyne D, Viljoen A (2009). Dual inoculation of *Fusarium oxysporum* endophytes in banana: effect on plant colonization, growth and control of the root burrowing nematode and the banana weevil. *Biocontrol Sci. Technol.*, 19: 639 – 655.
- Pereira JO, Azevedo JL, Petrini O (1993). Endophytic fungi of *Stylosanthes*: a first report. *Mycologia*, 85: 362 – 364.
- Petrini O, Dreyfuss MM (1981). Endophytische Pilze epiphytischen Araceae, Bromeliaceae und Prchidaceae. *Sydowia* 34 : 135 – 148.
- Petrini O, Sieber TN, Toti L, Viret O (1992). Ecology, metabolite production and substrate utilization in endophytic fungi. *Nat. Toxins*,

1: 185 – 196.

Petrini O (1991). Fungal endophytes in tree leaves. In: Andrews JH, Hirano SS (eds). Microbial Ecology of Leaves. Springer, New York. pp. 179 – 197.

Photita W, Lumyang S, Lumyang P, Hyde KD (2001). Endophytic fungi of wild banana (*Musa acuminata*) at Doi Suthep Pui National Park, Thailand. Mycol. Res., 105: 1508 – 1513.

Rao VP, Thakur RP (1988). *Fusarium semitectum* var. *majus* a potential biocontrol agent of argot (*Claviceps fusiformis*) of pearl millet. Indian Phytopathol., 41: 567-574.

Rodrigues KF (1994). The foliar fungal endophytes of the Amazonian palm *Euterpe oleracea*. Mycologia, 86: 376 – 385.

Saikkonen K, Wali P, Helander M, Faeth SH (2004). Evolution of endophyte-plant symbioses. Trends Plant Sci., 9: 275 – 280.

Schulz B, Wanke U, Draeger S, Aust HJ (1993). Endophytes from herbaceous plants and shrubs : effectiveness of surface sterilization. Mycol. Res., 97: 1447 – 1450.

Summerell BA, Salleh B, Leslie JF (2003). A utilitarian approach to *Fusarium* identification. Plant Dis., 87: 117–128.