

## Full Length Research Paper

# Multiple-locus variable-number tandem-repeat analysis of *Bacillus anthracis* isolated from a human-animal anthrax outbreak in the Luangwa valley of Zambia

Kalumbi Malama Mox<sup>1</sup>, Hang'ombe Bernard Mudenda<sup>2\*</sup>, Mulenga Evans<sup>2</sup>, Moonga Ladslav<sup>2</sup>, Muwowo Sergio<sup>3</sup>, Squarre D<sup>4</sup> and Mwansa James C. L.<sup>1</sup>

<sup>1</sup>Department of Pathology and Microbiology, University Teaching Hospital, Lusaka, Zambia.

<sup>2</sup>Microbiology Laboratory, Department of Paraclinical Studies, School of Veterinary Medicine, University of Zambia, Lusaka, Zambia.

<sup>3</sup>Chama District Health Management Team, Chama, Zambia.

<sup>4</sup>Zambia Wildlife Authority, Private Bag 1, Chilanga, Zambia.

Received 23 December, 2013; Accepted 21 February, 2014

The incidence of anthrax, caused by *Bacillus anthracis*, in human and animal population of Zambia has increased recently. In this study, 34 strains of *Bacillus anthracis* from soil, hippopotamuses and humans, isolated in the 2011 outbreak were analyzed using the multiple-locus variable-number tandem repeat analysis. The analysis revealed that a single anthrax clone may have been involved in the epidemic. Considering the cyclical nature of *B. anthracis*, a link could be established with anthrax spores in soil getting ingested by hippopotamuses, followed by human contact resulting into an animal-human epidemic. These data confirm the importance of molecular typing methods for in-depth epidemiological analyses of anthrax epidemics.

**Key words:** *Bacillus anthracis*, hippopotamuses, variable number tandem repeat (VNTRs), epidemic, human.

## INTRODUCTION

*Bacillus anthracis* is a zoonotic disease-causing agent for anthrax. It is a spore forming bacterium that causes disease in wild and domestic animals (Koehler, 2002; Habrun et al., 2011). Animals especially herbivores are mostly infected by oral ingestion of soil contaminated with anthrax spores, while humans are infected through contact with contaminated animal products (Spencer, 2003). In humans, the bacterium causes three major types of anthrax namely inhalational, gastrointestinal and

cutaneous (Baykam et al., 2009).

*B. anthracis* has a worldwide dissemination as it is easy to propagate and has a history of being used as an agent of bioterrorism (Gierczynski et al., 2009). With this background, it is therefore important that the bacteria be identified and typed accurately in order to determine microbial forensics associated with disease and act appropriately in controlling the infection (Koehler, 2002). To enhance such an understanding, the phenotypic and

\*Corresponding author. E-mail: mudenda68@yahoo.com or bhang'ombe@unza.zm. Tel: +260-977-326288. Fax: +260-211-293727.

genetic properties of isolates from one source to the other must be determined. *B. anthracis* is highly monomorphic in nature and differentiation of isolates from different regions has proved to be difficult (Brodzik and Francoeur, 2011). However attempts have been made to develop techniques that can resolve the minor genetic differences that can be found in *B. anthracis*. Genetic differentiation among very closely related individuals such as *B. anthracis* requires the use of molecular markers that exhibit very high diversity (Keim et al., 2004). These techniques include; multiple-locus variable-number tandem repeat analysis (MLVA), amplified fragment length polymorphism (AFLP), single nucleotide repeats (SNR) and single nucleotide polymorphism (SNP) (Vos et al., 1995; Okinaka et al., 2008). SNP detection is facilitated by whole genome discovery approaches which are useful for identifying long branches or key phylogenetic positions (Keim et al., 2004) while AFLP has good resolution analysis that require the use of MLVA. The MLVA is the most suitable technique to differentiate and subtype *B. anthracis* strains (Lindstedt, 2005). One powerful feature of the MLVA system is the ability to simultaneously employ multiple variable number tandem repeat (VNTRs) markers that exhibit varying levels of diversity and, therefore, high resolving power. In this study, we analyzed and compared isolates of *B. anthracis* from wildlife (hippopotamus), soil and humans reported in a human-animal anthrax outbreak in Zambia (Hang'ombe et al., 2012). Anthrax outbreaks have been recorded in Zambia, since 1989 (Turnbull et al., 1991; Turnbull, 1998; Tuchili et al., 1993; Siamudaala et al., 2006) without any report of direct molecular epidemiological linkage of *B. anthracis* strains from an epidemic scenario in Zambia.

## MATERIALS AND METHODS

In this study, we analyzed 34 *B. anthracis* strains associated with a single anthrax epidemic in wildlife and human populations. A vaccine Sterne strain was used as a positive control and was compared with the epidemic isolates. Each *B. anthracis* strain was streaked onto 5% sheep blood agar plates and then incubated at 37°C for 24 h. Genotyping was done as previously described (Keim et al., 2004). The cultured isolates were inactivated by heating at 95°C for 30 min after which DNA was extracted using a DNA extraction kit (Zymo Research, Irvine, CA, USA) according to the manufacturer's instructions. Following DNA extraction, a panel of 21 VNTR markers was used in this analysis (Table 1). These were variable repeat region (*vrr*) markers; *vrrA*, *vrrB1*, *vrrB2*, *vrrC1*, *vrrC2*, CG3, pXO1-*aat*, (Schupp et al., 2000) and Ceb-Bams (CB); CB-1, CB-3, CB-5, CB-7, CB-13, CB-15, CB-21, CB-22, CB-23, CB-24, CB-25, CB-28, CB-30 and CB-31 (Le Fleche et al., 2001). The PCR amplification was performed using the Phusion<sup>TM</sup> flash high fidelity PCR master mix (Finnzymes Oy, Finland). The reactions were performed in a final volume of 10 µl containing 5 µl Phusion flash PCR master mix, 0.5 µM of primer sets in 1 µl volume of each and 2 µl of PCR water. The Piko<sup>TM</sup> thermal cycler (Finnzymes Instruments Oy, Finland) was programmed at 95°C for 10 s for initial denaturation, followed by 35 cycles consisting of 95°C for 1 second, 58°C (changed according to the primers (Table 1)) for 5 s and 72°C for 15 s. Final extension was given 72°C for 1 minute. The thermal PCR-profile was used for all the VNTR markers. The

amplicons were visualized on 1.5% agarose gel (MP Biomedicals, Eschwege, Germany), stained with ethidium bromide and evaluated under UV transilluminator. The estimation of the sizes of PCR products was done according to the migration pattern of a 100-bp DNA ladder.

## RESULTS AND DISCUSSION

In this study, we used the MLVA genotyping comprising 21 VNTR markers to understand the relationship of 34 strains of *B. anthracis* isolated from soil (13 strains), hippopotamuses (16 strains) and human (5 strains) during an anthrax epidemic. Of these isolates, no difference was observed on a panel of 21 VNTR markers. The positive control isolate which is the vaccine strain (Sterne strain) had an observable difference with the outbreak isolates. The difference was observed with the CG3 marker where the Sterne vaccine was positive for CG3, while the outbreak strains were negative (Table 2). Further differences were noted with other markers which amplified a different molecular weight amplicon. The markers with different weight amplicons when compared with the Sterne vaccine strain were, CB-1, CB-3, CB-15, CB-21, CB-22, CB-23, CB-30 and CB-31. This MLVA analysis conducted on the isolates from human, soil and hippopotamus revealed relatedness using 21 VNTRs. The results indicated that all 34 isolates belonged to a single clone. The outbreak strains had the pXO1 and pXO2 plasmids required for virulence. Their presence indicates strain pathogenicity (Keim et al., 1997; Jackson et al., 1997; Habrun et al., 2011) as opposed to the vaccine strain which is negative for the pXO2 plasmid as expected. Analysis of other variable repeats found within *B. anthracis* provides a valuable tool to understand the epidemiology of anthrax outbreaks (Jackson et al., 1997). In this study, the confirmation through MLVA, indicates the outbreak may have been due to a single clone. The results may also indicate that changes from one host to another as a result of genetic instability at the locus during the culture or infection process are rare. Such sequence stability suggests the utility of the VNTR regions as markers to determine the source of an infection or outbreak.

## Conclusion

The cyclical nature of anthrax outbreaks determined by climatic factors such as extreme weather changes like drought followed by heavy rains, may lead to generation of high anthrax spore concentration causing disease in grazing animals such as hippopotamus which are bulky grazers. This outbreak indicated a single clonal source transmission, probably from soil to hippopotamuses and finally to humans through consumption of contaminated meat. The data presented here are limited and therefore call for continuous monitoring of human and animal clinical isolates and their environment (including soil) to

**Table 1.** Primers used in this study for MLVA analysis.

VNTR locus	Primer sequences (5' to 3')	Annealing temperature (°C)
vrrA	F: CACAACCTACCACCGATGGCACA R: GCGCGTTTCGTTTGATTCATAC	60
vrrB1	F: ATAGGTGGTTTTCCGCAAGTTATTC R: GATGAGTTTGATAAGAATAGCCTGTG	61
vrrB2	F: CACAGGCTATTCTTTATCAAACATCATC R: CCCAAGGTGAAGATTGTTGTTGA	60
vrrC1	F: GAAGCAAGAAAAGTGATGTAGTGGCA R: CATTTCCTCAAGTGCTACAGGTTTC	62
vrrC2	F: CGAGAAGAAGTGGAAACCTGTAGCAC R: GTCTTTCCATTAATCGCGCTCTATC	62
CG3	F: TGTCGTTTTATCTTCTCTCTCCAATAC R: AGTCATTGTTCTGTATAAAGGGCAT	59
pXO1	F: CAATTTATTAACGATCAGATTAAGTTCA R: TCTAGAATTAGTTGCTTCATAATGGC	57
CB-1	F: GTTGAGCATGAGAGGTACCTTGTCCTTTT R: AGTTCAAGCGCCAGAAGGTTATGAGTTATC	66
CB-3	F: GCAGCAACAGAAAACCTCTCTCCAATAACA R: TCCTCCCTGAGAACTGCTATCACCTTTAAC	64
CB-5	F: GCAGGAAGAACAAGAAACTAGAAAGAGCA R: ATTATTAGCAGGGGCCTCTCCTGCATTACC	64
CB-7	F: GAATATTCGTGCCACCTAACAAAACAGAAA R: TGTCAGATCTCTAGTTGGCCCTACTTTTCCTC	63
CB-13	F: AATTGAGAAATTGCTGTACCAAAC R: CTAGTGCAATTTGACCCTAACTTGT	58
CB-15	F: GTATTTCCCCAGATACAGTAATCC R: GTGTACATGTTGATTCATGCTGTTT	59
CB-21	F: TGTAGTGCCAGATTTGTCTTCTGTA R: CAAATTTTGAGATGGGAGTTTACT	58
CB-22	F: ATCAAAAATTCTTGGCAGACTGA R: ACCGTTAATTCACGTTTAGCAGA	57
CB-23	F: CGGTCTGTCTCTATTATTCAGTGGT R: CCTGTTGCTCCTAGTGATTTCTTAC	62
CB-24	F: CTTCTACTCCGTAATTGAAATTGG R: CGTCACGTACCATTTAATGTTGTTA	59
CB-25	F: CCGAATACGTAAGAAATAAATCCAC R: TGAAAGATCTTGAAAAACAAGCATT	56
CB-28	F: CTCTGTTGTAACAAAATTTCCGTCT R: TATTAACCAGGAGTTACTTACAGC	59
CB-30	F: AGCTAATCACCTACAACACCTGGTA R: CAGAAAATATTGGACCTACCTTCC	61
CB-31	F: GCTGTATTTATCGAGCTTCAAATCT R: GGAGTACTGTTTGTGAATGTTGTTT	59

help draw sound conclusions on the exact spread of *B. anthracis*.

#### ACKNOWLEDGEMENTS

The authors are grateful to the School of Veterinary Medi-

cine, University of Zambia, Department of Pathology and Microbiology, University Teaching Hospital, Lusaka, Zambia and the Wellcome trust (Grant number WT 087546MA) for one medicine Africa-UK research capacity development partnership for infectious diseases in Southern Africa under the Southern African Center for infectious disease surveillance.

**Table 2.** Genotyping results of *Bacillus anthracis* strains isolated from soil, hippopotamus and humans with the 21 VNTR locus.

Strain	VNTR markers																					
	vrRA1	vrRB1	vrRB2	vrRC1	vrRC2	CG3	pXO1	CB-1	CB-3	CB-5	CB-7	CB-13	CB-15	CB-21	CB-22	CB-23	CB-24	CB-25	CB-28	CB-30	CB-31	
Outbreak isolates	+	+	+	+	+	–	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Sterne Vaccine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

## REFERENCES

- Baykam N, Ergonul O, Ulu A, Eren S, Celikbas A, Eroglu M, Dokuzoguz B (2009). Characteristics of cutaneous anthrax in Turkey. *J. Infect. Dev. Ctries.* 3: 599-603.
- Brodzik AK, Francoeur J (2011). A new approach to in silico SNP detection and some new SNPs in the *Bacillus anthracis* genome. *BMC Res. Notes* 4: 114.
- Gierczynski R, Jakubczak A, Jagielski M (2009). "Extended multiple-locus variable-number tandem-repeat analysis of *Bacillus anthracis* strains isolated in Poland. *Pol. J. Microbiol.* 58: 3-7.
- Habrun B, Racic I, Kompes G, Spicic S, Benic M, Mihaljevic Z, Cvetinic Z (2011). The antimicrobial susceptibility and virulence factors of *Bacillus anthracis* strains isolated in Croatia. *Vet. Med-Czech.* 56: 22–27.
- Hang'ombe BM, Mwansa JCL, Muwowo S, Mulenga P, Kapina M, Musenga E, Squarre D, Mataa L, Thomas SY, Ogawa H, Sawa H, Higashi H (2012). Human – Animal Anthrax outbreak in the Luangwa valley of Zambia in 2011. *Trop. Doct.* 42:136-139.
- Jackson PJ, Eliza AW, Abdullah SK, Kara R, Debra MA, Karen KH, Cheryl RK, Gary A, Kenneth HW, Martin EHJ, Keim P (1997). Characterization of the Variable-Number Tandem Repeats in *vrRA* from different *Bacillus anthracis* isolates. *Appl. Environ. Microbiol.* 63: 1400–1405.
- Keim P, Kalif A, Schupp J, Hill K, Travis SE, Richmond K, Adair DM, Hugh-Jones M, Jackson P (1997). Molecular evolution and diversity in *Bacillus anthracis* as detected by amplified fragment length polymorphism markers. *J. Bacteriol.* 179: 818–824.
- Keim P, Van Ert MN, Pearson T, Vogler AJ, Huynh LY, Wagner DM (2004). Anthrax molecular epidemiology and forensics: using the appropriate marker for different evolutionary scales. *Infect. Genet. Evol.* 4: 205-213.
- Koehler TM (2002). *Bacillus anthracis* genetics and virulence gene regulation. *Curr. Top. Microbiol. Immunol.* 71: 143–164.
- Le Flèche P, Hauck Y, Onteniente L, Prieur A, Denoëud F, Ramisse V, Sylvestre P, Benson G, Ramisse F, Vergnaud G (2001). A tandem repeats database for bacterial genomes: application to the genotyping of *Yersinia pestis* and *Bacillus anthracis* *BMC Microbiol.* 1: 2.
- Lindstedt BA (2005). Multiple-locus variable number tandem repeats analysis for genetic fingerprinting of pathogenic bacteria. *Electrophoresis* 26: 2567-2582.
- Okinaka RT, Henrie M, Hill KK, Lowery KS, Van Ert M, Pearson T, Schupp J, Kenefic L, Beaudry J, Hofstadler SA, Jackson PJ, Keim P (2008). Single nucleotide polymorphism typing of *Bacillus anthracis* from Sverdlovsk tissue. *Emerg. Infect. Dis.* 14: 653-656.
- Schupp JM, Klevytska AM, Zinser G, Price LB., Keim P (2000). *vrRB*, a hypervariable open reading frame in *Bacillus anthracis*. *J. Bacteriol.* 182: 3989–3997.
- Siamudaala VM, Bwalya JM, Munang'andu HM (2006). Ecology and epidemiology of anthrax in cattle and humans in Zambia. *Jpn. J. Vet. Res.* 54:15-23.
- Spencer RC (2003). *Bacillus anthracis*. *J. Clin. Pathol.* 56: 182-187.
- Tuchili LM, Pandey GS, Sinyangwe PG, Kaji T (1993). Anthrax in cattle, wildlife and humans in Zambia. *Vet. Rec.* 132: 487.
- Turnbull PC, Bell RH, Saigawa K, Munyenyembe FE, Mulenga CK, Makala LH (1991). Anthrax in wildlife in the Luangwa Valley, Zambia. *Vet. Rec.* 128: 399 – 403.
- Turnbull PCB (1998). Guidelines for the Surveillance and Control of Anthrax in Human and Animals. 3rd edition. WHO/EMC/ZDI/98.6.
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M (1995). AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res.* 23: 4407-4414.