

Full Length Research Paper

# Tomato (*Solanum lycopersicum* L.) seedling growth and development as influenced by *Trichoderma harzianum* and arbuscular mycorrhizal fungi

Bombiti Nzanza\*, Diana Marais and Puffy Soundy

Department of Plant Production and Soil Science, Faculty of Natural and Agricultural Sciences, University of Pretoria, Pretoria 0002, South Africa.

Accepted 18 February, 2011

Recent trends in soil microbiology suggest that certain soil microbes have a positive effect on seedling growth and development. A study was conducted to investigate the interactive effect of the plant-growth promoting fungi *Trichoderma harzianum* and the arbuscular mycorrhizal fungi (AMF) in growth and development of tomato (*Solanum lycopersicum*) seedlings grown under greenhouse conditions. A 3 × 3 factorial experiment was laid out in a completely randomised design with six replications. At harvest (42 DAP), when compared with the control, *T. harzianum* and/or AMF treated plants improved shoot length, root length, dry shoot mass and dry root mass. Pre-inoculation with AMF increased shoot N, P and S content of tomato seedlings, whereas pre-sowing with *T. harzianum* alone increased the shoot N. Generally, shoot Zn and Mn content were affected by both fungi, with the best result observed when AMF was applied 2 weeks after *T. harzianum*. The percentage of roots colonised by AMF was less than 15% regardless of the time when *T. harzianum* was applied. However, the percentage of roots colonised by *T. harzianum* was greater than 90% at all times. In conclusion, this study suggested that *T. harzianum* and AMF have the potential to improve tomato seedling growth and development.

**Key words:** Essential mineral nutrients, mycorrhiza, plant-growth promoting fungi, seedling quality, *Solanum lycopersicum*.

## INTRODUCTION

The need to produce quality tomato seedlings, capable of withstanding adverse abiotic and biotic stresses after transplanting and improve mineral nutrient uptake, inspired producers to consider a combined pre-sowing inoculation of seedlings with *Trichoderma harzianum* and arbuscular mycorrhizal fungi (AMF). Nursery inoculation of tomato with AMF resulted in stronger and superior quality seedlings (Giannuzzi et al., 2001), higher crop uniformity (Waterer and Coltman, 1988), better mineral

nutrient uptake (Bethlenfalvay et al., 1988; Chandanie et al., 2009; Marschner and Dell, 1994), improved tolerance to soil borne diseases (Pozo and Azcón-Aguilar, 2007), and both reduced stress and increased yields (Chandanie et al., 2009; Lovato et al., 1996). Similarly, *T. harzianum* enhanced plant growth and development (Harman and Taylor, 1990; Liu et al., 2008; Samuels, 2006), and provided protection against soil-borne pathogens that cause damping-off in tomato seedlings (Harman and Taylor, 1990).

The symbiosis between *T. harzianum* and AMF is widely reported in literature (Meyer and Roberts, 2002; Raupach and Kloepper, 1998). *Trichoderma* species have both antagonistic (Camporota, 1985; McAllister et al., 1994;

\*Corresponding author. E-mail: [s24493024@tuks.co.za](mailto:s24493024@tuks.co.za). Tel: +27153952135. Fax: +27153952135.

Wyss et al., 1992) and stimulating effects on AMF (Calvet et al., 1992; McAllister et al., 1994) and *vice versa*. Antagonistic modes of action of Trichoderma to AMF include competition, myco-parasitism and production of antifungal metabolites (Lorito et al., 1993; Stefanova et al., 1999). Also, the species have a high reproductive capacity estimated at 12 h for spore germination (Liu et al., 2008; Woo et al., 2005). In spite of the increasing interest in the interaction between *T. harzianum* and AMF, information about these interactions in tomato seedling production is scanty (Fracchia et al., 1998; McAllister et al., 1994). The objective of this study was to investigate the interactive effects of nursery inoculation of *T. harzianum* and AMF on their impact on growth and development of tomato seedlings when applied at different times.

## MATERIALS AND METHODS

### Location

The experiment was conducted under greenhouse conditions at the Hatfield Experimental Farm, University of Pretoria, South Africa, during the 2008 growing season and repeated in 2009. The site is located at 23° 45' S latitude, 28° 16' E longitude, at 1372 m above sea level.

### Microbial inoculants

Commercial mycorrhizal inoculum Biocult® containing spores of *Glomus mossae*, was obtained from Biocult Ltd. (Somerset West, South Africa). Commercial Trichoderma inoculum T-GRO containing spores of *T. harzianum* isolate DB 103 ( $1 \times 10^9$  colony forming units  $g^{-1}$ , as a wettable powder) was obtained from Dagugat Biolab (Johannesburg, South Africa). The microbial inoculants were thoroughly mixed with peat moss and vermiculite before applying them into the pasteurised sand: coir (seedling trays) or peat moss (PVC pipe) mixtures used for seedling production. The microbial inoculants were introduced either before sowing the seed or before transplanting the seedlings (two weeks later).

### Experimental design and treatments

The nine treatment combinations, namely  $T_0M_0$  (untreated/control),  $T_0M_1$  (treated with AMF only, before sowing),  $T_0M_2$  (treated with AMF only, 2 weeks after sowing),  $T_1M_0$  (treated with *T. harzianum* only, before sowing),  $T_1M_1$  (treated with both fungi before sowing),  $T_1M_2$  (treated with *T. harzianum* before and AMF two weeks after sowing),  $T_2M_0$  (treated with *T. harzianum* only, 2 weeks after sowing),  $T_2M_1$  (treated with *T. harzianum* at 2 weeks after sowing and AMF before sowing) and  $T_2M_2$  (treated with both fungi 2 weeks after sowing), were arranged in a completely randomised design with six replications.

Seeds of tomato cv. 'Nemo-Netta' were sown into cell plug trays filled with a pasteurised sand and coir mixture at ratio 50:50 (v/v). Trays were transferred to the germination room for 3 days and then moved to the greenhouse. Two weeks after sowing, seedlings were transplanted into a 30 cm long PVC pipe (diameter: 3.5 cm) filled

with peat moss and supported by a cylinder base. Plants were fertilised with half strength modified Hoagland's solution (Spomer et al., 1997) and watered daily.

### Data collection

At harvest, 42 days after initiating the treatment, plant height, root length, stem diameter and leaf area were recorded. Roots were separated from shoots and sampled for defeminisation of colonisation with the two fungal species. Roots of randomly selected tomato seedlings were washed free of medium, stained with trypan blue in lactophenol (Phillips and Hayman, 1970) and quantified for percentage of AMF colonisation using the line-intersect method (Brundrett et al., 1996). Root colonisation by *T. harzianum* was also determined (Datnoff et al., 1995).

Shoots and the remaining roots were oven-dried at 50°C for 70 h to determine dry shoot and dry root mass. Dried shoots and roots were each ground in a Wiley mill to pass through 1 mm sieve. 1 g sample was digested in sulphuric acid at 410°C and N determined by an auto analyser. Other essential nutrient elements were digested with a 2:1 nitric/perchloric acid mixture at 230°C and nutrient elements determined by the inductive coupled plasma (ICP).

### Data analysis

Data were subjected to analysis of variance using SAS (SAS Institute Inc., Cary, NC, USA. (2002 to 2003). The degrees of freedom and their associated sum of squares were partitioned to provide the total treatment variation for different sources of variation (Little, 1981). Mean separation was achieved using Fisher's least significant difference test. Unless stated otherwise, treatments discussed were different at 5% level of probability.

## RESULTS

### Root colonisation by fungi

The *T. harzianum* × AMF on root colonisation was not significantly different during both growing seasons (Table 1). Roots of treated *T. harzianum* seedlings had more than 90% root colonisation; AMF-treated seedlings had less than 15% colonisation, whereas untreated roots had no colonisation. Using the partitioning of the degrees of freedom and their associated sum of squares *T. harzianum* contributed 99% to total treatment variation (TTV) in percentage Trichoderma colonisation, whereas AMF accounted for over 96% of the TTV in colonisation.

### Growth parameters

This analysis revealed a significant interactive effect of *T. harzianum* and AMF for plant height and root length, which only explained half of the total variability (Table 1). *T. harzianum* contributed ca. 40% of the TTV in the mean plant height. The treatment also explained 21 and 29% of

**Table 1.** Partitioning of the treatment sum of squares derived from the analysis of variance for the plant growth variables and root colonisation of 6-weeks old tomato seedlings as influenced by *Trichoderma harzianum* and AMF inoculation.

Source	Df	Mycorrhiza		Trichoderma		Plant height		Root length		Dry shoot mass		Dry root mass	
		SS	%	SS	%	SS	%	SS	%	SS	%	SS	%
<b>2008 growing season</b>													
<i>T. harzianum</i> (T)	2	19.7	2.1ns	107215	99.7*	455.39	41.2***	185.27	20.5***	92.01	45.0***	6.95	56.6***
AMF (M)	2	902.48	96.1*	15	0.0ns	87.66	7.9***	260.34	28.8***	32.63	15.9*	0.87	7.1ns
T×M	4	16.74	1.8ns	296	0.3ns	561	50.8***	459.21	50.8***	79.99	39.1**	4.45	36.3*
Total	53	938.92		107526		1104.05		904.82		204.62		12.27	
<b>2009 growing season</b>													
<i>T. harzianum</i> (T)	2	4.59	0.2ns	98415	99.8*	145.67	40.1**	135.38	29.3ns	37.39	81.1***	1.14	78.7*
AMF (M)	2	2013.37	99.3*	104	0.1ns	50.65	13.9ns	70.27	15.2ns	2.11	4.6ns	0.04	2.6ns
T×M	4	10.52	0.5ns	74	0.1ns	167.34	46.0**	256.99	55.5*	6.59	14.3ns	0.27	18.8ns
Total	53	2028.48		98593		363.65		462.64		46.09		1.44	

ns, \*, \*\*, \*\*\* are levels of significance at  $P \geq 0.10$ ,  $P \leq 0.05$ ,  $P \leq 0.01$ ,  $P \leq 0.00$ , respectively.

the TTV in mean root length in 2008 and 2009 growing seasons, respectively. In 2008, AMF contributed 29% of the TTV in mean root length but only 15% during the second growing season. During the first season, inoculating both fungi at sowing ( $T_1M_1$ ) increased plant height and root length by 40 and 30%, respectively, as compared to the control plants (Table 2). The highest plant height was obtained with late *T. harzianum* inoculation ( $T_2M_0$ ). In 2009, the highest plant height and root length were recorded with  $T_1M_1$  and  $T_2M_0$ , respectively, whereas the lowest counts were obtained in the untreated plants ( $T_0M_0$ ). In both seasons, all the microbial inoculated seedlings, except for late microbial inoculations ( $T_2M_2$ ), when compared with the control increased plant height and root length.

### Biomass production

There was a significant *T. harzianum* × AMF effect on dry shoot and root mass during the first growing season, which accounted for ca. 40% of the TTV of dry shoot mass. The major source of variability was due to *T. harzianum*, which contributed nearly 50% of the TTV of dry shoot mass. Interestingly, in 2009, *T. harzianum* accounted for ca. 80% of the TTV with small contributions from AMF and *T. harzianum* × AMF interactions. During the first season, compared to the control plants, the combined inoculation of *T. harzianum* and AMF before sowing resulted in 35% higher dry shoot mass, whereas inoculating both fungi simultaneously 2 weeks after sowing, resulted only in 13% increase. The highest

increase (52%) in dry shoot mass was obtained with  $T_1M_0$ . All microbial inoculants increased dry shoot mass (Table 2). Dry root mass was increased (up to 37%) when *T. harzianum* was inoculated before planting and AMF 2 weeks later ( $T_1M_2$ ).

However, a negative interaction between *T. harzianum* and AMF was observed when both fungi were applied 2 weeks after sowing ( $T_2M_2$ ), resulting in the lowest dry root mass. During the second season, irrespective of the AMF treatment, inoculating *T. harzianum* before sowing increased the dry mass of the shoot and root by 19 and 11%, respectively, whereas dry shoot and root mass in plants inoculated with *T. harzianum* 2 weeks later, did not differ from those of the control.

**Table 2.** Plant growth variables of 6-week old tomato seedlings as influenced by *Trichoderma harzianum* and AMF inoculation.

Treatment	Plant height (cm)			Root length (cm)			Dry shoot mass (g plant <sup>-1</sup> )			Dry root mass (g plant <sup>-1</sup> )		
	M <sub>0</sub>	M <sub>1</sub>	M <sub>2</sub>	M <sub>0</sub>	M <sub>1</sub>	M <sub>2</sub>	M <sub>0</sub>	M <sub>1</sub>	M <sub>2</sub>	M <sub>0</sub>	M <sub>1</sub>	M <sub>2</sub>
<b>2008 growing season</b>												
T <sub>0</sub>	16.73f	25.12c	21.40e	22.38e	33.74a	29.28bc	6.00d	8.80bc	6.93cd	1.89b	2.46ab	1.94b
T <sub>1</sub>	27.34b	28.11ab	28.56a	26.63d	34.23a	32.66a	12.50a	9.17bc	9.36bc	2.91a	2.83a	2.98a
T <sub>2</sub>	29.16a	23.08d	17.15f	29.86b	26.92cd	23.21e	10.31ab	6.91cd	6.89cd	2.89a	1.91ab	1.84b
<b>2009 growing season</b>												
T <sub>0</sub>	20.25d	26.52ab	25.15bc	21.82c	28.88ab	26.33bc	8.24d	9.33bcd	8.71cd	2.47c	2.67abc	2.50bc
T <sub>1</sub>	27.07ab	29.30a	27.31ab	27.80ab	29.68ab	30.10ab	10.58ab	10.70ab	10.80a	2.79abc	2.90ab	2.92a
T <sub>2</sub>	27.47ab	25.30bc	22.66cd	31.75a	30.00ab	24.82bc	9.75abc	9.46abcd	8.54cd	2.69abc	2.54abc	2.51bc

T<sub>1</sub>, T<sub>2</sub> and T<sub>0</sub>: *T. harzianum* inoculated before sowing, 2 weeks after sowing or uninoculated. M<sub>1</sub>, M<sub>2</sub> and M<sub>0</sub>: AMF inoculated before sowing, 2 weeks after sowing or uninoculated. Column means followed by the same letter were not significantly different at 5% level according to Fisher's least significant different test.

### Shoot chemical analysis

Neither *T. harzianum* nor AMF affected essential nutrient elements such as K, Ca, Mg, Mo and Na. There was a significant *T. harzianum* × AMF interaction term for the shoot Mn and Zn content, whereas P and S were only affected by AMF. Mean shoot N content of seedlings was affected by *T. harzianum* and AMF, but not their interaction (Table 3).

Inoculating *T. harzianum* before sowing (T<sub>1</sub>) increased the N shoot content by 6%, whereas later inoculation was similar to the uninoculated plants (T<sub>0</sub>). On the other hand, when compared with the control (M<sub>0</sub>), inoculating AMF before (M<sub>1</sub>) or 2 weeks after sowing (M<sub>2</sub>) increased the shoot N content by 9 and 10%, respectively (Table 4).

Inoculating AMF before (M<sub>1</sub>) or after sowing (M<sub>2</sub>) increased the shoot P content of tomato seedlings by ca. 18 and 16%, respectively. Shoot

S increased by 15% when AMF was inoculated before sowing (M<sub>1</sub>), whereas later inoculation (M<sub>2</sub>) had no effect on this nutrient element (Table 4). Inoculating *T. harzianum* and AMF before (T<sub>1</sub>M<sub>1</sub>) or after (T<sub>2</sub>M<sub>2</sub>) sowing increased Mn content by 18 and 9%, respectively. However, the highest Mn shoot content increase (33%) was obtained with a combination of early *T. harzianum* and late AMF application (T<sub>1</sub>M<sub>2</sub>) (Table 5). Similarly, for Zn shoot content, the highest increase (34%) was recorded with T<sub>1</sub>M<sub>2</sub>, while T<sub>1</sub>M<sub>1</sub> and T<sub>2</sub>M<sub>2</sub> yielded about 13 and 10% Zn increase, respectively.

### DISCUSSION

Nursery inoculation of tomato with *T. harzianum* and AMF improved most of the growth variables of tomato seedlings, increased nutrient element uptake and permit microbial root colonisation.

Uninoculated plants showed no *Trichoderma* or AMF or colonisation, indicating that these fungi were not indigenous to the specific growth media. The low mycorrhizal colonisation (< 15%) observed was in agreement with Chandanie et al. (2009), who argued that the 13% level of colonisation with AMF observed before transplanting should be considered adequate for successful establishment of mycorrhizal seedlings. According to Bierman and Linderman (1983), less than 13% root colonisation should not be a concern, as these fungi would spread rapidly to new roots after transplanting. On the other hand, the higher *Trichoderma* root colonisation was due to its high reproductive capacity (Woo et al., 2005).

Observations in this study suggested that, low mycorrhizal and high *Trichoderma* root colonisations were due to their inherent individual abilities to colonise tomato roots rather than their

**Table 3.** Results of ANOVA (P values) executed for the shoot mineral nutrient content for the 2008 growing season on tomato seedlings at 42 days after planting.

Response variables	N	P	K	Ca	Mg	S	Mn	Zn	Cu	Mo	Na
T (df = 2)	*	ns									
M (df = 2)	**	*	ns	ns	ns	*	*	*	ns	ns	ns
T×M (df =4)	ns	ns	ns	ns	ns	ns	*	*	ns	ns	ns

ns, \*, \*\*, \*\*\* are levels of significance at  $P \geq 0.10$ ,  $P \leq 0.05$ ,  $P \leq 0.01$ ,  $P \leq 0.00$ , respectively. T = *T. harzianum*; M= AMF.

**Table 4.** Macronutrients shoot content of 6-week old tomato seedlings as influenced by *T. harzianum* and AMF applied before sowing and at two weeks after sowing.

Response variable	N (%)	P (%)	K (%)	Ca (%)	Mg (%)	S (%)
<b>T (<i>T. harzianum</i>)</b>						
T <sub>0</sub>	4.42b	0.62	2.97	4.19	1.06	1.63
T <sub>1</sub>	4.72a	0.63	2.75	4.17	1.03	1.56
T <sub>2</sub>	4.45b	0.60	2.72	4.48	1.13	1.77
<b>M (AMF)</b>						
M <sub>0</sub>	4.23b	0.54b	2.80	4.00	1.05	1.57b
M <sub>1</sub>	4.65a	0.66a	2.86	4.47	1.13	1.83a
M <sub>2</sub>	4.71a	0.64a	2.77	4.37	1.05	1.56b

T<sub>1</sub>, T<sub>2</sub> and T<sub>0</sub>: *T. harzianum* inoculated before sowing, 2 weeks after sowing or uninoculated. M<sub>1</sub>, M<sub>2</sub> and M<sub>0</sub>: AMF inoculated before sowing, 2 weeks after sowing or uninoculated. Column means followed by the same letter were not significantly different at 5% level according to Fisher's least significant different test.

**Table 5.** Micronutrient shoot contents of 6-week old tomato seedlings as influenced by AMF pre-inoculation.

Treatment	Mn (ppm)			Zn (ppm)			Mo (ppm)			Na (ppm)		
	M <sub>0</sub>	M <sub>1</sub>	M <sub>2</sub>	M <sub>0</sub>	M <sub>1</sub>	M <sub>2</sub>	M <sub>0</sub>	M <sub>1</sub>	M <sub>2</sub>	M <sub>0</sub>	M <sub>1</sub>	M <sub>2</sub>
T <sub>0</sub>	19.67d	27.67ab	26.00abc	24.67d	35.00	32.67abc	14.86ab	16.18a	14.80ab	1967.7a	1851.0ab	1466.0b
T <sub>1</sub>	22.67cd	24.00bcd	29.33a	28.33cd	30.33bcd	37.33a	14.26ab	14.39ab	13.72	1922.7a	1935.3a	1717.3ab
T <sub>2</sub>	23.00cd	22.33cd	21.67cd	29.33bcd	27.67cd	27.33cd	14.20ab	12.64b	14.97ab	1750.7ab	1807.7ab	1621.7ab

T<sub>1</sub>, T<sub>2</sub> and T<sub>0</sub>: *T. harzianum* inoculated before sowing, 2 weeks after sowing or uninoculated. M<sub>1</sub>, M<sub>2</sub> and M<sub>0</sub>: AMF inoculated before sowing, 2 weeks after sowing or uninoculated. Column means followed by the same letter were not significantly different at 5% level according to Fisher's least significant different test.

competitive interactions. However, the observation was not in agreement with McGovern et al. (1992) who reported antagonistic effect of *Trichoderma* on AMF in tomato. Chandanie et al. (2009) observed a decreased *T. harzianum* growth due to AMF inoculation in cucumber (*Cucumis sativus*).

However, Green et al. (1999) observed a mutually inhibitory interaction between *T. harzianum* and the external mycelia of an AMF *Glomus intraradices*. Apparently, the interaction between *Trichoderma* and AMF is species and host-plant specific (Fracchia et al., 1998; Green et al., 1999; Rousseau et al., 1996).

*Trichoderma harzianum* and AMF, either inoculated

alone or in combination, increased the root length and plant height of tomato. Generally, improved plant growth had been observed under *Trichoderma* (Duffy et al., 1997; Ozbay and Newman, 2004) and AMF inoculations (Tahat et al., 2008). Improved plant growth observed in these experiments might be due to increased solubility of insoluble plant nutrients by *Trichoderma* species (Kaya et al., 2009) or enhanced immobile nutrient elements uptake by AMF (Bethlenfalvay et al., 1988; Chandanie et al., 2009; Marschner and Dell, 1994). Observations in this study suggested that, there were the desired beneficial effect of nursery inoculation with *T. harzianum* and/or AMF on dry matter production of tomato seedlings, which

was in agreement with Ozbay and Newman (2004), who observed an increased dry shoot mass due to *Trichoderma* inoculation and Tahat et al. (2008) who observed similar trends with AMF. Chandanie et al. (2009) demonstrated that, the combined inoculation of AMF with *Trichoderma* synergistically increased dry shoot mass when compared with inoculation of *Trichoderma* and AMF alone. McAllister et al. (1994) reported a decrease in dry shoot mass when *Trichoderma* was inoculated alone before sowing or at the same time with AMF. In this study, both fungi either applied alone or in combination, improved plant growth, except when simultaneously applied 2 weeks after sowing. The negative interaction when combined inoculation is applied at 2 weeks could be due to competition for nutrients or space.

Nursery microbial inoculation had no effect on K, Ca and Mg shoot content, which was in agreement with Karagiannidis et al. (2002), who did not find any positive effect of mycorrhiza on shoot K and Ca content. Increased K and Mg content have been reported in wheat inoculated with AMF (Tarafdar and Marschner, 1995), whereas *Trichoderma* species did not increase the shoot Ca, K and Mg content in tomato seedlings grown in hydroponics (Yedidia et al., 2000). Nevertheless, there were beneficial effects of AMF inoculation on shoot N, P and S in tomato seedlings. Increased N uptake due to AMF inoculation had been reported previously (Karagiannidis et al., 2002; Thomson et al., 1996). Similarly, increased shoot P contents following AMF inoculation were in agreement with other observations (Al-Karaki, 2006; Nurlaeny et al., 1996; Yedidia et al., 2000), whereas others did not observe any positive effect (Inbar et al., 1994). Late inoculation had no effect on shoot S, suggesting that early application was advisable for increased S uptake. Increased S content of plants with mycorrhiza had been reported previously (Rhodes and Gerdemann, 1978).

Shoot Zn and Mn increased in nursery inoculation, probably due to an increased absorptive capability for these nutrient elements when tomato roots are colonised by *Trichoderma* and AMF, as suggested for pepper plants (Kaya et al., 2009). However, this is in disagreement with a reduced concentration of Mn and Zn on leaves of AMF infected plants (Weissenhorn et al., 1995). Other micronutrients such as Cu, Mo and Na were unaffected by the nursery microbial inoculation, possibly due to their low concentration in the growing medium.

## Conclusion

Nursery inoculation of tomato with *T. harzianum* and/or AMF improved growth and development of tomato seedlings. *T. harzianum* and AMF synergistically

improved most of the growth variables in tomato seedlings. A negative *T. harzianum* × AMF interaction was only observed 2 weeks after sowing, probably due to competition for nutrient elements and/or infection sites. In contrast to *T. harzianum*, which had little effect on essential nutrient elements, AMF inoculation affected the nutrient uptake of key elements such as N, P, S, Zn and Mn. Although, the myco-parasitic effect of *Trichoderma* species is well known, results of this study demonstrated that, this plant-growth promoting fungi can successfully be inoculated with AMF for improved seedling health and development in tomato production.

## ACKNOWLEDGMENT

The authors acknowledge the support from Bertie van Zyl Pty (Ltd) ZZ2 for funding this work.

## REFERENCES

- Al-Karaki GN (2006). Nursery inoculation of tomato with arbuscular mycorrhizal fungi under subsequent performance under irrigation with saline water. *Sci. Hortic.*, 109: 1-7.
- Bethlenfalvay GJ, Brown MS, Ames RN, Thomas RS (1988). Effects of drought on host and endophyte development in mycorrhizal soybeans in relation to water use and phosphate uptake. *Physiol. Plant*, 72: 565-571.
- Bierman BJ, Linderman RG (1983). Increased geranium growth using pretransplant inoculation with a mycorrhizal fungus. *J. Am. Soc. Hortic. Sci.*, 108: 972-976.
- Brundrett M, Bougher N, Dell B, Grove T, Malajczuk N (1996). Working with Mycorrhizas in Forestry and Agriculture. *ACIAR*, 32: 374.
- Calvet C, Pera J, Barea J (1992). *In vitro* interactions between the vesicular-arbuscular mycorrhizal fungus *Glomus mosseae* and some saprophytic fungi isolated from organic substrates. *Soil Biol. Biochem.*, 24 : 775-780.
- Camporota P (1985). Antagonistic action of *Trichoderma spp vis-à-vis* de *Rhizoctonia solani* Kühn. *Agronomie*, 5: 613-620.
- Chandanie WA, Kubota M, Hyakumachi M (2009). Interaction between the arbuscular mycorrhizal fungus *Glomus mosseae* and plant growth-promoting fungi and their significance for enhancing plant growth and suppressing damping-off of cucumber (*Cucumis sativus* L.). *Appl. Soil Ecol.*, 41: 336-341.
- Datnoff LE, Nemecek S, Pernezny K (1995). Biological control of Fusarium crown and root rot of tomato in Florida using *Trichoderma harzianum* and *Glomus intraradices*. *Biol. Control.*, 5: 427-431.
- Duffy BK, Ownley BH, Weller DM (1997). Soil chemical and physical properties associated with suppression of take-all of wheat by *Trichoderma koningii*. *Phytopathol.*, 87: 1118-1124.
- Fracchia S, Mujica MT, Garcia-Romera I, Garci-Garrido JM, Martin J, Ocampo JA, Godeas A (1998). Interactions between *Glomus mosseae* and arbuscular mycorrhizal sporocarp-associated saprophytic fungi. *Plant Soil*, 200: 131-137.
- Giannuzzi S, Schuepp H, Barea JM, Haselwandter K (2001). Mycorrhizal technology in Agriculture: From genes to bioproducts. *Birkhauser*, Basel, Switzerland.
- Green H, Larsen J, Olsson PA, Jensen DF, Jakobsen I (1999). Suppression of the biocontrol agent *Trichoderma harzianum* by mycelium of the arbuscular mycorrhizal fungus *Glomus intraradices* in root-free soil. *Appl. Environ. Microbiol.*, 65: 1428-134.
- Harman GE, Taylor AG (1990). Development of an effective biological seed treatment system. In "Biological control of soil borne pathogens"

- (Hornby D and Cook, R.J, Eds). CAB International, Wallingford, UK., pp. 415-426.
- Inbar J, Abramsky M, Cohen D, Chet I (1994). Plant growth enhancement and disease control by *Trichoderma harzianum* in vegetable seedlings grown under commercial conditions. *Eur. J. Plant Pathol.*, 100: 337-346.
- Karagiannidis N, Bletsos F, Stavropoulos N (2002). Effect of *Verticillium* wilt (*Verticillium dahliae* Kleb.) and mycorrhiza colonization, growth and nutrient uptake in tomato and eggplant seedlings. *Sci. Hortic.*, 94: 145-156.
- Kaya C, Ashraf M, Sonmez O, Aydemir S, Tuna AL, Cullu MA (2009). The influence of arbuscular mycorrhizal colonization on key growth parameters and fruit yield of pepper plants grown at high salinity. *Sci. Hortic.*, 121: 1-6.
- Little TM (1981). Interpretation and presentation of results. *HortScience*, 16: 19-22.
- Liu B, Glenn D, Buckley K (2008). *Trichoderma* communities in soils from organic, sustainable, and conventional farms, and their relation with southern blight of tomato. *Soil Biol. Biochem.*, 40: 1124-1136.
- Lorito M, Harman GE, Hayes CK, Broadway RM, Tronsmo A, Woo SL, Di Petro A (1993). Chitinolytic enzymes produced by *Trichoderma harzianum*: antifungal activity of purified endochinase and chetobiosidase. *Phytopathol.*, 83: 302-307.
- Lovato PE, Gininazzi-Pearoon V, Trouvelot A, Gininazzi S (1996). The state of art of mycorrhizas and micropropagation. *Adv. Hortic. Sci.*, 10: 46-52.
- Marschner H, Dell B (1994). Nutrient uptake in mycorrhizal symbiosis. *Plant Soil*, 159: 89-102.
- Mcallister CB, Garcia-Romera I, Godeas A, Ocampo JA (1994). Interaction between *Trichoderma koningii*, *Fusarium solani* and *Glomus mosseae*: Effect on plant growth, arbuscular mycorrhizas and saprophytic populations. *Soil Biol. Biochem.*, 26: 1363-1367.
- McGovern RJ, Datnoff LE, Tripp L (1992). Effect of mixed infection and irrigation method on colonization of tomato roots by *Trichoderma harzianum* and *Glomus intraradices*. *Proc. Fla. State Hortic. Soc.*, 105: 361-363.
- Meyer SLF, Roberts DP (2002). Combinations of biocontrol agents for management of plant-parasitic nematodes and soilborne plant-pathogenic fungi. *J. Nematol.*, 34: 1-8.
- Nurlaeny N, Marschner H, George E (1996). Effects of liming and mycorrhizal colonization on soil phosphate depletion and phosphate uptake by maize (*Zea mays* L.) and soybean (*Glycine max* L.) grown in two tropical acid soils. *Plant Soil*, 181: 275-285.
- Ozbay N, Newman SE (2004). Effect of *Trichoderma harzianum* strains to colonize tomato roots and improve transplant growth. *Pak. J. Biol. Sci.*, 7: 253-257.
- Phillips JM, Hayman PS (1970). Improved procedure for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. Br. Myco. Soc.*, 55: 158-161.
- Pozo MJ, Azcón-Aguilar C (2007). Unraveling mycorrhiza-induced resistance. *Plant Biol.*, 10: 393-398.
- Raupach GS, Kloepper JW (1998). Mixtures of plant growth-promoting rhizobacteria enhance biological control of multiple cucumber pathogens. *Phytopathol.*, 88: 1158-1164.
- Rhodes LH, Gerdemann JW (1978). Hyphal translocation and uptake of sulfur by vesicular-arbuscular mycorrhizae of onion. *Soil Biol. Biochem.*, 10: 355-360.
- Rousseau A, Benhamou N, Chet I, Piche Y (1996). Mycoparasitism of the extrametrical phase of *Glomus intraradices* by *Trichoderma harzianum*. *Phytopathol.*, 86: 434-443.
- Samuels GJ (2006). *Trichoderma*: systematic, the sexual state, and ecology. *Phytopathology*, 96: 195-206.
- SAS Institute (2003). *Statistical Analysis Systems Computer Package*, Cary, New York, USA.
- Spomer LA, Berry WL, Tibbitts TW (1997). *Plant culture in solid media. Plant growth chamber handbook*. R.W Langham and T.W Tibbitts (Eds.). Iowa Agriculture and Home Economics Experiment Station Special Report.
- Stefanova M, Leiva L, Larrinaga L, Courrone MF (1999). Metabolic activity of *Trichoderma* spp. isolates for control of soilborne phytopathogenic fungi. *Rev. Fac. Agro (Luz)*, 16: 509-516.
- Tahat MM, Kamaruzaman S, Radziah O, Kadir J, Masdek HN (2008). Response of (*Lycopersicon esculentum* Mill.) to different Arbuscular Mycorrhizal Fungi Species. *Asian J. Plant Sci.*, 7: 479-484.
- Tarafdar JC, Marschner H (1995). Dual inoculation with *Aspergillus fumigatus* and *Glomus mosseae* enhances biomass production and nutrient uptake in wheat (*Triticum aestivum* L.) supplied with organic phosphorus as Na-phytate. *Plant Soil*, 173: 97-102.
- Thomson TE, Manian S, Udaiyan K (1996). Interaction of multiple VAM fungal species on root colonization, plant growth and nutrient status of tomato seedlings (*Lycopersicon esculentum* Mill.). *Agr. Ecosyst. Environ.*, 59: 63-68.
- Waterer DR, Coltman RR (1988). Phosphorus concentration and application interval influence growth and mycorrhizal infection of tomato and onion transplants. *J. Am. Soc. Hortic. Sci.*, 113: 704-798.
- Weissenhorn I, Mench M, Leyval C (1995). Bioavailability of heavy metals and arbuscular mycorrhizae in a sewage-sludge-amended sandy soil. *Soil Biol. Biochem.*, 27: 287-296.
- Woo SL, Scala F, Ruocco M, Lorito M (2005). The molecular biology of the interactions between *Trichoderma* spp., phytopathogenic fungi, and plants. *Phytopathology*, 40: 309-348.
- Wyss P, Boller TH, Wiemken A (1992). Testing the effect of biological control agents on the formation of vesicular-arbuscular mycorrhiza. *Plant Soil*, 147: 159-162.
- Yedidia I, Benhamou N, Kapulnik Y, Chet I (2000). Induction and accumulation of PR proteins activity during early stages of root colonization by the mycoparasite. *Trichoderma harzianum* strain T203. *Plant Physiol. Biochem.*, 38: 863-873.