

Full Length Research Paper

Quantitative structure activity relationship studies for new antimicrobial N²- substituted phenazines

Khairi M.S. Fehelbom^{1*} and Moawia Al-Tabakha²

¹Al-Ain University of Science and Technology, College of Pharmacy, Al-Ain 64141, United Arab Emirates.

²Ajman University of Science and Technology, Faculty of Pharmacy and Health Science, Fujairah 2202, United Arab Emirates.

Accepted January 23, 2009

The increased availability of iminophenazines with polar substitute in the N²- side chain (notably B826 with a (3-diethylaminopropyl) imino substituent) in the spleen of mice following oral administration prompted an investigation of a possible correlation between the partition coefficient (log P) values and electronic properties expressed by σ^* with the concentration of the iminophenazines in the spleen of mice. The correlation studies between log P values and the concentration of the iminophenazines in the spleen of mice showed a poor correlation coefficient ($r = 0.515$), while a much better correlation was obtained using the electronic parameter σ^* , where the correlation coefficient was 0.911, indicating the significance of the electronic factor in relation to the transport of the compounds to the spleen of mice, despite the high lipophilicity of the iminophenazines.

Key words: QSAR, iminophenazines, partition coefficient, electronic parameter.

INTRODUCTION

Iminophenazines (Figure 1) exhibit antibacterial activity against species of *Mycobacterium* including *M. leprae*, the causative organism in leprosy. Clofazimine (British Pharmacopoeia, 1988) has been used for many years, either alone or in combination with a sulphone, dapsone, in the treatment of leprosy.

The development of clofazimine-resistant strains of *M. leprae* has led to the synthesis and testing of new analogues of clofazimine. (O'Sullivan et al., 1988). The N²-substituted phenazines used in this study are shown in Table 1. These compounds are incompletely absorbed from the gastro-intestinal tract and accumulate in relatively large amounts in the spleen, liver, lungs and fat from which they are slowly released. The partitioning and distribution behaviors of solutes between the organic and aqueous phases have been widely studied in quantitative structure activity relationships (Lien et al., 1986; Dearden and Bresnen, 2005). Various solvents pairs have been studied (Beezer et al., 1980; Beezer et al., 1983; Rogers and Davis, 1980; Anderson et al., 1983; Kojima and

Davis, 1984; Betageri and Rogers, 1987). The n-octanol/water system is frequently employed as a model system (Hansch and Leo, 1979; Kellogg and Abraham, 2000).

Quantitative structure relationships for many drugs using log P and other steric and electronic parameters have been extensively studied (Johnson et al., 1998; Hansch and Leo, 1995; Warhurst et al., 1998; Thakur et al., 2004).

Quantitative structure/protein binding relationships indicate that lipophilicity is the principle physicochemical determinant correlating protein binding to chemical structure (Austel and Kutter, 1983). In addition to lipophilicity the polar character of substituents, as expressed by Hammett σ^* values was also found to correlate with the observed protein binding (Martin, (1979).

In a previous study the thermodynamics of the distribution and interfacial transfer of antimicrobial iminophenazines in two phases system of n-octanol/ aqueous buffer has been reported (Fahelbom et al., 1989). In the present work correlations study between the Partition coefficient and polar electronic constant σ^* with the concentration of the iminophenazines in the spleen of mice have been investigated in order to get a suitable quantitative structure relationship for these highly lipophilic

*Corresponding author. E-mail: Khairi_f@hotmail.com.
Fax: +971-3-7611198 Tel.: +971-3-7611185.




Figure 1. Chemical structure of Iminophenazines.

Table 1. The antimicrobial activity and accumulation of the iminophenazines in the spleen and fatty tissue.

Compound	R	Growth inhibition of <i>M. smegmatis</i>		Concentration ($\mu\text{g/ml}$)
		607($\mu\text{g/ml}$)		
		Sens. to clofazimine	Res. to clofazimine	Spleen
B663	CH Me ₂	1.3	30.0	16.5
B826	(CH ₂) ₃ NEt ₂	0.2	1.0	257.0
B3779	CH(Me)(CH ₂) ₃ NEt ₂	0.2	2.0	146.0
B3770	(CH ₂) ₃ N(CH ₂) ₄ CH ₂	0.2	2.0	250.0
B3785	(CH ₂) ₃ N(CH ₂) ₃ CH ₂	0.3	0.8	103.0
B749	CH ₂ CH ₂ N(C ₂ H ₅) ₂	0.6	12.0	262.0
B3955	CH ₂ (CH ₂) ₄ NHCH ₂ CH ₂	0.4	6.0	110.0

B663* = Clofazimine.

compounds.

MATERIALS AND METHODS

Chemicals

The seven iminophenazines used in this study were synthesized in the laboratories of the Medical Research Council of Ireland, Trinity College Dublin. The solvents n-octanol, methanol, glacial acetic acid and dichloromethane were all BDH- Analar grade. The Sorenson phosphate buffer was prepared using KH₂PO₄ and Na₂HPO₄ (both from BDH- Analar grade). Tris Buffer (BDH- Analar grade) was used in pKa determinations.

Instrumentation and procedures

Determination of ionization constants: The pKa values for seven iminophenazines were determined using the spectrophotometric method described by Albert and Serjeant, 1984). The absorbance of the iminophenazines solutions were recorded at 482 nm, using an SP8-100 spectrophotometer (Pye-Unicam). A pH-meter (Radiometer-Copenhagen) was used for the determination of the pH values at 20°C.

Determination of partition coefficient: The partition coefficients of the iminophenazines was studied using a phosphate buffer pH 5.15 and n-octanol system in a thermostatically controlled shaker water bath; a shaking time of 6 h was sufficient to produce equilibration in each case, and the absorbance of the organic layer was determined spectrophotometrically as described above. The following equation was used for the partition coefficient calculations:

$$P = P' (1-\alpha)^{-1}$$

where α is the ionization degree, and is given by:

$$\alpha = (1 + 10^{\text{pH} - \text{pKa}})$$

Mouse infection and treatment: This part of the work was conducted in the National Medical Research Center (Tripoli- Libya). 33 female Swiss mice (8 -12 weeks old) weighing 18 - 20 g, were used. Before infection with *M. leprae*, the mice were housed for two weeks. Each mouse was infected in both hind footpads with a 6×10^3 CFU in a volume of 0.03 ml (Shepard, 1967). One day after infection, five mice were killed to establish baseline values of AFB and CFU counts (Degas et al., 2000), the remaining 28 were divided into two groups; a control group of 7 mice, and the other groups of 21 mice were fed orally with the phenazines derivatives.

The oral administered phenazines were prepared by dissolving 50 mg of each compound in 48 ml of mineral oil and 2 ml of water, producing a drug concentration of 1 mg/ml of the oil and water emulsion. An aliquot of 0.5 ml of the final solution ($\approx 25\text{mg/kg}$ of body weight) was fed to the infected mice every 3 to 4 days over two weeks with an appropriate gavage needle. Control preparation of free phenazines derivatives were given also on the same schedule to each mouse every day with the appropriate gavage needle. The animals were killed 24 h later after the last treatment. The lung, liver and spleen of the treated mice were homogenized (one part tissue and nine parts methanol/glacial acetic acid 9:1).

Determination of iminophenazines concentration: The concentration of each iminophenazines was extracted from the spleen and fat tissues of the mice using a mixture of dichloromethane and sodium hydroxide, quantified using a reversed phase C18 column and determined using the modified previous HPLC method (O'Connor et al., 1996). The HPLC analysis was carried using the Jasco - Borwin 310 model (Japan) and absorbance measurements were obtained using a multi channel UV- Visible detector at λ_{max} 384 nm.

RESULTS AND DISCUSSION

The linear regression analysis of the partition coefficients

Table 2. The ionization constant (pKa), degree of ionization (1- α) and the partition coefficients (log P) of the iminophenazines

Compound	pKa	log(1- α)	log P
B663	8.511	3.361	4.396
B826	8.813	3.663	4.496
B3779	8.662	3.512	4.533
B3770	8.800	3.650	4.782
B3785	8.592	3.446	4.380
B749	8.085	2.936	4.307
B3955	8.480	3.330	3.795

Table 3. The value of lipophilic parameter (π) obtained by comparing log P values of substituted derivative with that of parent compound (R= H)

Compound	π
B663	1.096
B826	1.196
B3779	1.233
B3770	1.482
B3785	1.080
B749	1.007
B3955	0.496

in the n-octanol/buffer system of seven iminophenazines with the concentration found in the spleen of mice following oral administration gave the following equations:

$$\log (1/C) = 1.165 \log P + 3.015 \text{ ----- equation 1}$$

$$t = 1.34 (P = 0.237) n = 7$$

$$r = 0.515 s = 0.398; F = 1.81; \text{significance } 76.3\%$$

The correlation is obviously not good, and similar results were obtained using the parabolic equation:

$$\log (1/C) = 23.630 \log P - 2.83 (\log P)^2 - 51.344 \text{ -----}$$

$$\text{equation 2}$$

$$t = 0.52 (P = 0.237) t = 0.54 (P = 0.237)$$

$$n = 7; r = 0.562 s = 0.430$$

These results indicate that lipophilicity of the new potent analogous is not by itself a significant physicochemical parameter in determining the transport of these compounds to the spleen. The high log P values of clofazimine and derivatives with polar N²- substituents using n-octanol/buffer system and the lipophilic parameter (π) (Tables 2 and 3) may reflect an extensive binding to lipoproteins, as the mechanism of absorption of the phenazines has not been established. Electrophoretic studies of the blood serum of orally treated mice with clofazimine have shown the binding of clofazimine to lipoproteins of the α - and β - globulin fraction. These lipoprotein-clofazimine complexes are taken up by the macrophages

and concentrated in the reticulo-endothelial system. Following fusion with liposomes, it appears that the lipoprotein carrier is enzymatically split off from the complex, with the resultant as the crystallization of clofazimine (Barry et al., 1960; Conalty et al., 1971). It has also been advanced that clofazimine inhibits the multiplication of organisms by binding to the guanine bases of DNA (Morrison and Marley, 1976).

However, correlations of protein binding with the polar properties of the substituents of certain drug molecules have been reported, and this showed that the extent of plasma protein binding is significantly dependent on the frontier electron density on the hetero-atom at the terminal portion of the alkyl side chains. The greater the frontier electron density, the greater was the degree of the drug binding to plasma protein (Austel and Kutter, 1983). Several studies have been conducted on the anti-microbial and the structure activity relationships of clofazimine analogues (Kumar, 2008; Huygens et al., 2005; Oliva et al., 2004; Scollard et al., 2006; Carta et al., 2008).

The relatively high concentration of phenazines with polar substituents at the N²- position recovered from the spleen of mice may be due to the contribution of the polar groups to protein binding. This finding prompted the calculations of the polar substituent constants for the alkyl amino and cycloalkyl amino groups at the N²- position.

It was recommended that the use of σ^* might give a better discrimination between the various substituents than σ_1 . The σ^* was proposed by Taft, (1956) for evaluating the polar effects of substituents R in the ester RCO₂ R'.

$$\sigma^* = [\log (K/K_0) B - \log (K/K_0) A] 2.48 \text{ ----- equation 3}$$

where σ^* is the polar substitute constants for R, the rate constants (K) refer to the reactions of RCO₂ R' and K₀ to those of CH₃-CO₂ R' standard. B and A refer to basic and acidic hydrolyses carried out with same R' solvent and temperature. The factor 2.48 puts the σ^* values into about the numerical range as Hammett's σ which has been used extensively in QSAR.

The linear regression analysis using σ^* values for the polar N²- substituents of six compounds and the concentration extracted from the spleen of mice following oral administration gave the following equations:

$$\log (1/C) = - 3.740 \sigma^* - 2.094 \text{ ----- equation 4}$$

$$t = 3.88 (P = 0.018); n = 6$$

$$r = 0.889 s = 0.238; \text{significance } 98.2\%$$

A combination of log P (n-octanol/buffer) and σ^* values gave the following regression equation:

$$\log (1/C) = 0.706 \log P - 3.217 \sigma^* + 1.032 \text{ ----- equation 5}$$

$$t = -0.84 (P = 0.465); t = -2.72 (P = 0.073)$$

$$\text{significance} = 53.3\%; \text{significance} = 92.7\%$$

$$n = 6; r = 0.911 s = 0.248; F = 7.286 (\text{sig.} = 92.7\%) A$$

Table 4. The σ^* of the polar N²- substituted iminophenazines.

Compound	σ^*
B663	-0.190
B826	0.040
B3779	0.055
B3770	0.042
B3785	0.045
B749	0.125

linear regression analysis using the lipophilic parameter for the six phenazines gave a poorer correlation ($r = 0.640$). It is clear from the results that the electronic properties of the N²-substituents, as expressed by their σ^* values are of great significance in relation to the transport of these compounds to the spleen of the mice than the lipophilicity of the molecules. The negative value of the coefficient of the σ^* term (Table 4) indicates the enhancement of the electron-donating substituents. These recent results indicate that the absorption process is affected by not only lipophilicity, and the absorption may be dependent on protein binding of these lipophilic molecules.

Conclusion

The recent results suggest that despite the high log P values, transport to the spleen where the compounds accumulate is not, apparently, a simple absorption process but is more likely to be dependent to a significant extent on protein binding, as these iminophenazines with polar substituents have a pKa values within a narrow range (8.66 - 8.80). This suggests their capabilities, therefore, of binding to proteins by electrostatic interactions.

REFERENCES

- Albert A, Serjeant EP (1984). Determination of ionization constants: A Laboratory Manual" Chapman and Hall. London. pp 118-134.
- Anderson NH, Davis SS, James M, Kojima I (1983). Thermodynamics of distribution of p-substituted phenols between aqueous solution and organic solvents and phospholipid vesicles. *J. Pharm. Sci.* 72: 443-448.
- Austel V, Kutter E (1983). Quantitative structure-activity relationships of Drugs" ed. John G. Academic press, New York, Chapter 11. pp.112-134.
- Barry VC, Bugle K, Byrne J, Conalty M, Winder F (1960). Absorption, distribution and retention of the Rimino- compounds in experimental animal. *Irish J. Med. Sci.* 416: 345-452.
- Beezer AE, Hunter WH, Storey DE (1980). Quantitative structure relationships: the Van't Hoff heat of transfer of resorcinol monoethers from water to n-octanol. *J. Pharm. Pharmacol.* 32: 815-819.
- Beezer AE, Hunter WH, Storey DE (1983). Enthalpies of solution of a series of m-alkoxy phenols in water, n-octanol, and water, n-octanol mutually saturated. *J. Pharm. Pharmacol.* 35: 350-357.
- Betageri GV, Rogers JA (1987). Thermodynamics of partitioning of β -blockers in n-octanol - buffer liposome system. *Int. J. Pharm.* 36: 165-173.
- Carta A, Piras S, Palomba M, Jabes D, Molicotti P, Zanetti S (2008). Anti-Mycobacterial Activity of Quinolones, Triazoloquinolones a New Class of Potent Anti-Mycobacterial Agents. 7: 134-147
- Conalty M, Barry V C, Jina A (1971). The ant leprosy agent B663 (Clofazimine) and the reticulo-endothelial system. *Int. J. Lepr.* 39: 479-484.
- Dearden JC, Bresnen GM (2005). Thermodynamics of water-octanol and water-cyclohexane partitioning of some aromatic compounds. *Int. J. Mol. Sci.* 6: 119-129.
- Degas H, Robert J, Bonnafous P, Jarlier V (2000). Activities of several Antimicrobial against Mycobacterium ulcerans Infection in Mice, *J. Antimicrob. Agents Chemother.* 44: 2367-2372.
- Fahelbom KMS, Quigley JM, Timoney RF, Corrigan OI (1989). Thermodynamics of the distribution and interfacial transfer of antimicrobial iminophenazines in two phase system of n-octanol/ aqueous buffer. *Proc. R. Ir. Acad.* 89B: 287-298.
- Hansch C, Leo A. (1979). Substituent constants for correlation analysis in Chemistry and Biology, Wiley, New York. pp. 49-55.
- Hansch C, Leo A (1995). Exploring QSAR; Fundamentals and applications in Chemistry and Biology, American Chemical Society. Washington DC. pp. 169-221
- Rogers JA, Davis SS (1980). Functional group contributions to the partitioning of phenols between liposomes and water. *Biochem. Biophys. Acta.* 598: 392-404.
- Huygens F, O'Sullivan JF, van Rensburg CE (2005). Antimicrobial activities of seven novel tetramethylpiperidine-substituted phenazines against multiple-drug-resistant Gram-positive bacteria. *Chemotherapy.* 51:263-267.
- Johnson T, Khan IA, Avery MA, Grant J, Meshnick S.R (1998). Quantitative structure-activity relationship studies of a series of sulfa drugs as inhibitors of pcarinii dihydropteroate synthetase. *J. Antimicrob. Agents Chemother.* 42: 1454-1458.
- Kellogg GE, Abraham, DJ (2000). Hydrophilicity: is Log Po/w more than the sum of its parts? *Eur. J. Med. Chem.* 35: 651-661.
- Kojima I, Davis SS (1984). Thermodynamics of distribution of p-substituted phenols between organic solvents and sodium chloride solution. *Int. J. Pharm.* 20: 247-264.
- Kumar V, (2008). Does clofazimine (B663) reach Mycobacterium leprae persisting in Schwann cells and endothelial cells of endoneurial blood vessels in peripheral nerves? *Microscopy Research and Technique.* 71: 614-618.
- Lien EJ, Hansch C, Anderson SM (1968). Structure-activity correlations for antibacterial agents on gram-positive and gram-negative cells. *J. Med. Chem.* 11: 430-441.
- Martin, YC (1979). Advances in the methodology of Quantitative Drug Design, Drug design. Vol.VIII, pp 324-352.
- Morrison NE, Marley GM (1976). The mode of action of clofazimine; DNA binding studies. *Int. J. Lepr.* 44: 133-134.
- O'Connor R, O'Sullivan S, O'Kennedy R (1996). An improved method for analysis clofazimine in biological samples using HPLC, *J Chromatogr. B.* 681: 307-315.
- Oliva B, O'Neill AJ, Miller K, Stubbings W, Chopra I (2004). Anti-staphylococcal activity and mode of action of clofazimine. *J Antimicrob Chemother.* 53: 435-440
- O'Sullivan, JF, Conalty M L, Morrison, NE (1988). Clofazimine analogues active against a clofazimine-resistant organism, *J. Med. Chem.* 31: 567-572.
- Thakur M, Agawal A, Thakur A, Khadikar, PV (2004). QSAR study on Phenolic activity: need of positive hydrophobic term (log P) in QSAR. *Bioorg. Med. Chem.* 12(9): 2287-2293.
- Scollard, DM, Adams, LB, Gillis TP, Krahenbuhl JL, Truman RW, Williams DL (2006). The Continuing Challenges of Leprosy, *Clin Microbiol. Rev.* 19: 338-381.
- Shepard CC (1967). A kinetic method for the study of the activity of drugs against Mycobacterium Leprae. *Int. J. Lepr.* 35: 429-436.
- Taft RW (1956). Steric Effects in Organic Chemistry; Newman MS, Ed.; Wiley: New York. pp. 556 -577.
- Warhurst D, Craig JC, Adagu IS, Lee, DJ (1998). The relationship of physico-chemical properties at different antiparasitodal activity of the cinchona. *Malar. J.* 2: 26- 41.