

Full Length Research Paper

Effects of food spices on Gram-negative food indicator bacteria from some Nigerian ethnic fermented plant food condiments

Adenike A. O. Ogunshe^{1*}, Zibi M. Johnny² and Amala O. Arinze²

¹Applied Microbiology and Infectious Diseases Unit, Department of Microbiology, Faculty of Science, University of Ibadan, Ibadan, Nigeria.

²Department of Botany and Microbiology, Faculty of Science, University of Ibadan, Ibadan, Nigeria.

Accepted 11 October, 2011

Varied but multiple antibiotic resistance rates were exhibited by all the foodborne indicator bacterial species, *Citrobacter*, *Escherichia coli*, *Enterobacter*, *Klebsiella*, *Morganella morganii*, *Salmonella*, *Shigella*, *Pseudomonas* isolated from four most-popular Nigerian indigenous fermented plant food condiments *ogiri*, *iru*, *okpehe* and *ugba*. The generally most-resisted antibiotics (discs) were tetracycline (44.0 - 63.0%), cotrimoxazole (56.0 - 70.3%), nalidixic acid (28.0 - 74.1%), amoxicillin (28.0 - 88.9%) and augmentin (16.0 - 94.4%), while the least resisted antibiotics were ofloxacin (0.0% - 1.85%) and gentamicin (4.0 - 16.7%). Antibacterial activities of crude extracts of local spices and essential oils of plant origin, as well as carvone, lactic and acetic acids on the multiple antibiotic resistant Gram-negative foodborne bacterial species were determined using modified agar well-diffusion assay. Crude extracts of *Eugenia aromatica* (92.0%) and *Allium sativum* (72.0%) were maximally inhibitory; lactic (44.4%) and acetic (46.3%) acids were moderately inhibitory but essential oils of *Eugenia uniflora* (24.0%), *Ageratum conyzoides* (16.7%) and *Chrysophyllum albidum* juice (13.0%) were minimally inhibitory *in vitro*, indicating that crude extracts of *E. aromatica* and *A. sativum* can serve as easily prepared, non-chemical, plant-based, adjunct preservatives of fermented condiments both domestically and industrially, even by the traditional producers of the indigenous and similar condiments.

Key words: Cottage food production, essential oils, food safety, food processing, multiple antibiotic resistance, Nigerian indigenous fermented foods, plant foods.

INTRODUCTION

New strategic research areas have emerged and been developed, with profound effects on the capacity to produce food and manage natural resources and the environment (Taeymans, 2011), while the world Health Organisation food safety unit has also given high priority to the research area of fermentation as a technique for preparation and storage of food. In Africa and Asia, the

art of fermentation is widespread, including the processing of plant vegetable seeds into food condiments, which in Nigeria and many other countries of west and central Africa are popular strong-smelling fermented food culinary products that give pleasant aroma to soups, sauces and other prepared dishes.

It is evident that fermented food condiments are good sources of nutrients and could be used to produce complementary food supplements (Achi, 2005); however, it is quite obvious that fermented food condiments may harbour some bacterial pathogens that can be hazardous to human health. Nevertheless, the demand for locally

*Corresponding author. E-mail: adenikemicro@yahoo.com. Fax: (234)-2-8103043.

processed Nigerian fermented food condiments (NIFFC) such as *iru*, *ogiri*, *ugba*, *okpehe*, etc. (from seeds of *Parkia biglobosa*, *Citrullus vulgaris*, *Pentaclethra macrophylla* and *Prosopis africana* respectively) is currently on the increase; even among the elites, probably because none of the industrial seasonings or spices has been able to meet the indigenous organoleptic qualities of these fermented condiments of plant origin. Microorganisms have successfully adapted to changes in food production, processing and preservation techniques, thereby resulting in a number of new and emerging foodborne pathogens, as well as re-emergence of the ones that have been problematic in the past. To protect the public health, science must therefore, meet the challenges that result from the remarkable adaptability of foodborne pathogens to foods (Woteki and Kineman, 2003). In addition to recovery of foodborne bacterial species from these condiments, there is also the emergence of multiple antibiotic resistant bacteria, which pose a challenge in food systems (White et al., 2002, 2004). Food safety therefore, remains a major challenge to producers and consumers of these fermented food condiments. Consumers are also currently skeptical of chemical preservatives in their foods; thus, the need to discover other more acceptable means of protecting and preserving these highly popular indigenous foods becomes pertinent, more so since plant extracts are more natural.

Certain plant-based spices have been found to impart flavour and prolong the storage life of foods by bacteriostatic or bactericidal activities (López et al., 2007; Shan et al., 2007). Being natural foodstuffs, they also appeal to consumers who tend to question the safety of synthetic / chemical food additives (Vanit et al., 2010). The purpose of this study therefore, was to determine the *in vitro* antibiotic resistance patterns of Gram-negative, foodborne indicator bacterial species isolated from four most popular Nigerian fermented food condiments, *iru*, *ogiri*, *ugba*, *okpehe*. Using an easily reproducible method, the inhibitory and preservative activities of some local spices, organic acids and essential oils against the foodborne bacterial pathogens isolated from the fermented food condiments shall be evaluated.

MATERIALS AND METHODS

Collection of samples

The Nigerian fermented food condiments used in this study were market *iru* produced (from locust bean, *Parkia biglobosa*), *ogiri* (from melon seeds, *Citrullus vulgaris*), *okpehe* (from *Prosopis africana*) and *ugba* (from oil-bean seeds, *Pentaclethra macrophylla*), while the fresh food spices (*Allium sativum*, *Eugenia aromatica*, *Monodora fragrans*, *Myristica fragrans*, *Piper guineensis*, *Xylopiya aethiopica* and *Zingiber officinalis*) were all purchased from various markets in South-West Nigeria. The spices, two organic acids (lactic and acetic acids) and essential oils of

Ageratum, *Eugenia*, *Carvone*, *Anacardium* plants and *Chrysophyllum albidium* juice were assayed for their inhibitory effects on the foodborne pathogens from the four fermented condiments in this study. The condiments were obtained at 12-month interval between the first samples' collection (A) and the second samples' collection (B).

Isolation, characterisation and preservation of Gram-negative bacteria from fermented condiments

The Gram-negative bacteria isolates were selectively recovered from the various fermented food condiment samples, using sterile MacConkey broth and agar (MCC, Lab M, England), *Salmonella-Shigella* agar (SS, Lab M, England), *Pseudomonas* selective agar (BHI, Oxoid, England), thiosulphate citrate bile sucrose agar (TCBS, Lab M, England), while differential identification was determined on eosin methylene blue (EMB, Lab M, England) and triple sugar iron agar (TSI, Lab M, England). All the pure bacterial isolates were kept in triplicates on Brain Heart Infusion (BHI, Oxoid, England) agar slants, as working and stock cultures. Taxonomic studies were carried out on the purified bacterial isolates on the basis of their cultural, morphological, biochemical and physiological characteristics. General keys with reference were used for the identification (Buchanan and Gibbons, 1974; Crichton, 1996).

Antibiotic susceptibility determination of the Gram-negative bacteria

Antibiotic susceptibility / resistance of the Gram-negative foodborne bacterial species to various antibiotics AMX (Amoxicillin; 25 µg), COT (Cotrimoxazole; 25 µg), NIT (Nitrofurantoin; 250 µg), GEN (Gentamicin; 10 µg), NAL (Nalidixic acid; 30 µg), OFL (Ofloxacin; 30 µg), AUG (Augmentin; 30 µg) and TET (Tetracycline; 30 µg) was determined according to the Bauer et al. (1996) and NCCLS (2003) methods. The entire surface of each sterile Mueller-Hinton agar plate was seeded with each Gram-negative bacteria isolate using sterile swab sticks, after which the plates were left for about 15 min before aseptically placing the antibiotic discs on the agar surfaces with sterile forceps, followed by incubation at 35°C for 18 to 24 h. Zones of inhibition around the agar wells after incubation were measured and recorded in millimeter diameter, while zones of inhibition less than 10.0 mm in diameter or absence of zones of inhibition were recorded as resistant or negative.

In vitro antimicrobial inhibition bioassays of the Gram-negative bacteria isolates by local spices

Fresh local food spices (*Z. officinalis*, *A. sativum*, *X. aethiopia*, *P. guineensis*, *M. fragrans*, *M. fragrans*, *E. aromatica*, *Aframomum melegueta*) were surface-sterilised with ethanol and later cleaned with wet, sterile cotton wool. In order to obtain the spice's extracts, about 150 g of each cleaned spice were dry-milled with blender, which was hot-washed (boiled water at 100°C) five times after each use, based on the microbial loads of the rinses. A set of the milled extracts were soaked in warm sterile distilled water for 12 h to obtain the aqueous extracts of the spices and the other set were slightly wet with sterile distilled water to obtain the crude extracts (pastes) of the spices. About 10 ml of *C. albidium* juice was obtained by squeezing out the juice into 15 ml of sterile distilled water, while the essential oils of *Ageratum conyzoides*, *Eugenia uniflora*, *Anacardium occidentals*, *carvone* used in this study were prepared in the organic unit of the Department of Chemistry, University of Ibadan. The organic acids - lactic and acetic acids

Table 1. Percentage *in vitro* antibiotic resistance patterns of Gram-negative foodborne bacteria species isolated from fermented food condiments (A).

S/N	Bacterial isolate	Antibiotic disc (μg^{-1})							
		AMX	COT	NIT	GEN	NAL	OFL	AUG	TET
1	<i>Enterobacter</i> spp. [8]	12.5	50	0	12.5	25	0	12.5	37.5
2	<i>Klebsiella</i> spp. [6]	50	66.7	33.3	0	33.3	0	16.7	66.7
3	<i>Citrobacter</i> spp. [3]	33.3	33.3	0	0	66.7	0	33.3	100
4	<i>Salmonella</i> sp. [5]	0	80	0	0	0	0	0	20
5	<i>Ps. aeruginosa</i> [3]	66.7	33.3	33.3	0	33.3	0	33.3	0
No. / % Total resistance		7 (28.0)	14 (56.0)	3 (12.0)	1 (4.0)	7 (28.0)	0 (0.0)	4 (16.0)	11 (44.0)

Amx = amoxicillin; Cot = cotrimoxazole; Nit = nitrofurantoin; Gen = gentamicin; Nal = nalidixic acid; Ofi = ofloxacin; Aug = augmentin; Tet = tetracycline.

were BDH products.

In vitro antimicrobial inhibition bioassay on the Gram-negative bacteria species by the local spices, essential oils and organic acids was determined by the modification of the agar well-diffusion method of Tagg and McGiven (1971), in which 0.5% agar powder dissolved in appropriate quantity of water was sterilised and added (40°C) to the aqueous extracts of the spices, essential oils and organic acids to avoid spreading of the extracts and acids on the surface of the seeded agar plates. Using sterile cork borers, 6.0 mm wells in diameter were bored into sterile Mueller-Hinton agar plates followed by surface flaming of the agar surfaces. The entire surface of each sterile plate was seeded by streaking with each selected Gram-negative bacterial strains (previously inoculated into sterile tryptone soy broth and incubated at 37°C for 18 to 24 h). 500 μl of the different aqueous extracts and 0.5 gm of the crude extracts of the local spices, essential oils and organic acids were dispensed into each set of agar wells and then incubated un-inverted at 35°C for 24 to 48 h. Zones of inhibition surrounding the wells after incubation were measured and recorded in millimeters (mm) diameter, while wells with no inhibition zones or less than 10.0 mm in diameter were recorded as resistant.

RESULTS

Varied antibiotic resistance patterns recorded among the individual bacterial species and the overall phenotypic antibiotic resistance patterns exhibited by the foodborne indicator bacteria from both sets A and B were as shown in Tables 1 and 2. The foodborne bacteria from the first set (A) of sampled fermented food condiments exhibited lower overall antibiotic resistance towards the test antibiotics- nitrofurantoin (12.0%), augmentin (16.0%), amoxicillin and nalidixic acid (28.0%), with moderate antibiotic resistance exhibited towards tetracycline (44.0%) and cotrimoxazole (56.0%). All the bacterial isolates were however, susceptible to ofloxacin. The most resisted antibiotics by the bacterial flora from set B of the sampled fermented food condiments were tetracycline (63.0%), cotrimoxazole (70.3%), nalidixic acid (74.1%), amoxicillin (88.9%) and augmentin (94.4%) but lowest resistance was recorded towards ofloxacin (1.85%).

In set (A), relatively lower antibiotic resistance patterns were recorded- *Enterobacter* strains (12.5-50.0%) except in nitrofurantoin and ofloxacin (0.0%); *Klebsiella* and *Pseudomonas* strains (16.7 – 66.7% and 33.3 to 66.7%) respectively except in gentamicin and ofloxacin, as well as tetracycline in *Pseudomonas* strains (0.0%); *Citrobacter* strains (33.0 to 100.0%) except towards nitrofurantoin, gentamicin and ofloxacin (0.0%). Most of the *Salmonella* strains were susceptible to the test antibiotics but resistance rates of between 20.0 and 80.0% were still recorded among them towards tetracycline and cotrimoxazole respectively (Table 1). Multiple antibiotic resistance (MAR) was however, exhibited by all the bacterial species. Among the bacteria isolated in set B, low to very high phenotypic antibiotic resistance patterns were recorded in *Escherichia coli* strains (27.3 to 90.9%) except against ofloxacin (0.0%); *Salmonella* strains (71.4 to 100.0%) except against gentamicin and ofloxacin (0.0%); *Shigella* strains (16.7 to 100.0%) except against nitrofurantoin (0.0%); *Citrobacter* strains (22.2 to 100.0%) and *Klebsiella* strains (23.0 to 100.0%) except against ofloxacin (0.0%); *Enterobacter aerogenes* strains (28.6 to 85.7%) and *Morganella morganii* except against gentamicin and ofloxacin (0.0%) (Table 2).

In this study, the foodborne bacteria from the four types of fermented food condiments of plant origin were assayed *in vitro* against some local Nigerian spices to determine the possibility of the spices being used as domestic and industrial, non-chemical preservatives but only the crude extracts of the spices were inhibitory *in vitro* against the foodborne Gram-negative bacteria, while the aqueous extracts of the spices were non-inhibitory or exhibited very minimal inhibitory activities *in vitro*. Meanwhile, the overall *in vitro* susceptibility patterns of crude extracts of the spices indicated that the foodborne bacteria were mostly susceptible to *E. aromatic* and *A. sativum* but mostly resistant to *M. fragrans*, *M. fragrans*, *P. guineensis*, *X. aethiopia*, *Z. officinalis* and *A. megueta* (Table 3).

Table 2. Percentage *in vitro* antibiotic resistance patterns of Gram-negative food borne bacterial species isolated from fermented food condiments (B).

S/N	Bacterial isolate	Antibiotic disc (μg^{-1})							
		AMX	COT	NIT	GEN	NAL	OFL	AUG	TET
1	<i>E. coli</i> [11]	81.8	72.7	36.4	27.3	63.6	0.0	90.9	45.5
2	<i>Salmonella</i> spp. [7]	100.0	85.7	85.7	0.0	100.0	0.0	100.0	71.4
3	<i>Shigella</i> spp. [6]	83.3	16.7	0.0	16.7	50.0	16.7	100.0	50.0
4	<i>Citrobacter</i> [9]	100.0	77.8	44.4	22.2	88.9	0.0	100.0	77.8
5	<i>Klebsiella</i> spp. [13]	92.3	84.6	61.5	23.0	76.9	0.0	100.0	84.6
6	<i>Ent. aerogenes</i> [7]	85.7	71.4	28.6	0.0	71.4	0.0	85.7	42.9
7	<i>Morg. morganii</i> [1]	100	100	100	0.0	100	0.0	100	100
No. / % Total resistance		48 (88.9)	38 (70.3)	24 (44.4)	9 (16.7)	40 (74.1)	1 (1.85)	51 (94.4)	34 (63.0)

AMX = Amoxicillin; COT = Cotrimoxazole; NIT = Nitrofurantoin; GEN = gentamicin; NAL = Nalidixic acid; OFL = Ofloxacin; AUG = Augmentin; TET = Tetracycline. Values in parenthesis are percentage resistance.

Table 3. Percentage *in vitro* inhibition rates of crude extracts of spices against Gram-negative food borne bacterial species isolated from fermented food condiments

S/N	Lab codes of isolate	Crude extract of spices (g ml^{-1})							
		ZGO	ALS	XYA	PPG	MYF	MOF	EUA	AMG
1	<i>Enterobacter</i> [8]	0.0	87.5	0.0	25.0	0.0	0.0	87.5	12.5
2	<i>Klebsiella</i> [6]	16.7	66.7	16.7	0.0	0.0	0.0	100.0	33.3
3	<i>Citrobacter</i> [3]	0.0	100.0	0.0	33.3	0.0	0.0	66.7	0.0
4	<i>Salmonella</i> spp. [5]	0.0	40.0	20.0	0.0	0.0	0.0	100.0	0.0
5	<i>Pseudomonas</i> [3]	0.0	66.7	0.0	33.3	0.0	0.0	100.0	0.0
No. / % Total susceptibility		1 (4.0)	18 (72.0)	5 (20.0)	4 (16.0)	0 (0.0)	0 (0.0)	23 (92.0)	3 (12.0)

ZGO = *Zingiber officinalis*; ALS = *Allium sativum*; XYA = *Xylopiya aethiopia*; PPG = *Piper guineensis*; MYF = *Myristica fragrans*; MOF = *Monodora fragrans*; EUA = *Eugenia aromatica*; AMG = *Aframomum melegueta*.

An overall total of 44.4 and 46.3% of the bacterial strains from the fermented food condiments were inhibited by lactic acid and acetic acid respectively (Table 4), while 24.0, 16.7 and 13.0% of the bacteria were minimally inhibited by the essential oils of *E. uniflora*, *A. conyzoides* and *C. albidium* juice respectively (Table 4).

DISCUSSION

Foodborne pathogens have been found to be widely distributed in nature and causing considerable mortality and morbidity in the population (Indu et al., 2006); so, authors have started to focus on the study of survival and growth of foodborne pathogens including some of the indigenous / ethnic fermented foods and beverages of plant origin (Odibo et al., 1992; Ogunshe et al., 2006, 2007, Ogunshe and Olasugba, 2008). As earlier reported, the most commonly encountered bacterial pathogens in the Nigerian fermented food condiments include *Bacillus cereus*, *E. coli*, *Salmonella* sp., *Staphylococcus aureus*, *Vibrio cholerae*, *Aeromonas*, *Klebsiella*, *Campylobacter*

and *Shigella* spp., *Alkaligenes viscolactis*, *Corynebacterium* spp., *Enterobacter cloacae*, *Citrobacter cloacae*, *E. aerogenes*, *Pseudomonas aeruginosa*, *Proteus mirabilis* and *Micrococcus* sp. (Kolawole and Okonkwo, 1985; Nwafor, 1985; Barber et al., 1988). However, these Gram-negative foodborne bacteria have been implicated in diseases of public health importance, such as gastroenteritis (Ogunshe and Olasugba, 2008; Jiva et al., 1988; Salyers and Whitt, 1994; Tauxe, 2002), and have become increasingly important as causes of diarrhoeal diseases.

In addition to the health implications of the survival of these food indicator bacteria in the Nigerian food condiments, the issue of their antibiotic multi-resistance is also of great concern. According to Kiessling et al. (2002), antibiotic resistance in foodborne pathogens is a reality and recovery of strains of resistant foodborne pathogens to a variety of antimicrobials has become a major health concern, and similarly, the rise in carriage of antibiotic genes in virtually every species of bacterium has also been consistently documented. In this study, the Gram-negative bacteria species from the fermented food

Table 4. Percentage *in vitro* inhibition rates of organic acids and essential oils on Gram-negative bacterial species from fermented condiments.

S/N	Bacterial isolate	Organic acids			Essential oils			
		LA	AA	AGR	EUG	ANC	CAR	CHR
1	<i>E. coli</i> [11]	54.5	63.6	9.09	9.09	0.0	0.0	0.0
2	<i>Shigella dysenteriae</i> [6]	33.3	16.7	16.7	0.0	0.0	0.0	33.3
3	<i>Citrobacter</i> sp. [9]	66.7	66.7	22.2	44.4	11.1	11.1	11.1
4	<i>Salmonella</i> sp. [7]	28.6	28.6	14.3	0.0	14.3	14.3	14.3
5	<i>Klebsiella aerogenes</i> [13]	38.5	46.2	15.4	30.8	15.4	7.7	7.7
6	<i>Ent. aerogenes</i> [7]	28.6	42.9	28.6	57.1	0.0	0.0	28.6
7	<i>Morg. morganii</i> [1]	1	0	0	0	0	0	0
No. / % Total susceptibility		24 (44.4)	25 (46.3)	9 (16.7)	13 (24.0)	4 (7.4)	3 (5.56)	7 (13.0)

LA = lactic acid; AA = acetic acid; CHR = *Chrysophyllum albidum*; AGR = *Ageratum conyzoides*; EUG = *Eugenia uniflora*; ANC = *Anacardium occidentale*; CAR = carvone.

condiments were also found to exhibit antibiotic resistance, and in addition, almost all the bacterial strains were found to be resistant to multiple antibiotics. This is of considerable medical significance because they may also serve as hosts of antibiotic resistance genes that can be transferred to other bacteria and ultimately humans who consume such foods that harbour them. Search for new antimicrobials is therefore, very important in recent times, considering the escalating levels of antibiotic resistance among the foodborne bacteria.

Several food additives, such as herbs and spices, and which are generally regarded as safe (GRAS), are used to delay or inhibit the growth of pathogenic microorganisms and also to extend the shelf life of food (Zaika, 1988; Arora and Kaur, 1999; Newberne et al., 2000). In this study, crude extracts of garlic (*A. sativum*) and clove (*E. aromatica*) exhibited 72.0 and 92.0% *in vitro* inhibitory activities respectively against the Gram-negative foodborne bacterial species. These results agree with observations of previous researchers in which garlic was reported of having inhibitory activities against *E. coli* O157:H7, *Salmonella typhimurium*, *E. coli*, *S. aureus*, *B. cereus*, *Bacillus subtilis*, mycotoxigenic *Aspergillus* and *Candida albicans*. Moderate activities of ginger extract on *S. paratyphi* and *S. enteritidis*; onion extract against *E. coli*, *S. paratyphi* and *S. typhimurium* and antibacterial activities of extracts of garlic (*A. sativum*), nutmeg (*M. fragrans*), ginger (*Z. officinale*), onion (*Allium cepa*) and pepper (*Piper nigrum*) against different strains of *E. coli* and *Salmonella* [Dankert et al., 1979; Sasaki et al., 1979; Arora and Kaur, 1999; Indu et al., 2006].

Contrary to the studies of Indu et al. (2006), extracts of ginger (4.0%) and pepper (16.0%) did not show significant antibacterial activity against the test bacterial strains in the present study; although results obtained indicated that the *in vitro* inhibitory activities of the spices were not bacterial species dependent. Also as observed

in the current study, inhibitory activities were prominent in the crude extracts of the spices, which support the earlier assertions of Shelef (1983) that in general, higher spice levels are required to effect inhibition in foods than in culture media, and although the inhibitory activities of some spices and herbs are documented, the normal amounts added to foods for flavour is not sufficient to completely inhibit microbial growth because the antimicrobial activities vary widely, depending on the type of spice or herb, test medium and microorganisms.

In this study as well, the foodborne bacterial species from the fermented food condiments were found to be partially inhibited *in vitro* by lactic and acetic acids. Other studies which reported antibacterial potentials / bactericidal effects of organic acids include those of Cherrington et al. (1991), Hirshfield et al. (2003). Plants essential oils and phenolic secondary metabolites possess antimicrobial activities (Prabuseenivasan et al., 2006), and some of these plant extracts are used in the aroma and flavor industry, and are also generally regarded as safe (Shelef, 1983). Based on the result findings of the present study, it was observed that the addition of some commonly obtainable spices such as *A. sativum* and *E. aromatica* to fermented condiments can serve as biopreservatives against foodborne bacterial pathogens to a larger extent but the organoleptic results (results not shown) indicated that addition of *E. aromatica* to the fermented condiments was preferred to *A. sativum* in supporting dishes such as *ewedu*, a Nigerian soup prepared from *Corchorus olerarius* L. and *ila*, a Nigerian soup prepared from okra (*Abelmoschus esculentus* (L.) Moench), while the addition of *A. sativum* in minimal quantities was preferred in main and supporting stews like various vegetable soups such as *apon / ogbonno*, another Nigerian soup prepared from *Irvingia gabonensis*, as well as *ikokore* (a Nigerian indigenous delicacy prepared from bitter yam (*Dioscorea dumetorum* (Kunth) Pax) and popular among the Ijebus).

This indicates that *A. sativum* was acceptable in stews, soups and dishes that can mask its smell.

There are several industrial food spices in Nigeria, but they are mostly salt and aroma based; therefore, the popularities of the indigenous condiments such as *ogiri*, *iru* / *dawadawa*, *ugba*, *okpehe* etc. cannot be compared with the less desired industrial food spices (Ogunshe et al., 2008a, b). Consumers are so traditionally inclined to their ethnic fermented foods and beverages, such that it is very difficult to accept industrial alternatives to these indigenous foods. About two decades ago, an industrial condiment named *dadawa* was launched into the Nigerian markets with very captivating advertisements but less than a year afterwards, it was out of market because of its non-acceptance by the general populace, and till date, no industrially-prepared fermented condiment has been prepared for the Nigerian market. Consumers demand convenient, innovative, fresh foods, including new "minimally processed" products, while the food industry will utilise novel technologies to provide the new quality attributes demanded by consumers and ensure the all-important and expected assurance of food safety (Taeymans, 2011). The popularity of the Nigerian indigenous plant-based fermented condiments is specifically due to their organoleptic properties, that is, their aroma, flavour, taste and colour; though, few of the elites do not like these nutrient rich food products because of their pungent smell prior to cooking.

A sensory evaluation panel preferred the uncooked fermented condiments that were preserved with the essential oils in their evaluation for smells (results not shown). Therefore, the addition of such well-accepted local spices as, *A. sativum* and *E. aromatica*, essential oils of some spices and some natural organic acids, which have inhibitory characteristics on foodborne indicator bacteria incorporated as non-chemical, adjunct preservatives to domestically or industrially prepared condiments can alleviate the fear of non-safety of such enriched, indigenous fermented food products of plant origin. Addition of salt is the preservative currently employed by the traditional producers but it has not been effective in inhibiting the foodborne pathogens; thus, suggestion of adding crude plant extracts having inhibitory properties to the prepared condiments is imperative.

REFERENCES

- Achi OK (2005). The potential for upgrading traditional fermented foods through biotechnology. *Afr. J. Biotechnol.*, 4 (5): 375-380.
- Arora DS, Kaur J (1999). Antimicrobial activity of spices. *Int. J. Antimicrob. Agents.* 12(3): 257-262.
- Barber L, Achiewu SC, Ibiawa EA (1988). The microbiology of ogiri production from castor seed (*Ricinus commmunis*) In: Food microbiology, 4th Ed., Acad. Press. New York, pp. 177-182.
- Bauer AW, Kirby WM, Sherris JC, Turck M (1996). Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.*, 45: 493-496.
- Buchanan RE, Gibbons NE (1974). *Bergey's Manual of Determinative Bacteriology* (8th Ed.). Williams and Wilkins Comp. Baltimore, U.S.A.
- Cherrington CA, Hinton M, Mead GC, Chopra I (1991). Organic Acids: Chemistry, Antibacterial Activity and Practical Applications. *Adv. Microb. Physiol.*, 32: 87-108.
- Crichton PB (1996). Enterobacteriaceae: *Escherichia*, *Klebsiella*, *Proteus* and other genera. In Collee JG et al., eds. Mackie and McCartney practical Medical Microbiology, 14th ed. Baltimore, Churchill Livingstone, p. 161.
- Dankert J, Tromp TFJ, De Vries H, Klasen HJ (1979). Antimicrobial activity of crude juices of *Allium ascalonicum*, *Allium cepa*, and *Allium sativum*. *Zentrabl. Bakteriol. Parasitenkd. Infektkrankh. Hyg. Abt. 1 Org.*, 245: 229-239.
- Hirshfield IN, Terzulli S, O'Byrne C (2003). Weak organic acids: A panoply of effects on bacteria. *Sci. Prog.*, 86 (Pt 4): 245-269.
- Indu MN, Hatha AAM, Abirosh C, Harsha U, Vivekanandan G (2006). Antimicrobial activity of some of the South-Indian spices against Serotypes of *Escherichia coli*, *Salmonella*, *Listeria monocytogenes* and *Aeromonas hydrophila*. *Braz. J. Microbiol.*, 37: 2
- Jiva SFH, Krovacek K, Wadstom T (1988). Enterotoxigenic bacteria in food and water from an Ethiopian community. *Appl. Environ. Microbiol.*, 41: 1010-1019.
- Kiessling CR, Cutting JH, Lofis MK, Kiessling VW, Data AR, Sofos JN (2002). Antimicrobial resistance of food retail *Salmonella* isolates. *J. Food Protect.*, 65: 603-608.
- Kolawole DO, Okonkwo BA (1985). Microbiology of traditional fermented oil bean (*Pentaclethra macrophylla* Benth) seeds for ukpaka, a Nigerian condiment. *Nig. Food J.*, 3: 149-152.
- López P, Sanchez C, Battle R, Nerin C (2007). Vapor-phase activities of cinnamon, thyme, and oregano essential oils and key constituents against foodborne microorganisms. *J. Agr. Food Chem.*, 30: 55(11): 4348-4356.
- National Committee for Clinical Laboratory Standards (2003). Performance standards for antimicrobial disk susceptibility tests. 8th Ed., 23: 37-50.
- Newberne P, Smith RL, Doull J, Feron VJ (2000). GRAS Flavouring substances. *Food Technol.*, 54: 66-83.
- Nwafor UV (1985). A detailed study on the microflora of iru, a locally fermented African food condiment. *Nig. J. Microbiol.*, 5: 118-125.
- Odibo FJC, Ugwu DA, Ekeoha DC (1992). Microorganisms associated with the fermentation of Prosopis seeds for ogiri-okpei production. *J. Food Sci. Technol.*, (Mysore) 29: 306-307.
- Ogunshe AAO, Olasugba KO (2008). Microbial loads and incidence of food-borne indicator bacteria in most popular indigenous fermented food condiments from Middle-Belt and Southwestern Nigeria. *Afr. J. Microbiol. Res.*, 2: 332-339.
- Ogunshe AAO, Ayodele AI, Okonko IO (2006). Microbial studies on aisa, a potential indigenous laboratory fermented food condiment from *Albizia saman* (Jacq.) F. Mull. *Pakist. J. Nutr.*, 5 (1): 51-58.
- Ogunshe AAO, Omotoso MO, Ayansina ADV (2007). Microbial studies and biochemical characteristics of controlled fermented ajiyo- A Nigerian fermented food condiment from *Prosopis africana* (Guill and Perr.) Taub. *Pakist. J. Nutr.*, 6(6): 620-627.
- Ogunshe AAO, Jayeola AA, Ogundimu TC (2008b). Microbial studies on laboratory fermentation of iregi - A potential food condiment from *Delonix regia* (Boj. ex Hook.) Raf *FOOD.*, 2 (1): 61-64.
- Ogunshe AAO, Obiora-Okeke CM, Olurin TO (2008a). Bioinhibition of phenotypes of Gram-negative food indicator bacteria from some Nigerian fermented food condiments. *FOOD*, 2(2): 152-158.
- Prabuseenivasan S, Jayakumar M, Ignacimuthu S (2006). *In vitro* antibacterial activity of some plant essential oils. *BMC Complem. Altern. Med.*, 6: 39.
- Salyers AA, Whitt DD (1994). *Bacterial pathogenesis: A molecular approach*, Washington, D.C., ASM Press, USA.
- Sasaki J, Kita T, Ishita K, Uchisawa H, Matsue H (1999). Antibacterial activity of garlic powder against *Escherichia coli* O157. *J. Nutr. Sci. Vitaminol.*, (Tokyo) 45(6): 785-790.
- Shan B, Cai YZ, Brooks JD, Corke H (2007). The *In vitro* antibacterial activity of dietary spice and medicinal herb extracts. *Int. J. Food*

- Microbiol., 117(1): 112-119.
- Shelef LA (1983). Antimicrobial effects of spices. *J. Food Saf.*, 6: 29-44.
- Taeymans D (2011). New technologies for ensuring the quality, safety and availability of food. In: *Food, Nutrition and Agriculture*. FAO Corporate Document Repository, Agriculture and Consumer Protection. <http://www.fao.org/docrep/003/x7133m/x7133m04.htm> Accessed March 30, 2011.
- Tagg JR, McGiven AR (1971). Assay system for bacteriocins. *Appl. Microbiol.*, 21: 943.
- Tauxe RV (2002). Emerging foodborne pathogens. *Int. J. Food Microbiol.*, 78: 31-41.
- Vanit S, Suppakul P, Jinkarn T (2010). Antimicrobial effects of coating solution containing clove oil and hydrophobic starch for coating paperboard. *Asian J. Food Agro-Ind.*, 3(2): 204-212.
- White DG, Zhao S, Simjee S, Wagner DD, McDermott PF (2002). Antimicrobial resistance of foodborne pathogens. *Microb. Infect.*, 4(4): 405-412.
- White DG, Zhao S, Singh R, McDermott PF (2004). Antimicrobial resistance among Gram-negative foodborne bacterial pathogens associated with foods of animal origin. *Foodborne Pathol. Dis.*, 1(3): 137-152.
- Woteki CE, Kineman BD (2003). Challenges and approaches to reducing foodborne illness. *Ann. Rev. Nutr.*, 23: 315-344.
- Zaika LL (1988). Spices and herbs: Their antimicrobial activity and its determination. *J. Food Saf.*, 9: 97-118.