

Review

A review of the farming of common carp (*Cyprinus carpio* L.) in Malawi: Policy research directions for aquaculture development in Malawi

E. R. Chirwa^{1, 2*}, D. Kassam², W. L. Jere² and A. Mtethiwa²

¹Mzuzu University, P/Bag 201, Luwingu, Mzuzu 2, Malawi.

²Lilongwe University of Agriculture and Natural Resources (LUANAR), P. O. Box 219, Lilongwe, Malawi.

Received 10 April, 2017; Accepted 15 May, 2017

The lack of better performing native fish species for aquaculture led the government of Malawi to import the exotic common carp (*Cyprinus carpio* L.) from Israel in 1976. Growth trials at Domasi and Kasinthula Experimental Stations had shown that common carp grew faster and to a larger size than the indigenous fish species. The government decided to distribute the fish to farmers for grow-out. Barely five years into common carp distribution to farmers, the government reversed its policy and banned the use of the species in aquaculture. The government not only became unpopular but also lost the confidence of the farmers who had begun to see positive impacts of common carp to their livelihoods. The farmers are as unconvinced today as they were before with the reasons behind the banning of common carp. This paper explores the background to common carp farming in Malawi, why the fish was later banned, and the impacts of the fish's ban on the status of Malawi's aquaculture. The paper further highlights the farmers' call for a return of common carp to Malawi's aquaculture and the research needed to be undertaken to inform government's policy for the development of a sustainable aquaculture industry in Malawi.

Key words: Aquaculture, common carp, fish introduction, exotic fish, Malawi.

INTRODUCTION

Fish is the most affordable source of animal protein in Malawi, contributing over 70% of animal protein to the diet of Malawians (Mahony et al., 2014; Chidammodzi et al., 2015; Sanudi et al., 2015). Most of the fish consumed by Malawians come from capture fisheries. Aquaculture contributes about 2% to the total fish supply in Malawi. Over the past decades, fish production from capture fisheries has plateaued, with little or no prospect for

further expansion (Weyl et al., 2010). Overfishing and weak enforcement of fisheries regulations have been blamed for dwindling catch rates from lakes and rivers. With the increasing human population (growing at 3% p.a.), the scarcity of fish in Malawi has had many ramifications.

Foremost is the increase in demand and prices of fish (GoM, 2011). As 65% of the people in Malawi are poor,

*Corresponding author. E-mail: erchirwa@gmail.com.

living on less than 1 US \$ per day, many people may not afford to buy fish. This is notwithstanding that compared with other animal products, fish still remains relatively cheap. Consequently, many Malawians may lack sufficient animal protein in their diets, leading to stunting and other protein-deficiency problems (IFFRI, 2012). Further, fish availability has declined, resulting in decreasing per capita fish consumption from 14.7 kg/person/year in 1970 to 4.9 kg/person/year in 2011 (Russell et al., 2008; Nagoli et al., 2009; Sanudi et al., 2015), and is projected to reach zero by 2034 (GoM, 2011) (Figure 1).

As fish is important to people's health and the economy of Malawi, the government has been considering various options of increasing fish supply in the country. Aquaculture is seen as the most viable option (Sanudi et al., 2015). However, the main constraint in Malawi's aquaculture is the slow growing and small-sized native fish species cultured (GoM, 2011, 2012). This problem has been observed since the 1960s, but efforts to identify more suitable indigenous fish species have been unsuccessful. This led the government of Malawi to import the common carp (*Cyprinus carpio* Linnaeus, 1758) (Figure 2) from Israel in 1976.

Test trials for growth showed that the common carp grew fast and to a large marketable size. The government distributed the fish to farmers in southern Malawi from 1985 to 1990. Further importation and distribution of the fish was stopped in 1991 and a ban imposed on its culture in 1992. But farmers wanted a reversal on carp ban to promote growth in the aquaculture sector (GoM, 2011). Details on the history of aquaculture development in Malawi are covered in Pruginin (1976), Balarin (1987) and ICLARM/GTZ (1990). This paper reviews the origin and status of carp farming in Malawi, the issues surrounding the ban of the fish, and research areas to inform policy on common carp farming in Malawi are suggested.

METHODS OF INVESTIGATION

Information on the farming of common carp in Malawi was obtained from the following sources:

- (1) A compilation of existing literature on the origin and status of common carp farming in Malawi. Since farming of this fish in Malawi has been restricted, some of the literature on the subject is admittedly quite old (Betram et al., 1942; Pruginin, 1976; Balarin, 1987; Welcomme, 1988; Vanden Bossche and Bernacsek, 1990; Msiska and Costa-Pierce, 1993).
- (2) Personal involvement as a research assistant in Zomba district in 1989/1990 under the International Center for the Living Aquatic Resources Management¹ (ICLARM/GTZ Africa Project).

Further information was obtained through attendance of meetings organized by ICLARM in 1989/1990 and in later years by the Department of Fisheries².

RESULTS

History of common carp farming in Malawi

The farming of fish in Malawi started in 1906 with the introduction of rainbow trout (*Onchorhynchus mykiss* Walbaum, 1792) for angling (Balarin, 1987). As the human population at this time was low, fish stocks in the capture fisheries were considered adequate and in healthy state. A nutritional survey following the League of Nations (1935), Report on the Psychological Basis of Nutrition recommended that the farming of fish for food in upland areas of Malawi needed consideration to redress the nutritional deficiency in the diet of people living far from lakes (Betram et al., 1942). Thus, fish farming for food began in 1956/1957 using the indigenous tilapias *Oreochromis shiranus* Boulenger, 1897 and *Tilapia rendalli* Boulenger, 1897 (Pruginin, 1976). In 1969, a joint Malawi Fisheries Department (FD)/FAO survey of fish yields of Malawian species was carried out to assess the performance of these fishes. Results indicated slow growth rate (< 1 g/day), leading to low fish yields from ponds and dams of 0.1 to 0.2 t/ha/year for Northern Malawi and 0.5 to 1.0 t/ha/year for Southern Malawi (Msiska, 1993).

Presently, there are five main indigenous fish species used in Malawi's aquaculture, the tilapias *O. shiranus*, *Oreochromis karongae* Trewavas, 1941, *Oreochromis mossambicus* Peters, 1852, *T. rendalli*, and the catfish *Clarias gariepinus* Burchell, 1822 (Figure 3). Tilapias and catfish make up 93 and 5% of aquaculture production, respectively. *Oreochromis shiranus* is the most widely cultured fish in Malawi, followed by *T. rendalli*. *O. mosambicus* is cultured in the Lower Shire river basin. These tilapias grow slowly and to small sizes, with *O. shiranus* and *O. mosambicus* reaching sexual maturity as small as 6 g and breed precociously (M'balaka et al., 2012).

The perceived absence of a fast-growing local species and the need to provide animal protein and farm employment to rural people prompted the government of Malawi, as suggested by a consultant named Pruginin, to decide importing common carp, from Israel in 1976, for aquacultural purpose (Mkoko, 1993). Five hundred common carp of both scaled and mirror carp were imported and acclimated at the Kasinthula Experimental Station (Moreau and Costa-Pierce, 1997). Common carp is native in the piedmont zone of the Danube River to the Black, Caspian and Aral Sea basins, with western dispersants in central Asia and eastern dispersants in Siberia (Kirpichnikov, 1999). However, the fish has been

¹Now WorldFish Center

²The primary author has worked with Malawi's Department of Fisheries for 15 years as a Fisheries Officer




Figure 1. Trends in fish supply, demand and consumption in Malawi (Department of Fisheries, Lilongwe and National Statistical Office, Zomba)



Figure 2. Common carp (Donkers, 2004).




Figure 3. Native fish species used in Malawi's aquaculture (National Aquaculture Center and Atlas of Southern African Fresh water fishes).




Figure 4. Catchment area of Lake Malawi, southern Africa (ILEC, 2005). Carp was distributed to farmers outside the catchment (not shaded).

translocated and introduced to non-native regions since Roman times (Balon, 1995). The fish is regarded as one of the best growing aquaculture species in the world, and has been referred to as a “biological miracle” for its excellent growth and production performance in aquaculture systems (Msiska and Costa-Pierce, 1993). Thus, the common carp was the first to be introduced outside its natural range for aquaculture farming (Balon, 1995; Alves et al., 1999) and remains the most widely distributed fish in the world (ISSG, 2000; Zhou et al., 2003; Casal, 2006). In 2010, the fish accounted for 9% of the total fish production in freshwater aquaculture (Rahman, 2015). Some European countries obtain as much as 80% of their freshwater aquaculture production from common carp alone (Woynarovich et al., 2010; Rahman, 2015).

During test trials from 1976 to 1983 at the Domasi and Kasinthula Experimental Stations in Malawi, the carp performed well in ponds with growth increments averaging 4 g/day, leading to yields of 5 t/ha/year (Msiska, 1993). For example, in the Chingale area of Zomba district of Malawi, common carp was able to grow up to 9 kg in 2.5 years (Moreau and Costa-Pierce, 1997). Average weights for carp were 400 g; *O. shiranus*, 57 g and *T. rendalli*, 78 g (Noble, 1993). Following encouraging performance of carp in test trials, the government considered distributing the fish to farmers.

Before the distribution of carp to farmers, the Fisheries Department of Malawi formulated conditions for the distribution: (1) only farmers outside the Lake Malawi catchment area would be allowed to raise common carp; (2) no farmer would be allowed to breed the fish; (3) all

carp fingerlings were to be supplied from government fisheries stations (Domasi and Kasinthula) at a nominal fee; (4) farm ponds must have screens on inlets and outlets to prevent carp escapes; (5) at harvest all fish must be killed and sold in the presence of a Fisheries Officer; (6) all farmers growing carp must submit records on their carp stocks and possible information on carp transfers to neighbors (Mkoko and Mutambo, 1993).

The distribution, carried out between 1985 and 1990, was largely confined to the southern region, outside the Lake Malawi catchment area (Figure 4), except for a small population of carp stocked in experimental ponds at Bunda College of Agriculture (now LUANAR) for experimental purposes (Moreau and Costa-Pierce, 1997). The fish was distributed to 36 individual farmers in southern Malawi, mainly in Zomba district, and two estates (Satemwa Tea Estate in Mulanje district and SUCOMA Sugar Estate (now Illovo) in the Lower Shire) (Msiska, 1993). Because of its fast growth rate and large market size, common carp was raised mainly for commercial purposes (NRCM, 1999). A fuller utilization of this fish was realized when raised in association with other fish like *T. rendalli*, *O. shiranus* and *O. mossambicus* or *C. gariepinus*. The ease with which this species could be cultured, fast growth under poor input conditions and breeding without slowing down growth, and adaptability to wide environmental conditions made this species attractive to farmers (Kapeleta, 2001). Consumers also liked the fish for its flavor, and the large harvest size made a lot of farmers realize income they never had before (Andrew et al., 2003).

By 1989, two incidences were reported that would

change the course of carp farming in Malawi. Firstly, escapes were reported into Likangala stream in the Lake Chilwa basin and in the Lower Shire system. This implied that escapement of common carp was possible even with the best trained and well-experienced farmers. Secondly, the fish was reported to reproduce in the weedy margins of some of the farmers' ponds, contrary to the belief that the fish would not spawn under natural conditions. The Fisheries Department had never thought that these incidences would occur (Msiska and Costa-Pierce, 1996; Andrew et al., 2003). These two incidences heightened the concerns about the potential impacts of the fish on native fish biodiversity, particularly, in Lake Malawi, the world's most species-rich freshwater lake. Malawi has been particularly concerned about the possible negative impacts of carp on Lake Malawi fish biodiversity (Vanden Bossche and Bernacsek, 1990).

In 1990, the government of Malawi stopped further distribution of common carp fingerlings, pending a decision on the future of carp farming. This, coupled with drought in that year that dried up more than 50% of farmers' ponds, drastically reduced the number of carp farms to just about four or five in the country (Moreau and Costa-Pierce, 1997). In 1991/1992, carp withdrawn from farmers and the remaining stocks in ponds were eradicated in Malawi. All exotic fishes, including carp and even *Oreochromis niloticus*, were prohibited in the Lake Malawi catchment area by the Malawi Government, in order to conserve the lake's unique assemblage of native species (Msiska and Costa Pierce, 1993; Moreau and Costa-Pierce, 1997). The restriction of exotic fish farming became legalized in the 1997 Fisheries Conservation and Management Act [Part XI section 41(1) c] (Hecht and Maluwa, 2003). However, the farming of common carp was still carried out (although negligibly) in some parts of the Lower Shire and other areas in Southern Malawi (Msiska and Costa-Pierce, 1996; Andrew et al., 2003).

Reasons for banning common carp in Malawi

The decision to import common carp did not consider the ecological effects of the fish on aquatic ecosystems (Costa-Pierce et al., 1993). When reports of devastating ecological impacts of the Nile perch introduced into Lake Victoria in the 1950s began to spread in the early 1990s, scientists in Malawi were awakened to the negative effects that introduced fish species can have on native biota. Although, common carp was highly valued and already being distributed to farmers, scientists began to ponder about the potential negative impacts of this species on the unique Lake Malawi fish biodiversity. Fortunately, the fish had already been introduced elsewhere in the world and to more than 21 countries in Africa (Table 1 and Figure 5) from which lessons of its ecological effects could be learned.

Lessons from other countries in Africa suggested that

carp's habit of digging up lake's sediment could destroy tilapia breeding areas, thereby lowering tilapia recruitment due to disruption of nesting. The stirring of sediment by carp also hasten eutrophication by mobilizing sediment-bound nutrients (mostly phosphorus) into the water column (Costa-Pierce and Pullin, 1989; Breukelaar et al., 1994). However, considering the high economic value of carp farming, no country in Africa, beside Malawi, has rejected the fish on account of its ecological effects. The Malawi government was concerned about the potential threat of carp to the unique fish biodiversity of Lake Malawi if it escaped into the lake (Vanden Bossche and Bernacsek, 1990; Costa-Pierce et al., 1993).

Impacts of common carp ban in Malawi

Declining contribution of carp to fish supply and continued slowing of aquaculture growth

The contribution of common carp to aquaculture production declined from about 9% of total aquaculture production in the early 1990s to less than 0.5% by the early 2000s (Figure 6). During the same period, the contribution of common carp to global aquaculture production increased from 5.4% of global aquaculture production to 5.9%.

In Malawi, fish farming became less profitable for most of the farmers who were used to carp, prompting over 80% of them to quit fish farming altogether. Farmers' trust and confidence in the Malawian Fisheries Department declined sharply, setting the government on frantic but futile confidence rebuilding campaigns (Msiska and Costa-Pierce, 1993). Growth in Malawi's aquaculture has slowed. For instance the contribution of aquaculture to total fish supplies in Malawi has remained low, estimated at 2% (Sanudi et al., 2015). It is widely believed that if Malawi were to adopt common carp farming, aquaculture development would accelerate (GoM, 2011).

Search for indigenous aquaculture species

Aquacultural farmers demanded a replacement of the common carp to maintain profitable fish farming in Malawi. The search for suitable native aquaculture species had already proved difficult when such efforts began in the 1960s. However, the scientists believed a lack of success in this direction was attributable to an absence of sustained project commitment to screen and test indigenous fish species (Msiska and Costa-Pierce, 1993). Such a project came along in late 1999 with funding from Japan International Cooperation Agency (JICA). A number of indigenous fish species were assessed. By the end of the 5-year project period (1999-2004), no suitable indigenous fish species was identified

Table 1. Common carp introductions in Africa (Welcomme, 1988; Moreau and Costa-Pierce, 1997; FAO Inland Water Resources and Aquaculture Service, 2003).

Country	Origin	Year	Established?	Ecological effects?
South Africa	Germany	1859	Yes (Some reservoirs)	Yes
Kenya	South Africa	1910	Yes	Probably
Kenya	Uganda	1910	Yes	Probably
Madagascar	Unknown	1914	Yes (Lakes)	Probably
Zimbabwe	South Africa	1925; 1963	Yes (Some reservoirs)	Unknown
Morocco	France	1925	Yes	Yes
Egypt	Indonesia	1934	Yes	Unknown
Ethiopia	Italy	1940	Yes	Unknown
Zambia	Israel	1980	No	Unknown
Zambia	South Africa	1946	No	Unknown
Nigeria	Austria	1954	Probably	Unknown
Rwanda	Israel	1960	Yes	Unknown
Uganda	Israel	1962	Yes	Unknown
Ghana	Unknown	1962	Probably	Unknown
Tunisia	Germany/France	1965	Probably	Probably
C.A.R.	Israel	1966	Yes	Unknown
Cameroon	Israel	1970	Yes	Unknown
Malawi	Israel	1976	No	Improbable
Sudan	India	1975	No	Unknown
Mauritius	India	1976	Yes	Unknown
Cote d'Ivoire	Italy	1976	Yes	Unknown
Burundi	Rwanda	1980-1989	Unknown	Unknown
Algeria	Hungary	1985	Yes (Lakes)	Yes
Togo	Israel	1965; 1971	No	Unknown
Tanzania	India	1981	Unknown	Unknown
Mozambique	South Africa	1988	Yes (Limpopo R)	Unknown
Namibia	South Africa	Unknown	Yes	Unknown
Swaziland	South Africa	Unknown	Unknown	Unknown
Lesotho	South Africa	1965	Yes (Orange R)	Unknown

**Figure 5.** Main producer countries of *Cyprinus carpio* (FAO, 2004-2017).




Figure 6. Impact of carp ban on carp production in Malawi (FAO, 2005-2016; NAC, 2003).

that could replace carp (SSC, 2005).

Pressure on policy review

Failure by government to identify a better-performing native aquaculture species has attracted widespread calls from farmers for a reversal on common carp ban (GoM, 2011). Slow-growing and stunting native fish species have been cited as a major impediment to aquaculture growth in Malawi (GoM, 2011, 2012). The National Aquaculture Strategic Plan (NASP) (2005-2015) calls for impact studies to provide information that would form the basis for policy review on the use of carp in Malawi's aquaculture. The National Biodiversity Strategy and Plan (NBSP) also calls for impact assessment of

alien species as potential candidates for aquaculture development in Malawi (Environmental Affairs Department, 2006). The government has emphasized the need to do more research on the ecological impacts of carp before it can consider reviewing its policy (Msiska and Costa-Pierce, 1993; Bandula, 1997; SSC, 2005).

Research directions related to policy on common carp farming

Government insistence on judicious research on common carp before policy review is understandable and logical. However, government has not suggested any potential policy research direction on the issue. This paper suggests the following possible research avenues:

(1) Comparison of ecological impacts between common carp and *C. gariepinus*: Both of these species are benthivorous (Koekemoer and Steyn, 2002; Rahman, 2015). Anecdotal field observations of the impacts of these fishes in Malawi suggest they may impact on ecosystems in the same way. These observations contradict Msiska and Costa-Pierce (1993) who opined that common carp occupies a niche that no other farmed fish occupies in Malawi. Reports of common carp displacing *C. gariepinus* from its benthic niche in Zimbabwean reservoirs (Costa-Pierce et al., 1993) corroborate observations of similarity of niche occupation by the two species. If these observations are true, the introduction of one of them into a system already containing the other may not significantly alter the ecosystem.

(2) Assessment of colonization and establishment of common carp and its environmental impacts in the Lake Chilwa and Lower Shire drainage systems: Common carp is reported to have escaped from fish ponds into natural waters in these ecosystems in the 1988-1990 period (Msiska and Costa-Pierce, 1993). The escapees have not been followed to determine if they are established and what impacts, if any, they cause.

(3) Evaluation of common carp farming in the Lake Malawi catchment area in Tanzania and Mozambique: These riparian countries are reported to be farming common carp in their side of the lake's catchment (Costa-Pierce et al., 1993; Chirindza, 2010). Mozambique is one of the main African producer countries of common carp (Figure 5). If the species is already in the lake's watershed, the questions of what impacts the common carp is causing and what justification Malawi has for its unilateral rejection of the fish when other countries in the same watershed are farming it will need addressing.

(4) Invasion history of common carp in lakes of similar morphometry and physico-chemical conditions to Lake Malawi: Across-ecoregion analysis has shown that the invasivity of common carp is regulated by a number of ecological filters such as depth and trophic status of a water body (Bajer et al., 2015). However, no studies have been conducted to establish invasion history of common carp farmed in catchment areas of lakes with depth and trophic status similar to Lake Malawi. Lessons learned in these ecosystems can be used to make inference about the potential vulnerability or invasion potential of Lake Malawi to common carp.

Conclusions

Common carp was introduced in Malawi to complement tilapia aquaculture production with an aim of increasing overall production from the fish farming industry. The

indigenous tilapias were slow-growing, stunting and breeding precociously. The catfish *C. gariepinus* was economically a difficult fish for the majority of farmers as the species' protein requirements made it costly to feed. The species was also difficult to breed under prevailing pond conditions. Thus, these species had received farmers' disapproval as early as the 1960s. With common carp (1985-1991), the farmers' interest in fish farming surged as profits from fish farming began to increase. Government's withdrawal of common carp and its ultimate ban left many farmers disillusioned and wondering what the real justification was. The government insisted it was concerned with the effects the fish would have on Lake Malawi once the species found its way to the lake's catchment. Farmers were promised that a more suitable indigenous aquaculture species would be identified for use in Malawi's aquaculture sector.

To date, a more suitable native aquaculture species has not been identified in Malawi, despite the existence of well-resourced project investments in this effort. Lack of better performing indigenous fish species is continually being cited as a major constraint in the growth of aquaculture industry in Malawi. The Malawi government has persistently resisted calls to reverse its ban of common carp until it could be shown that carp's farming in the Lake Malawi catchment would not negatively affect the lake's unique fish biodiversity. This paper has outlined potential research areas that are relevant to policy decision-makers on the question of whether common carp would harm Lake Malawi.

Due to inconsistent and fragmentary documentation of the farmers involved in carp farming, this study has not been able to chronicle the trends of carp farmers in Malawi from the time of distribution of the fish to farmers to the time the fish was withdrawn from the farmers. Although it is recorded that carp was still being farmed after banning it (Andrew et al., 2003), the farmers could not make public declaration of the activity for fear of government reprisals. The study has therefore not been able to provide up-to-date records of carp production in Malawi. In addition, there are unconfirmed reports that farmer-to-farmer distribution of carp has occurred in Malawi, and that the fish is illicitly farmed in the Lake Malawi catchment area. This study has not been able to verify these reports as the farmers fear to provide information.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGEMENTS

Support for this research was made possible through a

capacity building competitive grant “Training the next generation of scientists” provided by Carnegie Cooperation of New York through the Regional Universities Forum for Capacity Building in Agriculture (RUFORUM). Technical guidance throughout the study was offered by the Graduate Teaching Assistantship (GTA) host Lilongwe University of Agriculture and Natural Resources (LUANR) through its Department of Aquaculture and Fisheries Science of the Faculty of Natural Resources. Martin Potgieter and another unknown reviewer provided invaluable comments that were used to improve an earlier manuscript.

REFERENCES

- Alves CBM, Vono V, Vieira F (1999). Presence of the walking catfish *Clarias gariepinus* (Burchell) (Siluriformes, Clariidae) in Minas Gerais state hydrographic basins, Brazil. *Rev. Bras. Zool.* 16(1):259-263.
- Andrew TG, Weyl OLF, Andrew M (2003). Aquaculture master plan development in Malawi: Socioeconomic survey report.
- Bajer PG, Cross TK, Lechelt JD, Chizinski CJ, Weber MJ, Sorensen PW (2015). Across-ecoregion analysis suggests a hierarchy of ecological filters that regulate recruitment of a globally invasive fish. *Divers. Distrib.* 21(5):500-510.
- Balarin JD (1987). National reviews for Aquaculture Development in Africa. Malawi. FAO Fisheries Circular, No. 770.12. 83 p.
- Balon EK (1995). The common carp, *Cyprinus carpio*: its wild origin, domestication in aquaculture, and selection as colour nishikigoi. *Guelph Ichthyol. Rev.* 3:1-55.
- Bandula DD (1997). Status of the common carp, (*Cyprinus carpio*) as a candidate exotic fish for fish farming in Malawi: Report of the Technical Consultation on Species for Small Reservoir Fisheries and Aquaculture in Southern Africa. Livingstone, Zambia, 7-11 November 1994, FAO, Harare (Zimbabwe), 1997, pp. 33-34, ALCOM report. Harare [Alcom Rep.], no. 19.
- Betram CK, Borley HJ, Trewavas E (1942). Report on fish and fisheries of Lake Nyasa. Crown Agents for the Colonies on behalf of the Government of Nyasaland, London, England. 91 p.
- Brekelaar AW, Lammens EHRR, Klein Breteler JGP, Tatrai I (1994). Effects of benthivorous bream (*Abramis brama*) and carp (*Cyprinus carpio*) on sediment resuspensions and concentrations of nutrients and chlorophyll *a*. *Freshwater Biol.* 32:113-121.
- Casal CMV (2006). Global documentation of fish introductions: the growing crisis and recommendations for action. *Biol. Inv.* 8(1):3-11.
- Chidammodzi C, Muhandiki V (2015). Development of indicators for assessment of Lake Malawi Basin in an Integrated Lake Basin Management (ILBM) framework. *Int. J. Commons* 9(1).
- Chirindza IA (2010). A survey of small-scale rural aquaculture in Mozambique. United Nations University Fisheries training Program, Iceland [final project]. <http://www.unuftp.is/static/fellows/document/isac09prf.pdf>
- Costa-Pierce BA, Moreau J, Pullin RSV (1993). New Introductions of Common Carp (*Cyprinus carpio* L) and Impacts on Indigenous Species with reference to Sub-Saharan Africa. ICLARM Contribution No. 744.
- Costa-Pierce BA, Pullin RSV (1989). Stirring ponds as a possible means of increasing aquaculture production. *Aquabyte* 2:5-7.
- Environmental Affairs Department (2006). National Biodiversity Strategy and Action Plan. Lilongwe, Malawi.
- Food and Agriculture Organization (FAO) (2005-2016). National Aquaculture Sector Overview. Malawi. National Aquaculture Sector Overview Fact Sheets. Text by Chimatiro SK and Chirwa BB. In: FAO Fisheries and Aquaculture Department [online]. Rome. Updated 1 January 2005. [Cited 27 August 2016]. http://www.fao.org/fishery/countrysector/naso_malawi/en
- Food and Agriculture Organization (FAO) (2004-2017). Cultured Aquatic Species Information Program. *Cyprinus carpio*. Cultured Aquatic Species Information Program. Text by Peteri A. In: FAO Fisheries and Aquaculture Department [online]. Rome. Updated 1 January 2004. [Cited 9 May 2017]. http://www.fao.org/fishery/culturedspecies/Cyprinus_carpio/en
- Government of Malawi (GoM) (2012). National Fisheries Policy 2012-2017. 2nd edn. Department of Fisheries, Lilongwe.
- Government of Malawi (GoM) (2011). Annual Economic Report, Malawi: Ministry of Development Planning and Cooperation, Lilongwe.
- Hecht T, Maluwa A (2003). Situation analysis of aquaculture in Malawi. Envirofish Africa (Pty) Ltd, Grahamstown, South Africa, and National Aquaculture Centre, Domasi, Malawi.
- ICLARM/GTZ (1990). The context of small-scale integrated agriculture-aquaculture systems in Africa: a case study of Malawi. *ICLARM Stud. Rev.*, No.18. 302 p.
- ILEC (International Lake Environment Committee) (2005). Managing Lakes and their Basins for Sustainable Use: A report for Lake Basin managers and stakeholders. International Lake Environment Committee Foundation, Kusatsu, Japan.
- ISSG (Invasive Species Specialist Group) (2000). IUCN guideline for the prevention of biodiversity loss caused by alien invasive species. IUCN, Gland.
- Kapeleta MV (2001). Aquaculture development in Malawi: Tilapia culture. In Promotion of sustainable commercial aquaculture in Sub-Saharan Africa. Experiences of selected developing countries, FAO Fisheries circular No *Cyprinus carpio* 971 FIEP/C971 (Bi).
- Kirpichnikov VS (1999). Genetics and breeding of common carp. Paris: INRA, p97.
- Koekemoer JH, Steyn GJ (2002). Fish community study of Hartbeespoort dam. Final Report. Department of Agriculture, Conservation, Environment and Tourism. R.S.A.
- Mahony C, Clark H, Bolwel M, Simcock T, Potter R, Meng J (2014). Where Politics Borders Law: The Malawi-Tanzania Boundary Dispute. New Zealand Centre for Human Rights Law, Policy and Practice – Working Paper 21, February 2014.
- M'balaka M, Kassam D, Rusuwa B (2012). The effect of stocking density on the growth and survival of improved and unimproved strains of *Oreochromis shiranus*. *Egypt. J. Aquat. Res.* 38:205-211.
- Mkoko BJ, Mutambo JT (1993). Formulation and Status of Government Policy on Common Carp. In: Msiska, O.V. & B.A. Costa-Pierce (Eds) History, Status and Future of Common Carp (*Cyprinus carpio* L.) as an exotic species in Malawi. WorldFish Center—ICLARM, Manila.
- Moreau J, Cost-Pierce B (1997). Introduction and present status of exotic carp in Africa. *Aquac. Res.* 28:717-732.
- Msiska OV, Costa-Pierce BA (Editors) (1993). History, Status and Future of Common Carp (*Cyprinus carpio* L.) as an exotic species in Malawi. WorldFish Center—ICLARM, Manila.
- Msiska OV (1993). Status and Performance of Common Carp on Estates in Malawi. In: Msiska, O.V. & B.A. Costa-Pierce (Eds) History, Status and Future of Common Carp (*Cyprinus carpio* L.) as an exotic species in Malawi. WorldFish Center—ICLARM, Manila.
- NAC (2003). National Aquaculture Center Annual Report, 2002/2003.
- Nagoli J, Phiri EM, Kambewa E and Jamu D (2009). Adapting Agriculture Aquaculture for HIV and AIDS – Affected Households: The case of Malawi. The WorldFish Center Working Paper 1957. The WorldFish center, Penang, Malaysia.
- Noble RP (1993). Status and Performance of Common Carp on smallholder Farms in Zomba District, Malawi. In: Msiska, O.V. & B.A. Costa-Pierce (Eds) History, Status and Future of Common Carp (*Cyprinus carpio* L.) as an exotic species in Malawi. WorldFish Center—ICLARM, Manila.
- NRCM (National Research Council of Malawi) (1999). Malawi agricultural sciences and natural resources research master plan. NRCM Agricultural Sciences Committee. 167 p.
- Pruginin Y (1976). Fish farming in Malawi. FI: DP/MLW/71/516/8. FAO. Rome.
- Rahman MM (2015). Role of common carp (*Cyprinus carpio*) in aquaculture production systems. *Front Life Sci.* 8(4):399-410.
- Russell AJM, Grötz PA, Kriesemer SK, Pems DE (2008). Recommendation Domains for Pond Aquaculture. Country Case Study: Development and Status of Freshwater Aquaculture in Malawi. WorldFish Center Studies and Reviews No. 1869. The WorldFish Center, Penang, Malaysia. 52 p.

- Sanudi F, Jere B, Mzengereza K, Chirwa BB (2015). Effect of Stocking Density and Feed on Growth of Improved (F5) Mono-Sex *Oreochromis shiranus* Reared in Tanks. *J. Fisheries Livest. Prod.* 3(4):1-5.
- SSC (System Science Consultants Inc) (2005). National Aquaculture Strategic Plan (NASP) (2006–2015). Master Plan Study on Aquaculture Development in Malawi. Main Report. Japan International Cooperation Agency, Tokyo.
- Vanden Bossche J-P, Bernacsek GM (1990). Source book for the inland fishery resources of Africa: 1. CIFA Technical Paper No. 18.1. Rome, FAO. 240 p.
- Welcomme RL (1988). International Introductions of Inland Aquatic Species. FAO, U.N.
- Weyl OLF, Ribbink AJ, Tweddle D (2010). Lake Malawi: fishes, fisheries, biodiversity, health and habitat. *Aquat. Ecosyst. Health Manag.* 13(3):241-254.
- Woynarovich A, Moth-Poulsen T, Peteri A (2010). Carp polyculture in Central and Eastern Europe, the Caucasus and Central Asia: a manual. Rome: FAO.
- Zhou JF, Wu QJ, Ye YZ, Tong JG (2003). Genetic divergence between *Cyprinus carpio carpio* and *Cyprinus carpio haematopterus* as assessed by mitochondrial DNA analysis, with emphasis on origin of European domestic carp. *Genetica* 119:93-97.