

Full Length Research paper

## Biochemical and histopathological changes in liver of albino rats fed diets incorporated with *Vernonia amygdalina* and *Vernonia colorata* leaves

Ijeh, Ifeoma Irene<sup>1\*</sup> and Obidoa Onyechi<sup>2</sup>

<sup>1</sup>Department of Biochemistry, College of Natural and Applied Sciences, Michael Okpara University of Agriculture, Umudike, P. M. B. 7267, Umuahia, Abia State, Nigeria.

<sup>2</sup>Department of Biochemistry, Faculty of Biological Sciences, University of Nigeria, Nsukka, Nsukka, Enugu State, Nigeria.

Accepted 21 September, 2010

*Vernonia amygdalina* and *Vernonia colorata* are widely used in medicinal plant preparations in sub-Saharan Africa. Certain dietary constituents can potentiate or reduce the potency of toxins such as AFB<sub>1</sub>. The effect of dietary incorporation of *V. amygdalina* and *V. colorata* on some biochemical and histopathological indices in albino rats was studied. *V. amygdalina* and *V. colorata* were incorporated into diets compounded to isocaloric and isonitrogenous and fed to mature male albino rats administered 250 mg/kg Aflatoxin B<sub>1</sub> over a 12 week period. Bioactivity studies using brine shrimp lethality test indicate that methanolic extracts of *V. amygdalina* and *V. colorata* had higher bioactivity (ED<sub>50</sub> 121.47 µg/ml and 761.8 µg/ml respectively) than their aqueous extracts (819.5 and 1,999.85 µg/ml respectively). Alanine amino transaminase (ALT) activity increased significantly ( $P \leq 0.05$ ) relative to control group (Gp1) in groups fed *V. colorata* with or without AFB<sub>1</sub> treatment but showed no significant increase in groups fed *V. amygdalina* with or without AFB<sub>1</sub>. Treatment of *V. colorata* fed animals with AFB<sub>1</sub> (Gp3) resulted in marked increase in alkaline phosphatase activity. Increase in Aspartate amino transaminase (AST) activities were not significantly different from those of animals fed only with *V. colorata* without AFB<sub>1</sub>. Administration of AFB<sub>1</sub> resulted in a significant increase in alkaline phosphatase activity in groups fed *V. colorata*, but not in groups fed *V. amygdalina*. Groups fed *V. amygdalina* + AFB<sub>1</sub> showed no significant changes in ALT and AST. These findings suggest that while the feeding of *V. amygdalina* may have hepatoprotective effects, the feeding of *V. colorata* may potentiate the toxic effects of toxins such as AFB<sub>1</sub>. Histopathological studies on the liver show that the feeding of the two vegetable varieties affect the liver in different ways.

**Key words:** *Vernonia amygdalina*, *Vernonia colorata*, AFB<sub>1</sub>, serum enzymes, bioactivity.

### INTRODUCTION

*Vernonia amygdalina* is widely used in the preparation of soups and porridges in south Eastern Nigeria. Two major varieties are used domestically. These two vary in their degree of bitterness (*Vernonia colorata* which is less bitter) and (*V. amygdalina* del which is very bitter). *V. amygdalina* del is usually macerated extensively before

use in soups and porridges but often used in unprocessed form in medicinal preparation. Its extract is used as a digestive tonic, appetizer, febrifuge and as antidiabetic tonic (Singha, 1966).

*V. amygdalina* var, is used in unprocessed forms as snack or in porridges. A number of the above uses have been investigated viz; antidiabetic activity (Akah and Okafor, 1992), antimalarial (Philipson et al., 1993) antiparasitic activity (Ohigashi, 1995) and antimicrobial activity against organisms commonly incriminated in wounds (Ijeh et al., 1996).

\*Corresponding author. E-mail: [ijehirene@yahoo.com](mailto:ijehirene@yahoo.com), [ijeh.ifeoma@mouau.edu](mailto:ijeh.ifeoma@mouau.edu).

**Table 1.** Composition of diets for different groups of rats (g%).

Ingredient/feedstuff	Diet 1	Diet 11	Diet 111
Maize (flour)	50.50	50.50	50.50
Crayfish	4.20	3.36	3.36
Groundnut (meal)	9.29	7.43	7.43
<i>V. colorata</i>	-	4.61	-
<i>V. amygdalina. del</i>	-	-	4.33
Bone meal	4.00	4.00	4.00
Pre-mix	0.50	0.50	0.50
Oil	5.00	5.00	5.00
Corn-flour	26.50	24.50	24.88
Total (approx)	100.00	100.00	100.00

**Table 2.** Brine shrimp Lethality tests.

Extract	ED (50 µg/ml)
Aqueous extract <i>V.a</i> var	1,999.85
Methanolic extract <i>V.a</i> var	761.89
Aqueous extract <i>V.a</i> del	819.59
Methanolic extract <i>V.a</i> del	121.47

Terpenoid extracts from *V. amygdalina* have been demonstrated to have hepatoprotective effects against Carbon tetrachloride induced liver damage in rats. Arhoghro et al. (2009) also reported that aqueous extracts of *V. amygdalina del* had hepatoprotective effects against Carbon tetrachloride induces liver damage in albino rats.

Liver diseases especially liver cancer is major cause of mortality and morbidity in the tropics. Hepatocellular carcinoma is the third leading cause of cancer deaths world wide with prevalence 16 to 32 times higher in developing countries where storage facilities are poor and risk of aspergillus flavus infestation of stored food products relatively higher than developed countries and temperate regions of the world (Liu and Wu, 2010). Numerous epidemiological studies have observed a correlation between areas of high aflatoxin B<sub>1</sub> exposure and high incidence of hepatocellular carcinoma (Jackson and Groopman, 1999).

The feeding of AFB<sub>1</sub> along side *V. amygdalina* in this study is borne out of the need to investigate its possible effects on AFB<sub>1</sub> toxicity since research evidence has shown that certain dietary constituents can potentiate or reduce the potency of toxins such as AFB<sub>1</sub>. AFB<sub>1</sub> is a major cause of hepatocarcinogenicity in Africa and Asia (Liu and Wu, 2010). Antioxidant activities of flavonoids from *V. amygdalina del* have been previously reported (Igile et al., 1994). The two varieties of Vernonia most commonly consumed in the South east of Nigeria are often confused by inexperienced users. Several steroidal saponins and proteins found in *V. amygdalina* have been

shown to have anticancer effect (Gresham et al., 2008).

## MATERIALS AND METHODS

### Plant materials

Stem cuttings of leaves of *V. amygdalina* and *V. colorata* were obtained from Okigwe Imp State, Nigeria and were botanically identified at the Botany Department of the University of Nigeria Nsukka. They were dried in mild sun for three days and then milled into fine particles using a Moulinex blender. They were stored in water proof bags in a deep freezer until ready for use.

### Acute toxicity studies

Brine shrimp lethality tests were carried out using aqueous and methanolic extracts of the leaves of *V. amygdalina. del* and *V. colorata* as described by Maclaughin et al. (1991).

### Experimental design and treatment of animals

Thirty male rats aged five to six weeks weighing 40 to 50 g were randomly assigned to six groups of five animals each. The animals were exposed to 12 h light and dark cycles under humid tropical conditions in stainless steel cages. Feed and water were supplied *ad libitum*.

After 7 days habituation on the basal diet, the animals were assigned randomly to separate cages and introduced to basal diet and diets compounded with the two different varieties under study. The composition of the diets is shown on Table 1. Diets were compounded to be isocaloric and isonitrogenous. Nutrients were supplied in amounts adequate for normal growth. Locally sourced ingredients were used. 250 µg/kg body weight AFB<sub>1</sub> was administered *per os* by intubation for ten days to groups, 2, 4, and 6 commencing on the 10<sup>th</sup> day of feeding the animals (post-equilibration). After AFB<sub>1</sub> dosing all the animals were fed for another nine weeks to complete 12 weeks of feeding post-equilibration. Animals were sacrificed by anaesthetizing with diethyl ether and venous blood was drawn by cardiac puncture and organs were blotted on filter paper and weighed. They were fixed in formal saline for histopathological studies.

### Enzyme assays


Assays of Alanine amino transaminase activity (ALT, EC, 2.6.1.2) and Aspartate amino transaminase activity (AST, EC. 2.6.1.1) were carried out as described of King (1978). Assay of Alkaline phosphatase activity was done as described by Teitz (1987).

### Statistical analysis


Data obtained were analyzed using SPSS. The analysis of variance procedure for completely randomized design was used. Treatment means were separated. Multiple range tests were carried out using Scheffe's tests to separate statistically significant means at a significance level of 0.05

## RESULTS

Table 2 shows result of bioactivity test. Methanolic were observed as shown on Plates 1 to 4. Alkaline extracts of



**Plate 1.** Photomicrograph of liver sections of rats fed the basal diet showing normal lobular architecture [H and E stain X 180] in rats fed basal diet.



**Plate 2.** Photomicrograph of liver sections showing hepatocyte degeneration and necrosis with mononuclear cells infiltration of periportal areas [H and E stain X 180] in rats fed Basal diet + AFB<sub>1</sub>.

*V. amygdalina* del showed the highest bioactivity (ED<sub>50</sub> 121.47 µg/ml) while aqueous extracts of *Vernonia colorata* showed the lowest bioactivity (ED<sub>50</sub> 1, 999.8 µg/ml). Remarkable changes in liver pathology phosphatase activity (Table 3) increased significantly in all groups treated with AFB<sub>1</sub> and also in the group fed *V. amygdalina* del only. ALT activity (Table 3) increased significantly in group fed *V. colorata* but not *V. amygdalina* del.

There was a non-significant ( $p < 0.05$ ) increase in AST activity (Table 4) in all the vegetable diet groups with or without AFB<sub>1</sub> administration.

## DISCUSSION


Results from bioactivity studies indicate that the leaves of *V. amygdalina* del may have higher bioactivity in low doses than *V. colorata*. It also indicates that the bioactivity and pharmacological properties of these leaves may reside more in the methanolic fraction.

Alkaline phosphatase is a cholestasis marker being a membrane bound enzyme. Elevation in its activity is associated with hepatobiliary diseases and is more marked the more complete the obstruction (Teitz, 1987). The variance in the pattern of elevation of alkaline

**Table 3.** Effects of feeding of two varieties of *Vernonia* on some serum enzymes.

S/N	Group treatment	ALT (I.U/L)	AST (I.U/L)	ALK (I.U/L)
1.	Basal diet	12.00 ± 0.00 <sup>a</sup>	12.67 ± 3.00 <sup>a</sup>	34.12 ± 3.75 <sup>a</sup>
2.	Basal diet + AFB <sub>1</sub>	12.00 ± 0.00 <sup>a</sup>	6.67 ± 3.06 <sup>a</sup>	160 ± 17.85 <sup>a</sup>
3.	<i>Vernonia colorata</i> diet	36.00 ± 0.00 <sup>c</sup>	26.66 ± 11.55 <sup>a</sup>	27.22 ± 4.10 <sup>a</sup>
4.	<i>Vernonia amygdalina</i> del diet	8.0 ± 3.46 <sup>a</sup>	14.00 ± 6.93 <sup>a</sup>	140.15 ± 1.78 <sup>b</sup>
5.	<i>Vernonia colorata</i> diet + AFB <sub>1</sub>	37.67 ± 0.29 <sup>c</sup>	28.00 ± 19.29 <sup>a</sup>	421 ± 195.91 <sup>c</sup>
6.	<i>Vernonia amygdalina</i> del diet = AFB <sub>1</sub>	5.00 ± 1.00 <sup>a</sup>	30.00 ± 16.00 <sup>a</sup>	168 ± 76.32 <sup>a</sup>

\*Means with the same superscripts are not significantly different.




**Plate 3.** Liver section showing hyperplasia of the duct epithelium. [H and E stain X 180] in rats fed *Vernonia colorata*.

phosphatase activity in this study with the feeding of the indicative of differences in patterns of toxic lesion the liver associated with the two different cultivars. This inference is further substantiated by the histopathological studies which indicate milder lesions in periportal area of the liver in the animals fed *V. amygdalina* del with AFB<sub>1</sub> treatment. Severe hepatocellular lesions with parenchymal cell necrosis is usually accompanied by a marked elevation of the mitochondrial enzyme AST which is distinctly increased over ALT. This pattern is seen in the group fed *V. colorata* with or without AFB<sub>1</sub> (Plates 3 and 4). These findings are in consonance with the findings of Arhoghro et al. (2009) who also reported hepatoprotective effects of *V. amygdalina* against Carbon tetrachloride induced hepatic damage. Ijeh and Obidoa (2004) reported the induction of Phase 11 biotransformation enzymes such as glutathions S- tranferase activity in rats fed *V. amygdalina* incorporated diets along with AFB<sub>1</sub>. GST is known to biotransform AFB<sub>1</sub> to less toxic metabolites

thereby conferring hepato-protection.

ALT and AST activities in rats fed the *V. amygdalina* without AFB<sub>1</sub> did not differ significantly from rats fed basal diet indicating that the vegetable diet had no chronic toxicity effects. The elevation over ALT is marked in group 5 compared to group 6 indicating a possibility that this vegetable potentiates the toxic effects of AFB<sub>1</sub>. There is a variance in the pattern of increase in AST activities in the groups fed *V. colorata* only and *V. amygdalina* del only without AFB<sub>1</sub> treatment. AST is a cellular enzyme of the liver lobules and changes in its activity vary in serum depending on the severity, type and stage of liver damage (Rosalki and Wilkriion, 1976).

ALT and AST activities in group 6 fed the *V. amygdalina* del compares with those fed the basal diet indicating that the feeding of this vegetable variety may not be toxic to the liver cell itself but could potentiate the toxic effects of AFB<sub>1</sub> possibly by inducing phase one microsomal enzymes. This has been observed by earlier



**Plate 4.** Photomicrograph of liver sections showing proliferation of bile ductules [H and E stain X 180] in rats fed *Vernonia colorata* diets +AFB<sub>1</sub>.

workers (Ezekwe, 1991).

## Conclusion

Our studies are indicative that the two varieties of *Vernonia* under study may have different pharmacological properties and effects on body organs especially the heart and liver. Our findings are indicative that while *V. amygdalina* del may be hepatoprotective against certain forms of liver damage it may cause obstructive liver damage as indicated by elevation of alkaline phosphates activity. *V. colorata* may potentiate the toxic effects of toxins such as AFB<sub>1</sub> as indicated by elevation of alanine amino transaminase activity and aspartate amino transaminase activity.

## ACKNOWLEDGEMENTS

The work was carried out in Prof. O. Obidoa's laboratory in the Department of Biochemistry, University of Nigeria, Nsukka. We acknowledge financial assistance given to Dr Ifeoma Irene Ijeh by the Abia State University Uturu, Nigeria.

## REFERENCES

- Akah PC, Okafor CC (1992). Blood sugar lowering effect of *Vernonia amygdalina*. Del in an experimental rabbit model. *Phytother. Res.*, 6: 171-173.
- Arhoghro EM, Ekpo KE, Anosile EO, Ibeh GO (2009) Effect of aqueous extracts of bitter leaf *Vernonia amygdalina*. Del on Carbon Tetra chloride induced liver damage in Wistar albino rats. *Eur. J. Sci. Res.*, 26(1): 122-130.
- Babalola OO, Anetor JI, Adeniyi FA (2001). Amelioration of carbon tetrachloride-induced hepatotoxicity by terpenoid extract from leaves of *Vernonia amygdalina*. *Afr. J. Med. Sci.*, 30(1-2): 91-93.
- Ezekwe CI (1991). Some Biochemical effects of the vegetables *Vernonia amygdalina*, *Gnetum africanum* and *Telfaria occidentalis* on rat liver microsomes. M. sc thesis submitted to the department of Biochemistry, University of Nigeria Nsukka.

- Gresham JL, Ross J, Izevbogie EB (2008) *Vernonia amygdalina*: Anticancer activity authentication and adulteration, *Int. J. Environ. Res. Pub. Health*, 5(5): 342-348
- Igile AO, Oleszek W, Jurzysta M, Burda S, Fafunso M, Fasamide A A (1994). Flavonoids from *Vernonia amygdalina* and their antioxidant activity. *J. Agric. Fd. Chem.*, 42: 2445-2448.
- Ijeh II, Nwaugo VO, Obidoa O (1996). Comparative studies on the Nutritive, phytochemical and antimicrobial studies on 2 varieties of *Vernonia amygdalina*. *Plant Prod. Res. Commun.*, 1: 77-75.
- Ijeh II, Obidoa O (2004) Effect of Dietary incorporation of *Vernonia amygdalina*. del on AFB<sub>1</sub> Induced hepatotoxicity in Weanling Albino rats. *Jamaican J. Sci. Tech.*, 15: 32-36.
- Jackson EP, Groopman JD (1999) Aflatoxin and Liver cancer. *Best Practice and Res. Clin. Gastroenterology*. 13(4): 545-555.
- King J (1978). Aspartate amino transferase (EC 2.6.1.1) and alanine amino transferase (EC 2.6.1.2). In *Clinical Biochemistry, Principles and methods* (H. C. Ch. Curtis and Macroth Eds), 11: 1148-1151.
- Liu Y, Wu F (2010). Global burden of Aflatoxin- induced hepatocellular Carcinoma -A risk assessment. *Health Perspectives* 118(6):818-824.
- Loliger J (1991). The use of antioxidants in foods. In : *Free radicals and Additives*. Aruoma, O.I., Halliwell, B. Eds. Taylor and Francis, London, p. 121.
- Maclaughin LJ, Chang C, Smith DL (1991). Bench top bioassays for the discovery of bioactive natural products an update. Atta Rahman (eds). *Studies on natural products chemistry*. 3. Elsevier science publishers. Amsterdam, pp. 383-409.
- Ohigashi H, Jisaka M, Takagaki T, Nozaki H, Tada T, Huffman M A, Nishida T, Koji M, Koshimizu K (1991). Bitter principle and related steroid glucoside from *Vernonia amygdalina* a possible medicinal plant for wild chimpanzees. *Agric. Biol. Chem.*, 55: 1201-1203.
- Philipson JO, Wright CW, Kirby GC, Warhurst DC (1992). Phytochemistry of some plants used in traditional medicine for the treatment of protozoal diseases. Abstracts, International symposium of phytochemical society of Europe, University of Lausanne, Switzerland, p. 12.
- Rosalki SB, Wilkron J H (1976). The principles and practice of diagnostic enzymology. Arnold London, pp. 664-667.
- Singha SC (1966), *Medicinal plants in Nigeria*. National press Ltd. Apapa. Nigeria, p. 46.
- Teitz NW (1987). *Fundamentals of clinical chemistry*, W.B. Saunders Co. Thorne CNR (1984). Techniques for determining protein concentration. In *Techniques for life sciences*. Vol. B1/1, B/104/8 Elsevier North Holland, pp. 383-385.