

Full Length Research Paper

Breeding of parental and tolerant hybrids of *Theobroma cacao* L. to *Phytophthora megakarya* Bras. and Griff.

Martine Louise Ondobo¹, Pierre Onomo Effa^{1,2*}, Jude Ndjaga Manga^{1,3}, Jules Christian Kouam Djoko^{1,3} and Pierre François Djocgoue^{1,3}

¹Laboratory of Plant Physiology, Department of Biological Sciences, Higher Teacher's Training College, P. O. Box 47, Yaounde, Cameroon.

²Department of Biochemistry, Faculty of Science, P. O. Box 812, Yaounde, Cameroon.

³Department of Plant Biology, Faculty of Science, P. O. Box 812, Yaounde, Cameroon.

Received 26 February, 2017; Accepted 26 April, 2017

The development of resistant varieties can be done according to a genetic approach. Estimation of genetic parameters for the quantitative traits of cocoa genotypes (*Theobroma cacao* L.) was made from a 5 × 5 diallel mating design. The objective of this study was to identify tolerant genotypes to black pod disease (BPD) through genetic analyzes. The inoculation of the leaves of parental cocoa genotypes and their offsprings with *Phytophthora megakarya* was performed in the nursery for two seasons (dry and rainy). Percentage success for crosses made using hand pollination was low (31.22%) and fair for grafting (60.68%). Observations made on necrosis length on the 2nd, 4th and 6th days after inoculation showed increasing sensibility of the clones to BPD in the order SNK 413<T 79/467<T 79/501<SNK 16<SCA 12. 84.37 and 76.04% of hybrid genotypes exhibited positive heterosis (hybrid vigor) in dry and rainy seasons respectively. Narrow sense (h^2) and broad-sense (H^2) heritabilities was high in two reciprocal crosses [F30 ($h^2= 0.699$ and $H^2=0.624$) and F70 ($h^2= 0.601$ and $H^2=0.643$)].

Key words: *Theobroma cacao*, *Phytophthora megakarya*, tolerant, heterosis, heritability.

INTRODUCTION

Cocoa is native to tropical America and is a cash crop in many tropical countries. It is an important source of foreign exchange, but several factors account for its low productivity. Some of these factors include aging plantations (Tijani, 2005), poor farming practices, presence of insects, presence of rodents and disease infestations especially black pod disease (BPD) (Ndoumbe-Nkeng, 2002). In Cameroon, approximately 80% of its production losses caused by BPD in areas

where climatic conditions are favorable (Despréaux et al., 1988).

Presently, chemical control can be used to manage the disease, but the costs of pesticides is usually too expensive for African farmers. To ensure sustainable production, improvement programs are increasingly focused on developing hybrids that are tolerant to BPD but information on its genetic control has not been fully elucidated.

*Corresponding author. E-mail: peffafr@yahoo.fr. Tel: (+237) 677 405 949.

The selection of cocoa genotypes with reduced susceptibility to black pod disease therefore remains a priority. Despite, many researches on cocoa (Tarjot, 1969; Blaha and Lotodé, 1976), selection for BPD resistant genotypes still remains futile. Many authors suggest the reaction of different *Phytophthora* spp. stem from partial resistance, probably polygenic (Partiot, 1975; Blaha and Lotodé, 1977). Different plant material evaluation methods have been proven (Blaha, 1974). This hypothesis was eventually disproved when experimental evidences showed that the interaction between the host followed the principles of gene-for-gene concept (Brading et al., 2002; Kema et al., 2002). Given that different types of gene actions are important in different crosses, the breeding strategy for developing a desirable genotype should be based on the gene action involved in that particular cross (Manga et al., 2016).

The hybridization technique for cocoa based on vigor, precocity, productivity and disease resistance, has helped to obtain improved genotypes (Eskes and Lanaud, 1997; Djocgoue et al., 2006). This hybridization is usually accompanied a heterotic effect. Heterosis is characterized by faster growth, better vegetative and generative development, better performance and improved tolerance to adverse environmental conditions. Hybrids are characterized by high productivity and environmental adaptability is mainly due to the non-additive gene effect of the parental genotypes (Tahi et al., 2000; Djocgoue et al., 2006; Manga et al., 2016). For this reason, the use of heritability is an effective genetic parameter for selection of quantitative traits in a determined reproductive system. Nyasse et al. (1995) and Djocgoue et al. (2011) reported that additive gene effects were important for transmission of character length of necrosis (tolerance gene at *P. megakarya*). Effa et al. (2015) and Manga et al. (2016) showed that selection based on family performance or progeny test should be more effective. Heritability tends to be only moderate (Simon et al., 1998), but progress in breeding for resistance may still be possible.

Studies on the analysis of data collected from an experiment involving diallel mating design showed that resistance to BPD is an heritable trait, and it is controlled by polygenes and additive gene effect (Djocgoue et al., 2006). Therefore it would be possible to develop cocoa genotypes that would possess genes that are tolerant to BPD. The objective of the present work was to identify hybrids of *T. cacao* that are tolerant to BPD.

MATERIALS AND METHODS

Cocoa plant and *P. megakarya* Oomycete material

This study was carried out on the experimental farms of SODECAO (Cameroon Cocoa Development Corporation) station in Mengang. Five parental clones of *T. cacao* and their progenies used in this study are listed in Table 1. These parents were crossed using 5x5 full diallel mating design. The use of the five parents produced

twenty different families. But in this study, only eight families made up of 8 to 10 individuals each according to their reaction to BPD have been selected.

A strain «Lebdi» of *P. megakarya* obtained from the Central laboratory of Phytopathology at the IRAD (Research Institute for Agricultural Development) Nkolbisson (central region, Cameroon). It was isolated in from a cocoa plantation of Leboudi (central region, Cameroon). The fungus maintained its aggressiveness by culturing on V8 agar medium, and the isolate is inoculated onto cocoa pods.

Hand-pollination and grafting

Crossing using hand-pollination techniques (Cilas, 1991) was conducted at the field in March and May, 2012. Seeds from the pods harvested from the experimental farms were sown in the nursery and 1086 hybrids plants were obtained. The parents were grafted in the nursery using bud wood. This vegetative-propagated (grafting) was done on non-specific young cocoa plantlets. This two techniques (hand-pollination and grafting) study was carried out at the SODECAO station at Mengang.

Leaf inoculation

The pathogenicity test was carried out in the nursery on leaves scarified along the midrib as described by Djocgoue et al. (2006) and Ondobo et al. (2014). The leaf test is an artificial inoculation method that can be used to assess the level of resistance in the genotypes. The inner surface of leaves was sterilized with ethanol 70%. Agar discs (6 mm diameter) cut from 5-day-old fungal and straminopilous isolates (oomycete) were laid on the midrib after creating wounds with a sterilized razor blade. The scars were then covered with cotton that had been immersed in sterilized water. The necrosis length was measured using a graduated ruler. The necrosis length was measured by a graduated ruler every two days after inoculation. The experiment was repeated thrice.

Estimation of the heterosis and heritability

The estimate of mid-parent heterosis was computed using the formula from Zahour (1992):

$$HF(\%) = \frac{F1 - \frac{P1 + P2}{2}}{\frac{P1 + P2}{2}} \times 100$$

Where: HF (%) = Mid-parent heterosis in percent; F1 = Means of hybrid genotype; P1 and P2 = Means of the two parents.

For necrosis length, heritabilities were estimated according to Falconer and Mackay (1996) following the formulas:

1. Broad-sense heritability

$$H^2 = \frac{V_G}{V_P} = \frac{V_G}{V_G + V_E}$$

2. Narrow sense heritability:

$$h^2 = \frac{V_A}{V_P} = \frac{V_A}{V_G + V_E}$$

Where: V_G = Genetic variance; V_P = Phenotypic variance; V_A = Additive genetic variance; V_E = Environmental variance.

Table 1. Origin and sensibilities of cocoa parental clones and their progenies of cocoa.

S/N	Parental clones/families	Origin	Collection	Group
1	SNK 16	Cameroon	Nkoemvone	Trinitario
2	SNK 413	Cameroon	Nkoemvone	Trinitario
3	T 79/467	Ghana	Tafo	Forastero
4	T 79/501	Ghana	Tafo	Forastero
5	SCA 12	Equateur	-	Forastero
6	F10: SCA 12 × T 79/467	Cameroon	Mengang	Forastero x Forastero
	F80: T 79/467 × SCA 12	Cameroon	Mengang	Forastero x Forastero
7	F15: T 79/501 × SCA 12	Cameroon	Mengang	Forastero x Forastero
	F61: SCA 12 × T 79/501	Cameroon	Mengang	Forastero x Forastero
8	F16: SNK 16 × T 79/501	Cameroon	Mengang	Trinitario x Forastero
	F79: T 79/501 × SNK 16	Cameroon	Mengang	Forastero x Trinitario
9	F30: SNK 413 × T 79/467	Cameroon	Mengang	Trinitario x Forastero
	F70: T 79/467 × SNK 413	Cameroon	Mengang	Forastero x Trinitario

Total phenolic compound contents

Total phenolic content was determined following the method describe by Singleton and Rossi (1965). A sample (50 mg) was extracted with 1 mL of 70% aqueous ethanol at room temperature, then centrifuged at 1000 g for 15 min. The supernatant (200 µL) was mixed with 1.5 mL of Folin-Ciocalteu reagent, and allowed to stand at room temperature for 5 min; then 1.5 mL of sodium bicarbonate solution (0.566 M) was added to the mixture. After 60 min, absorbance was read at 725 nm (Hitachi spectrometer U-200). Results were expressed as gallic acid equivalents. The concentration used was in a range between 0.02 and 0.1 mg/mL.

Total soluble amino acid and total soluble sugar contents

Amino acid contents were determined by using ninhydrine method of Yemm and Cocking (1955) with slight modifications. The incubation mixture containing 100 mL of the ethanol extract, 1 mL of 80% ethanol, 1 mL of 0.2 M citrate buffer (pH 5) and 2 mL of acetonic ninhydrin solution (1% ninhydrin and 0.006% KCN in acetone) was incubated for 15 min at 100°C. The mixture was cooled for 5 min under tap water before adding 8 mL of distilled water. The absorbance of the purple product was recorded at 570 nm (Hitachi spectrometer U-200). Glycine equivalents were calculated from a standard curve obtained with pure analytical grade glycine.

For carbohydrate determination, proteins were removed from the ethanolic extract after treatment with basic lead acetate. The carbohydrate extracts were then determined by using anthron method: 1mL of the extract was incubated in 5 mL of anthron solution (0.12 g anthron in 100 mL 6.5 M H₂SO₄) at 90°C for 10 min. The absorbance of the green product was measured at 630 nm. Results were expressed in µg eq. glucose by reference to the standard.

Proline contents

Proline contents were determined spectrophotometrically (Bates et al., 1973). Acid-ninhydrin was prepared by heating 0.7 g ninhydrin in 15 mL glacial acetic acid and 10 mL of 6 M phosphoric acid, with agitation till dissolution and stored at 4°C. Approximately 0.3 mg of plant material was homogenized in 8 mL of 3% aqueous sulfosalicylic acid and the homogenate filtered through Whatman

No2 filter paper. 2 mL of filtrate was reacted with 2 mL acid-ninhydrin and 1.5 mL of glacial acetic acid in a test tube for 1 h at 100°C, and the reaction terminated in an ice bath. The reaction mixture was extracted with 4 mL toluene, mixed vigorously with a test tube stirrer for 20 s. The chromophore containing toluene dissolved in the aqueous phase, warmed to room temperature and the absorbance read at 520 nm (Hitachi spectrometer U-200) using toluene as a blank. The proline concentration was determined from a standard curve and calculated on a fresh weight basis as follows: $[(\mu\text{g proline/mL} \times \text{mL toluene}) / 115.5 \mu\text{g}/\mu\text{mole}] / [(\text{g sample})/5] = \mu\text{moles proline/g of fresh weight material.}$

GABA contents

The GABA content was determined by following the procedure of Baum et al. (1996) with a minor modification. Ground cocoa leaves (200 mg) were added to a solution (800 mL) of methanol: chloroform: water mixture (12:5:3, volume basis) in a centrifuge tube. The tube was vortexed and then centrifuged at 12,000 g at 4°C for 15 min. The supernatant was collected in a flask and the residue was extracted again with a chloroform: water (3:5, v/v) solution (800 mL). The second supernatant was combined with the first supernatant. The collected sample was dried and then re-dissolved in water. The sample was then filtered through a 0.45 mm filter and the GABA content was measured by an amino acid analysis system (Waters, Milford, MA) after 6-aminoquiolyl-N-hydroxysuccinimidyl carbonate (AQC) derivatization.

Statistical analysis

Data presented are the means ± SE of three independent experiments. Analysis of variance (ANOVA) and the Tukey's test were performed using SPSS version 20.0. Principal component analysis (PCA) was used to describe the variability of necrosis length data. This analysis was performed with SPAD version 5.5.

RESULTS

Hand-pollination and grafting

Table 2 presents the results obtained from hand-

Table 2. Percentage of success of hand pollination of different hybridizations.

S/N	Famillies	Reciprocal crossing	Tests	Successful	% of successful	Observation
1	F10	(♀) SCA 12 × (♂) T 79/467	105	22	20.95	Viable
	F80	(♀) T 79/467 × (♂) SCA 12	108	25	23.14	Viable
2	F15	(♀) T 79/501 × (♂) SCA 12	105	43	40.95	Viable
	F61	(♀) SCA 12 × (♂) T 79/501	90	33	36.66	Viable
3	F16	(♀) SNK 16 × (♂) T 79/501	102	42	41.17	Viable
	F79	(♀) T 79/501 × (♂) SNK 16	100	31	31	Viable
4	F30	(♀) SNK 413 × (♂) T 79/467	92	36	39.13	Viable
	F70	(♀) T 79/467 × (♂) SNK 413	110	18	16	Viable
Total			812	250	31.12	

Table 3. Percentage of successful transplants parental clones.

Parents	Test graft	Successfull graft	Percent of successful
T 79/501	60	48	80
T 79/467	60	25	41.66
SNK 16	60	32	53.33
SNK 413	60	44	73.33
SCA 12	60	33	55
Total	240	148	60.68

pollination. The overall rate of success was 31.12%, with F10, F80 and F70 families having a lesser success of 16, 20.95 and 23.14% respectively. On the other hand, F15 (40.95%) and F16 (41.17%) families had the best percentage. The viability of all the seeds from reciprocal crosses showed the compatibilities of clones. However, after obtaining the hybrid plantlets four parental clones were obtained by grafting. The success rates varied between 41.66 and 80% (Table 2).

Development of the lesion

The rate of development of infection was genotype dependent. In all trials, there was a highly significant variation between the reactions of parental clones and their progenies to BPD (Table 3). On 2nd day after infection, necrotic lesions were observed in all parental and almost all hybrid genotypes, except for F15.02, F61.07 and F70.04 in the dry season (ds) and F15.08 and F70.08 genotypes in the rainy season (rs) (Table 4). From the fourth day, the necrosis evolved regularly in all the inoculated genotypes till the sixth day (Table 4). On the 4th day after infection, lowest necrotic lesions was recorded in 100 and 100% individuals of F10, 100 and 87.5% for F15, 100 and 62.5% for F30 in ds and rs respectively (Table 4). On the 6th day after infection by mycelium, the highest necrosis length was recorded for

genotypes F79.01 (7.36±0.63 cm), F79.03 (5.35±0.77 cm), F61.04 (5.27±1.67 cm), and F79.02 (5.03±0.50 cm) in the ds, and subsequently in the rs, F79.05 (6.30±0.17 cm), F79.07 (6.42±0.22 cm), F30.07 (6.63±0.38 cm) F79.08 (6.15±0.12 cm) and F30.02 (5.77±0.39 cm) genotypes also had the highest necrosis length (Table 4).

Heterosis and heritability

The percentage of genotypes with positive heterosis was higher in the ds (84.37%) than in rs (76.04%). Reciprocal families F10/F80 (best pair) had a hybrid vigor of 100 and 83.33% in dry and rainy seasons, respectively (Table 5). On the other hand, F16/F79 families had the lowest heterosis in ds (52.08%) and rs (68.75%) (Table 5).

The heritabilities (narrow-sense and broad-sense) for necrosis length were obtained for the eight families from reciprocal crosses. Concerning narrow sense heritability, the value obtained in the F10 family ($h^2 = 0.229$) was not significant to that obtained in F80 ($h^2 = 0.306$) (Table 6). However, such observations were made in reciprocal families F15-F61, F16-F79 and F30-F70. These values are coupled, respectively 0.212-0.203, 0.278-0.132 and 0.699-0.601 (Table 6). On the other hand, the broad-sense heritability showed that reciprocal families F10 ($H^2 = 0.329$) and F80 ($H^2 = 0.396$), F15 ($H^2 = 0.156$) and F61 ($H^2 = 0.232$), F16 ($H^2 = 0.375$) and F79 ($H^2 = 0.313$) had

Table 4. Length of necrosis of the parents and their offsprings in the nursery.

Genotypes	Length necrosis in cm (dry season)			Length necrosis in cm (rainy season)			
	Day 2	Day 4	Day 6	Day 2	Day 4	Day 6	
Parents	SCA 12	1.42±0.47 ^b	3.45±1.02 ^d	4.90±0.78 ^d	1.20±0.15 ^{ab}	2.33±0.29 ^b	5.85±1.05 ^e
	T 79/467	1.04±0.02 ^b	2.20±0.19 ^c	3.07±0.11 ^c	1.47±0.22 ^b	2.31±0.23 ^b	4.36±0.42 ^d
F10	F10.01	0.17±0.05 ^a	0.37±0.05 ^a	2.67±0.46 ^{bc}	0.69±0.15 ^a	2.33±0.17 ^b	3.81±1.21 ^c
	F10.02	0.23±0.05 ^a	0.54±0.05 ^a	2.10±0.36 ^b	0.23±0.15 ^a	0.47±0.05 ^a	1.43±0.05 ^a
	F10.03	0.27±0.25 ^a	0.70±0.20 ^a	1.73±0.58 ^a	0.62±0.12 ^a	1.27±0.20 ^a	2.07±0.61 ^b
	F10.04	0.10±0.17 ^a	0.60±0.26 ^a	2.13±0.89 ^b	0.60±0.17 ^a	1.36±0.14 ^a	4.63±1.16 ^d
	F10.05	0.30±0.17 ^a	0.75±0.32 ^a	1.36±0.40 ^a	1.13±0.27 ^{ab}	1.63±0.40 ^a	3.90±0.17 ^c
	F10.06	0.23±0.05 ^a	0.70±0.10 ^a	1.40±0.10 ^a	0.30±0.10 ^a	0.63±0.15 ^a	1.23±0.20 ^a
	F10.07	0.17±0.05 ^a	0.53±0.35 ^a	1.07±0.11 ^a	0.57±0.11 ^a	1.47±0.21 ^a	2.88±0.44 ^b
	F10.08	0.10±0.03 ^a	0.93±0.23 ^a	2.27±0.92 ^b	0.80±0.23 ^a	1.63±0.28 ^a	2.97±0.28 ^b
F80	F80.01	0.17±0.05 ^a	0.70±0.26 ^a	1.57±0.55 ^a	0.97±0.28 ^a	3.83±0.67 ^c	4.73±1.33 ^d
	F80.02	0.17±0.07 ^a	0.53±0.23 ^a	0.90±0.10 ^a	1.26±0.22 ^{ab}	4.00±1.20 ^{cd}	4.97±1.66 ^d
	F80.03	0.23±0.15 ^a	0.53±0.20 ^a	1.73±0.55 ^a	0.53±0.05 ^a	1.20±0.16 ^a	2.36±0.29 ^b
	F80.04	0.20±0.11 ^a	1.60±0.36 ^b	2.30±0.60 ^b	2.05±0.81 ^c	3.16±1.23 ^c	4.40±0.37 ^d
	F80.05	0.23±0.05 ^a	0.90±0.45 ^a	1.93±0.60 ^{ab}	2.12±0.23 ^c	2.63±0.35 ^b	3.91±1.03 ^c
	F80.06	0.27±0.05 ^a	0.43±0.11 ^a	1.00±0.30 ^a	0.83±0.47 ^a	1.69±0.36 ^a	2.95±0.66 ^b
	F80.07	0.13±0.05 ^a	0.63±0.15 ^a	1.03±0.15 ^a	0.67±0.27 ^a	1.47±0.81 ^a	2.67±0.58 ^b
	F80.08	0.67±0.27 ^a	1.47±0.81 ^b	2.67±0.58 ^{bc}	1.37±0.09 ^b	1.73±0.23 ^a	2.53±0.44 ^b
Parents	SCA 12	1.42±0.47 ^b	3.45±1.02 ^{bc}	4.90±0.78 ^{cd}	1.20±0.15 ^b	2.33±0.29 ^b	5.85±1.05 ^e
	T 79/501	0.72±0.06 ^a	2.10±0.17 ^{ab}	3.67±0.48 ^{bc}	1.56±0.31 ^b	2.13±0.20 ^b	4.13±0.38 ^c
F15	F15.01	0.60±0.17 ^a	2.20±0.79 ^{ab}	2.80±0.92 ^{ab}	0.63±0.11 ^a	1.20±0.17 ^a	3.83±0.28 ^c
	F15.02	0 ^a	2.53±0.46 ^{ab}	3.00±0.08 ^b	0.60±0.17 ^a	1.14±0.25 ^a	2.77±0.20 ^b
	F15.03	0.50±0.10 ^a	1.93±0.12 ^a	2.13±0.23 ^a	0.67±0.05 ^a	1.33±0.11 ^a	2.93±0.15 ^b
	F15.04	0.47±0.21 ^a	2.83±0.15 ^b	3.33±0.49 ^b	0.70±0.15 ^a	1.00±0.05 ^a	3.25±0.25 ^c
	F15.05	0.60±0.10 ^a	1.10±0.26 ^a	1.34±0.35 ^a	1.03±0.20 ^a	2.95±0.15 ^b	4.64±0.25 ^d
	F15.06	0.40±0.16 ^a	1.30±0.07 ^a	1.87±0.12 ^a	0.56±0.16 ^a	0.77±0.21 ^a	1.97±0.35 ^a
	F15.07	0.20±0.05 ^a	1.50±0.52 ^a	2.46±0.06 ^{ab}	0.43±0.23 ^a	0.63±0.36 ^a	1.58±0.05 ^a
	F15.08	0.53±0.12 ^a	2.40±0.35 ^{ab}	4.23±0.58 ^c	0 ^a	1.47±0.34 ^a	2.63±0.23 ^b
F61	F61.01	0.67±0.15 ^a	1.37±0.47 ^a	1.70±0.35 ^a	0.73±0.15 ^a	2.10±0.40 ^b	3.40±0.28 ^c
	F61.02	0.50±0.17 ^a	1.97±0.25 ^a	2.83±0.42 ^{ab}	0.67±0.20 ^a	2.46±0.51 ^b	4.87±0.17 ^d
	F61.03	0.53±0.06 ^a	1.57±0.15 ^a	2.06±0.21 ^a	0.63±0.11 ^a	1.30±0.16 ^a	2.57±0.51 ^b
	F61.04	0.72±0.12 ^a	4.80±0.69 ^{cd}	5.27±1.67 ^d	0.66±0.20 ^a	1.78±0.31 ^{ab}	2.73±0.30 ^b
	F61.05	0.75±0.15 ^a	3.00±0.87 ^b	3.33±0.99 ^b	0.70±0.12 ^a	2.50±0.29 ^b	3.90±0.30 ^c
	F61.06	0.67±0.15 ^a	1.17±0.57 ^a	1.50±0.44 ^a	0.90±0.28 ^a	2.84±0.17 ^b	5.34±0.23 ^e
	F61.07	0 ^a	2.53±0.46 ^{ab}	3.03±0.31 ^b	1.03±0.21 ^a	2.93±0.23 ^b	4.00±0.25 ^c
	F61.08	0.73±0.12 ^a	2.20±0.61 ^{ab}	3.80±0.20 ^{bc}	0.77±0.17 ^a	2.50±0.23 ^b	3.70±0.11 ^c
Parents	SNK 16	2.24±0.22 ^c	3.09±0.12 ^{bc}	4.09±0.44 ^c	1.06±0.21 ^a	2.97±0.06 ^b	5.27±0.44 ^d
	T 79/501	0.72±0.06 ^a	2.10±0.17 ^a	3.67±0.48 ^{bc}	1.56±0.31 ^b	2.13±0.20 ^b	4.13±0.38 ^c
F16	F16.01	1.76±0.32 ^b	3.33±0.04 ^{bc}	4.16±0.52 ^c	2.07±0.21 ^c	3.17±0.35 ^{bc}	4.23±0.37 ^c
	F16.02	1.33±0.40 ^{ab}	2.46±0.41 ^b	3.32±0.73 ^b	0.60±0.10 ^a	1.40±0.17 ^a	3.31±0.25 ^b
	F16.03	1.93±0.28 ^{bc}	4.80±0.69 ^d	5.18±0.04 ^{cd}	0.67±0.06 ^a	3.20±0.26 ^{bc}	4.13±0.21 ^c
	F16.04	1.66±0.45 ^b	3.50±1.00 ^c	4.46±0.27 ^c	0.60±0.13 ^a	1.53±0.15 ^a	2.63±0.12 ^a
	F16.05	1.63±0.65 ^b	3.20±0.72 ^{bc}	3.52±0.86 ^{bc}	0.83±0.12 ^a	1.83±0.14 ^a	3.44±0.20 ^b
	F16.06	1.20±0.52 ^{ab}	2.13±0.11 ^a	2.90±0.52 ^b	1.36±0.23 ^{ab}	4.73±0.25 ^{cd}	6.20±0.26 ^e
	F16.07	1.56±0.11 ^b	2.86±0.23 ^b	3.63±0.40 ^{bc}	0.83±0.23 ^a	2.53±0.21 ^b	4.70±0.20 ^c
	F16.08	0.73±0.05 ^a	2.00±0.51 ^a	2.46±0.25 ^a	1.22±0.68 ^{ab}	1.43±0.46 ^a	2.48±0.84 ^a
F79	F79.01	1.23±0.25 ^{ab}	5.83±0.76 ^e	7.36±0.63 ^e	0.70±0.14 ^a	1.53±0.25 ^a	2.90±0.17 ^a

Table 4.Contd.

	F79.02	1.10±0.10 ^a	4.23±0.25 ^d	5.03±0.50 ^{cd}	0.90±0.17 ^a	3.26±0.25 ^c	5.03±0.15 ^d
	F79.03	2.60±0.40 ^c	3.63±0.70 ^c	5.35±0.77 ^{cd}	0.78±0.06 ^a	1.33±0.15 ^a	3.30±0.30 ^b
	F79.04	0.63±0.11 ^a	1.00±0.06 ^a	2.00±0.17 ^a	0.80±0.20 ^a	2.94±0.20 ^b	4.57±0.31 ^c
	F79.05	1.83±0.23 ^b	3.33±0.66 ^{bc}	4.12±0.06 ^c	1.20±0.26 ^{ab}	4.37±0.23 ^{cd}	6.30±0.17 ^e
	F79.06	0.70±0.04 ^a	1.76±0.49 ^a	2.46±0.30 ^a	0.87±0.15 ^a	3.15±0.32 ^{bc}	5.00±0.50 ^d
	F79.07	1.46±0.72 ^b	2.45±0.89 ^b	3.83±0.56 ^{bc}	1.26±0.21 ^{ab}	4.70±0.26 ^{cd}	6.42±0.22 ^e
	F79.08	0.70±0.09 ^a	1.76±0.49 ^a	1.46±0.30 ^a	0.80±0.10 ^a	3.03±0.45 ^{bc}	6.15±0.12 ^e
Parents	T 79/467	1.04±0.02 ^{ab}	2.20±0.19 ^c	3.07±0.11 ^c	1.47±0.33 ^b	2.31±0.05 ^b	4.36±0.12 ^d
	SNK 413	0.98±0.01 ^a	2.37±0.02 ^c	2.76±0.01 ^c	1.16±0.31 ^b	2.06±0.16 ^b	3.70±0.46 ^c
	F30.01	0.23±0.08 ^a	1.33±0.64 ^b	1.93±0.10 ^b	0.55±0.15 ^a	1.23±0.25 ^{ab}	2.21±0.25 ^b
F30	F30.02	0.33±0.01 ^a	1.00±0.80 ^a	1.80±0.01 ^b	0.50±0.12 ^a	2.88±0.50 ^b	5.77±0.39 ^e
	F30.03	0.43±0.12 ^a	1.77±0.14 ^b	2.34±0.36 ^b	0.23±0.11 ^a	1.33±0.64 ^{ab}	1.92±0.13 ^a
	F30.04	0.45±0.15 ^a	2.20±0.47 ^c	2.60±0.58 ^b	0.80±0.17 ^a	2.70±0.20 ^b	4.15±0.42 ^d
	F30.05	0.17±0.06 ^a	1.60±0.44 ^b	2.10±0.10 ^b	1.33±0.25 ^b	2.26±0.35 ^b	3.33±0.31 ^c
	F30.06	0.26±0.12 ^a	0.62±0.12 ^a	1.07±0.40 ^a	0.23±0.02 ^a	0.60±0.12 ^a	1.07±0.40 ^a
	F30.07	0.13±0.05 ^a	0.50±0.06 ^a	0.72±0.25 ^a	0.63±0.15 ^a	2.43±0.30 ^b	6.63±0.38 ^f
	F30.08	0.20±0.09 ^a	0.53±0.15 ^a	1.13±0.76 ^a	1.16±0.31 ^b	2.06±0.16 ^b	3.70±0.46 ^c
	F70.01	0.23±0.06 ^a	1.10±0.17 ^a	2.03±0.61 ^b	0.50±0.13 ^a	0.86±0.25 ^a	2.26±0.19 ^b
F70	F70.02	0.33±0.15 ^a	1.41±0.85 ^b	2.17±0.32 ^b	0.50±0.20 ^a	2.73±0.26 ^b	3.70±0.12 ^c
	F70.03	0.27±0.03 ^a	1.20±0.55 ^b	1.43±0.35 ^a	0.33±0.10 ^a	2.80±0.21 ^b	3.57±0.24 ^c
	F70.04	0 ^a	0.32±0.03 ^a	0.68±0.12 ^a	0.67±0.06 ^a	2.53±0.72 ^b	4.55±0.64 ^d
	F70.05	0.30±0.07 ^a	0.54±0.05 ^a	0.73±0.25 ^a	0.75±0.17 ^a	2.93±0.52 ^b	4.73±0.46 ^d
	F70.06	0.34±0.06 ^a	1.00±0.68 ^a	1.60±0.72 ^b	0.89±0.31 ^a	1.73±0.27 ^{ab}	3.98±0.38 ^c
	F70.07	0.13±0.04 ^a	0.47±0.25 ^a	1.27±0.49 ^a	0.13±0.05 ^a	0.47±0.25 ^a	1.27±0.49 ^a
	F70.08	0.27±0.12 ^a	2.37±0.25 ^c	2.76±0.66 ^c	0 ^a	0.33±0.03 ^a	0.67±0.12 ^a

*Values with the same letter in the same column and in the same family are not significant ($P < 0.05$) different. Values are means of 3 replicates.

low values (Table 6). F30 ($H^2 = 0.624$) and F70 ($H^2 = 0.643$) families had relatively high values (Table 6).

Principal component analysis (PCA)

Principal component analysis (PCA) was carried out on the data from all the clones and hybrids studied during the development of necrosis. For the dry season, the first two principal component (PC) generated from all data represented 97.90% of the total variability of necrosis. In the first principal component, necrosis at 4th and 6th days after inoculation contributed to 86.64% of the total variability while in the second principal component, necrosis at the 2nd day after inoculation contributed 11.27% of the total variability (Figure 1). Examining a two-dimensional scores plot in the space defined by PC1 and PC2 shows that distribution of genotypes followed a specific pattern. Genotype F79.01 (highly susceptible to BPD) represented Group 1; group 2 constituted of three genotypes (F61.04, F79.02 and F16.03) that were susceptible to BPD; group 3 consisted of two parental clones (SCA 12 and SNK 16) and four hybrids (F16.04, F16.01, F79.06 and F79.03) that were moderately susceptible to BPD. Group 4 was very homogeneous and

also distinguished the best parents in terms of increasing tolerance (T 79/501, T 79/467 and SNK 413) and 25 hybrids (list them) that were tolerance to BPD The genotypes in group revealed high tolerance (Figure 1).

For the rs, the first two principal components represented 95.30% of the total variability of necrosis length. Days 4, and 6 were the dominant features in the first PC1 (76.39% of the total variability) while necrosis at day 2 was the highest feature in the second PC2 (18.91% of the total variability) (Figure 2). Only F79.01 represented the first group, marked by a higher necrosis length. The second group included the susceptible parents (SCA 12 and SNK 16) and 23 hybrids that presented significant necrosis length; however, it was less important in the third group consisting of the best parents T 79/501, T 79/467, SNK 413 and hybrids that formed the group. The fourth group embodied 30 hybrids, characterized by lower necrosis length (Figure 2).

The ranking of parents for necrosis length, allowed classifying more efficient genotypes other than the best parent "SNK 413". F10, F15, F30, F61, F70, F80 families respectively presented 6 (F10.02, F10.03, F10.05, F10.06, F10.07 and F10.08), 3 (F15.03, F15.06 and F15.07), 4 (F30.03, F30.05, F30.07 and F30.08), 1 (F61.03), 2 (F70.04, F70.05), 3 (F70.01, F70.07 and

Table 5. Estimation of mid-parent heterosis (%) of necrosis length from the parents and their offsprings in nursery.

Genotypes	Dry season			Rainy season			Genotypes	Dry season			Rainy season			
	Day 2	Day 4	Day 6	Day 2	Day 4	Day 6		Day 2	Day 4	Day 6	Day 2	Day 4	Day 6	
F10	F10.01	-86.18	-86.90	-60.60	-48.31	+0.43	-25.29	F16.01	+18.92	+28.32	+7.22	+58.02	+24.31	-10
	F10.02	-81.30	-80.88	-77.42	-82.77	-79.74	-71.96	F16.02	-10.14	-5.20	-14.43	-54.20	-45.10	-29.57
	F10.03	-78.05	-75.22	-56.59	-53.56	-45.26	-59.41	F16.03	+30.41	+84.97	+33.51	-48.85	+25.49	-12.13
	F10.04	-91.87	-78.76	-42.28	-55.06	-41.38	-9.22	F16.04	+12.16	+34.87	+14.95	-54.20	-40	-44.04
	F10.05	-75.61	-73.45	-51.57	-15.36	-29.74	-23.53	F16.05	+10.14	+23.31	-9.28	-36.64	-28.24	-26.81
	F10.06	-81.30	-75.22	-74.91	-77.53	-72.84	-75.88	F16.06	-18.92	-17.92	-25.26	+3.82	+85.49	+31.91
	F10.07	-86.18	-81.24	-74.15	-57.30	-36.64	-43.53	F16.07	+5.41	+10.21	-6.44	-36.64	-0.78	0
	F10.08	-91.87	-67.08	-33	-40.07	-29.74	-41.76	F16.08	-50.68	-22.93	-36.60	-6.87	-43.92	-47.23
F80	F80.01	-86.18	-75.22	-60.60	-27.34	+65.09	-7.25	F79.01	-16.89	+124.66	+89.69	-46.56	-40	-38.30
	F80.02	-86.18	-81.24	-77.42	-5.62	+72.41	-2.55	F79.02	-25.68	+63.01	+29.64	-31.30	+27.84	+7.02
	F80.03	-81.30	-81.24	-56.59	-60.30	-48.28	-53.73	F79.03	+75.68	+39.88	+37.89	-40.46	-47.84	-29.79
	F80.04	-83.74	-43.36	-42.28	+53.56	+36.21	-13.73	F79.04	-57.43	-61.46	-48.45	-38.93	+15.29	-2.77
	F80.05	-81.30	-68.14	-51.57	+58.80	+13.36	-23.33	F79.05	+23.65	+28.32	+6.19	-8.40	+71.37	+34.04
	F80.06	-78.05	-84.78	-74.91	-37.83	-27.16	-42.16	F79.06	-52.70	-32.18	-36.60	-33.59	+23.53	+6.38
	F80.07	-89.43	-77.70	-74.15	-49.81	-36.64	-47.65	F79.07	-1.35	-5.59	-1.29	-3.82	+84.31	+36.60
	F80.08	-45.53	-47.96	-33	+2.62	-25.43	-50.39	F79.08	-52.70	-32.18	-62.37	-38.93	+18.82	+30.85
F15	F15.01	-43.93	-20.72	-34.66	-54.35	-46.19	-23.25	F30.01	-77.23	-41.79	-33.79	-58.17	-43.71	-45.16
	F15.02	-100	-8.83	-29.99	-56.52	-48.88	-44.49	F30.02	-67.33	-56.24	-38.25	-61.98	+31.81	+43.18
	F15.03	-53.27	-30.45	-50.29	-51.45	-40.36	-41.28	F30.03	-57.43	-22.54	-19.73	-82.51	-39.13	-52.36
	F15.04	-56.07	+1.98	-22.29	-49.28	-55.16	-34.87	F30.04	-55.45	-3.72	-10.81	-39.16	+23.57	+2.98
	F15.05	-43.93	-60.36	-68.73	-25.36	+32.29	-7.01	F30.05	-83.17	-29.98	-27.96	+1.14	+3.43	-17.37
	F15.06	-62.62	-53.15	-56.36	-59.42	-65.47	-60.52	F30.06	-74.26	-72.87	-63.29	-82.51	-72.54	-73.45
	F15.07	-81.31	-45.95	-42.59	-68.84	-71.75	-68.34	F30.07	-87.13	-78.12	-75.30	-52.09	+11.21	+64.52
	F15.08	-50.47	-13.51	-1.28	-100	-34.08	-47.29	F30.08	-80.20	-76.81	-61.23	-11.79	-5.72	-8.19
F61	F61.01	-37.38	-50.63	-60.33	-47.10	-5.83	-31.86	F70.01	-77.23	-51.86	-30.36	-61.98	-60.64	-43.92
	F61.02	-53.27	-29.01	-33.96	-51.45	+10.31	-2.40	F70.02	-67.33	-38.29	-25.56	-61.98	+24.94	-8.19
	F61.03	-50.47	-43.42	-51.93	-54.35	-41.70	-48.50	F70.03	-73.27	-47.48	-50.94	-74.90	+28.15	-11.41
	F61.04	-32.71	+72.97	+22.99	-52.17	-20.18	-45.29	F70.04	-100	-86	-76.67	-49.05	+15.79	+12.90
	F61.05	-29.91	+8.11	-22.29	-49.28	+12.11	-21.84	F70.05	-70.30	-76.37	-74.96	-42.97	+34.10	+17.37
	F61.06	-37.38	-57.84	-64.99	-34.78	+27.35	+7.01	F70.06	-66.34	-56.24	-45.11	-32.32	-20.82	-1.24
	F61.07	-100	-8.83	-29.29	-25.36	+31.39	-19.84	F70.07	-87.13	-79.43	-56.43	-90.11	-78.49	-68.49
	F61.08	-31.78	-20.72	-11.32	-44.20	+12.11	-25.85	F70.08	-73.27	+3.72	-5.32	-100	-84.90	-83.37




Figure 2. Principal component analysis based on the length of necrosis on *T. cacao* leaf 2nd, 4th and 6th days after inoculation for parental and hybrid cocoa genotypes in rainy season.




Figure 3. Classification of the different parents and their progenies according to their susceptibility to black pod.

individuals (Figure 5).

DISCUSSION

The production of parental and hybrid genotypes of cocoa was achieved through hand pollination and

grafting. The percentage of successful crosses recorded for hand pollination (31.12%) was low. This could be due to genetic incompatibility between clones and also the period of pollination. This result is similar to those found by Mossu (1990) and Ondobo et al. (2014).

The percentage of success for the regeneration parents formed by grafting varied from 41.66 to 80% depending




Figure 4. Relationship between length of necrosis in rainy and dry seasons on cocoa leaves after inoculation with mycelium of *P. megakarya* on (a) 2nd day (b) 4th day (c) 6th day.

on the clone used. According to Akinnifesi et al. (2008), they observed that the success of grafting for *Uapaca kirkiana* depends on the skill of the grafter, management practices used after grafting, and the technique used. In this study, the period of grafting and the non-compatibility of rootstock and graft could have caused the low success rate.

The appearance of necrosis on the midrib of the leaves infected in nursery, confirms the presence of *P. megakarya* mycelium. This result is in agreement with the findings of Nyasse (1997). from the leaf disc test done in the laboratory, Efombagn et al. (2011) on pods in field, Djocgoue et al. (2010) and Ondobo et al. (2014) on the leaves attached to the plant in nursery. There were significant ($P < 0.05$) differences in the necrosis length of susceptible SCA 12 and SNK 16 as compared to T 79/501, T 79/467 and SNK 413.

Forty hybrid genotypes would have shown lower necrosis length to *P. megakarya* relative to the best parent (SNK 413). Earlier studies (Nyasse et al., 2002, Efombagn et al., 2011; Ondobo et al., 2014) showed that tolerance to black pod disease was under genetic control and can be improved genetically. Otherwise, the effects

of global combinations aptitudes to parental genotypic was substantial for the rot rate of cocoa pods (Cilas et al., 2004; Ondobo 2014), suggesting a primary and additive transmission of resistant characters (Tan and Tan, 1990).

The manifestation of hybrid vigor was observed between dry season (84.37%) and rainy season (76.04%) in all the hybrid families. Cilas et al. (1998) and Djocgoue et al. (2006) showed that, individuals which exhibit hybrid vigor, would imply the presence of the additive and dominant gene effect in the transmission of character. However, negative heterosis observed in hybrid genotypes could be explained by the epistatic effect of some genes which tend to mask genes controlling tolerance to *P. megakarya*. Similar results were also obtained by Djocgoue et al. (2007) and Ondobo et al. (2014) when leaves of *T. cacao* were inoculated with *P. megakarya*. Early studies of Mohammadi et al. (2012) showed that, resistance to *Septoria tritici* blotch is controlled by additive, dominance and epistatic gene action, but the role of the dominant gene effect is greater than the others.

The values of narrow sense (h^2) and broad-sense (H^2) heritabilities of necrosis length (dry and rainy season) in




Figure 5. (a) Content of phenolic compounds, (b) proline, (c) GABA and (d) sugars during dry season (DS) and rainy season (RS) on cocoa leaves 6th day after inoculation with mycelium of *P. megakarya*.

the reciprocal crossings were not significant. The studied character, the absence of a significant difference between the heritability values from reciprocal crossing portrays the absence of maternal heritability. This finding suggested that the transmission of this character would not cytoplasmic but nuclear (Djocgoue et al., 2011). Alone values of heritability in the two reciprocal crossings [F30 ($h^2 = 0.699$ and $H^2 = 0.624$) and F70 ($h^2 = 0.601$ and $H^2 = 0.643$)] was showed a variation of character strongly inheritable ($h^2 > 0.4$). These results matched with those of Djocgoue et al. (2010) and Manga et al. (2016). In addition, in the experimental conditions of the present investigations, the parental and hybrid genotypes were planted in the same plot, and this had the effect of minimizing environment related effects, rendering the heritability estimations more trustworthy (Cilas, 1991). Therefore, selection should be conducted in the advanced generations of selfing when the breeding materials can be duplicated for extensive evaluation. Then, selection based on family performance or progeny test should be more effective (Phudenpa et al., 2004).

PCA based on necrosis length categorized all the

families into five (dry season) and four (rainy season) groups. Each group consists of similar individuals characterized by low and high length of necrosis. The tolerant hybrids characterized by low and intermediate length of necrosis were considered as elites. This result confirms a good aptitude (cross) to parental gene combinations (Cilas et al., 2004; Ondobo et al., 2013).

Regression studies showed that *P. megakarya* is more virulent in the rainy season than in dry season and this virulence decreases with time of infection. During rainy season, oomycete develop their sporangia rapidly because of high humidity. The sporangia of many oomycetes may germinate directly to form an infection hypha, or else in the presence of abundant water they may differentiate, through specialized cleavage vesicles, into 10 to 30 zoospores that can individually disperse to initiate sites of infection (Birch and Cooke, 2004). In this study, necrosis appeared two days after young leaves of cocoa were inoculated.

Variation of metabolites in cocoa genotypes 6th day after inoculation with *P. megakarya* was seen in the increase of phenols, proline and GABA and a decrease in

sugar contents. This increase is more important in rainy season than dry season. Phenols and amino acids have been considered by several authors (Djougoue et al., 2011; Omokolo and Boudjeko, 2005) as markers of resistance to fungal infections in plants. Del Rio et al. (2003) in the study of the enhancement of phenolic compounds in olive plants (*Olea europaea* L.) and their influence on resistance against *Phytophthora* sp. noticed that the HPLC-MS studies pointed to an increase in the phenol content of leaves 120 days after treatment with 0.3% Brotomax. These authors stated that oleuropein, catechin and tyrosol are some of the main phenolic compounds produced after *Phytophthora* sp. attack plants. Increase in proline content in the rainy season as compared to the dry season contrasted the result of Szabados and Savoure (2009). They noticed that during osmotic stress, proline biosynthesis is augmented in the chloroplasts, and this is controlled by the stress induced P5CS1 gene in Arabidopsis.

Conclusion

Resistance tests on leaf were used to select clones and hybrid families less sensitive. The estimation of genetic parameters has achieved comparative hybrid tests of genotypes, for the most effective characters among those analyzed and to identify the parental genotypes that show good combining potentials. 84.37 and 76.04% of hybrid genotypes exhibited positive heterosis (hybrid vigor) in dry and rainy seasons respectively. However, inoculation periods (dry and rainy season) did not show a high distinction. Transmission of the character would thus seem to be governed by primarily additive gene effect suggesting a nuclear origin of the transmission of these characters.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGMENTS

A part of this study was supported by the Agence Universitaire de la Francophonie (AUF) via a grant to Pierre Effa Onomo (Grant No BACGL-2015-35). The authors express gratitude to the Cameroon Cocoa Development Corporation (SODECAO) for the field used.

REFERENCES

Akinnifesi FK, Sileshi G, Mkonda A, Ajayi OC, Mhango J, Chilanga T (2008). Germoplasm supply, propagation and nursery management of miombo fruit trees. CABI Publishing, Wallingford, UK 341-368.
 Bates LS, Waldren RP, Teare ID (1973). Rapid determination of free proline for water-stress studies. *Plant Soil* 39(1): 205-207.

Baum G, Lev-Yadun S, Fridmann Y, Arazi T, Katsnelson H, Zik M (1996). Calmodulin binding to glutamate decarboxylase is required for regulation of glutamate and GABA metabolism and normal development in plants. *The EMBO J.* 15:2988-2996.
 Birch PRJ, Cooke D (2004). Mechanisms of infection: Oomycetes. *Encyclopedia of Plant Crop Sci.* 1:697-700.
 Blaha G (1974). Methods of testing for resistance. In: *Phytophthora Disease of Cocoa* (PH Gregory, ed), Longman, Londres, Royaume-Uni, 259-268.
 Blaha G and Lotodé R (1976). Un caractère primordial de sélection du cacaoyer au Cameroun : la résistance à la pourriture brune des cabosses. *Café Cacao Thé.* 20:97-116.
 Blaha G, Lotodé R (1977). Contribution à la connaissance des modalités de la transmission héréditaire de la résistance du cacaoyer à la pourriture des cabosses (*Phytophthora palmivora*) au Cameroun. *Café Cacao Thé.* 21:179-196.
 Brading PA, Verstappen ECP, Kema GHJ, Brown JKM (2002). A gene-for-gene relationship between wheat and *Mycosphaerella graminicola*, the *Septoria tritici* blotch pathogen. *Phytopathol.* 92:439-445.
 Cilas C (1991). Estimation of some genetics parameters of different crosses plans of cocoa. *Café Cacao Thé.* 25: 3-13.
 Cilas C, Lanaud C, Paulin D, Nyasse S, Ngoran JA, Kebe BI, Ducamp M, Flamant MH, Risteruci AM, Pieretti, Saunigo O, Thevenon JM, Despraux D (1998). La résistance à la pourriture des cabosses due à *Phytophthora* spp. Recherche des composantes de la résistance. *Plantation, Recherche, Développement.* 5: 441-445.
 Cilas C, Ndounbé-Nkeng M, Bidzanga N, N'goran J (2004). Incidence de la maladie et de résistance sur le terrain. Dans: Cilas C. et Despraux D. (eds). *Amélioration de la résistance des arbres de cacao à la maladie de Phytophthora. Repères*, CIRAD, Paris, France. pp. 171-193.
 Del Rio JA, Baideza AG, Botiab JM, Ortuno A (2003). Enhancement of phenolic compounds in olive plants (*Olea europaea* L.) and their influence on resistance against *Phytophthora* sp. *Food Chem.* 83:75-78.
 Despraux D, Cambrony D, Clément D, Partiot M (1988). Étude de la pourriture brune des cabosses du cacaoyer au Cameroun : définition de nouvelles méthodes de lutte. In : *Proc 10th Int Cocoa Res Conf*, Santo Domingo, République dominicaine (Stephen Austin and sons, eds), pp. 407-412.
 Djougoue PF, Boudjeko T, Nankeu DJ, Efombagn MIB, Nyassé S, Omokolo DN (2006). Comparative assessment of the resistance of Cocoa (*Theobroma cacao* L.) progenies from SNK10 x SNK413; ICS84 x ICS95 to *Phytophthora megakarya* in Cameroon by measuring size of necrotic lesion along the midrib. *Plant Pathol. J.* 5:329-333.
 Djougoue PF, Boudjeko T, Mbouobda HD, Nankeu DJ, El Hadrami I, Omokolo ND (2007). Heritability of Phenols in the Resistance of *Theobroma cacao* against *Phytophthora megakarya*, the Causal Agent of Black Pod Disease. *J. Phytopathol.* 155:519-525.
 Djougoue PF, Simo C, Mbouobda HD, Boudjeko T, Nankeu DJ, Omokolo ND (2010). Assessment and heritability of productivity and tolerance level to *Phytophthora megakarya* in two hybrid populations of *Theobroma cacao*. *J. Plant Pathol.* 92:607-617.
 Djougoue PF, Mbouobda HD, Boudjeko T, Effa OP, Omokolo ND (2011). Amino acids, carbohydrates and heritability of resistance in the *Theobroma cacao/Phytophthora megakarya* interaction. *Phytopathol. Mediterr.* 50:370-383.
 Effa OP, Niemenak N, Djougoue PF, Ondobo ML, Omokolo ND (2015). Heritability of polyphenols, anthocyanins and antioxidant capacity of Cameroonian cocoa (*Theobroma cacao* L.) beans. *Afr. J. Biotechnol.* 36:2672-2682.
 Efombagn MIB, Bieysse D, Nyassé S, Eskes AB (2011). Selection for resistance to *Phytophthora* pod rot of cocoa (*Theobroma cacao* L.) in Cameroon: Repeatability and reliability of screening tests and field observations *Crop Protection.* 30:105-110.
 Eskes A, Lanaud C (1997). Le cacaoyer. In: *L'amélioration des plantes tropicales*. CIRAD. Montpellier, France, pp. 141-170.
 Falconer DS, Mackay TFC (1996). *Introduction to Quantitative Genetics*, Longmans Green Ed. 4. Harlow, Essex, UK, P 125.
 Kema GHJ, Goodwin SB, Hamza S, Verstappen ECP, Cavaletto JR,

- Van der Lee TAJ, De Waard M, Bonants PJM, Waalwijk C (2002). A combined amplified fragment length polymorphism and randomly amplified polymorphism DNA genetic linkage map of *Mycosphaerella graminicola*, the *Septoria tritici* leaf blotch pathogen of wheat. *Genet.* 161:1497-1505.
- Manga NJ, Effa OP, Ondobo ML, Djoko KJC, Djocgoue PF (2016). Heritability of the tolerance to *Phytophthora megakarya* Bras. and Grif. of *Theobroma cacao* L. in terms of their necrosis length, phenolic contents and activity of enzymes. *Int. J. Bio. Sci.* 5:249-261.
- Mohammadi M, Ramezani S, Navabpour S, Soltanloo H, Kalateharabi M, Kia S (2012). Genetic analysis and heritabilities of resistance to *Mycosphaerella graminicola* in wheat. *Crop Breed. J.* 1:35-42.
- Mossu G (1990). Le cacaoyer. Maisonneuve et Larousse (Paris). p.321.
- Ndombé-Nkeng NM (2002). Incidence des facteurs agro-écologiques sur l'épidémiologie de la pourriture brune des fruits du cacaoyer au Cameroun: contribution à la mise en place d'un modèle d'avertissements agricoles. Thèse de Doctorat de l'institut National agronomique (Paris- Grignon), p.151.
- Nyasse S, Cilas C, Héral C, Blaha G (1995). Leaf inoculation as an early screening test for cocoa (*Theobroma cacao* L.) resistance to *Phytophthora* black pod disease. *Crop Protection.* 14:657-663.
- Nyasse S (1997). Etude de la diversité de *Phytophthora megakarya* et caractérisation la résistance du cacaoyer *Theobroma cacao* L. à cet agent pathogène. Thèse de doctorat, Institut National de Polytechnique de Toulouse (France), P 133.
- Nyasse S, Despréaux D, Cilas C (2002). Validity of a leaf inoculation test to assess the resistance to *Phytophthora megakarya* in cocoa (*Theobroma cacao* L.) diallel mating design. *Euphytica.* 123: 395-399.
- Omokolo ND, Boudjeko T (2005). Comparative analyses of alterations in carbohydrates, amino acids, phenols and lignin in roots of three cultivars of *Xanthosoma sagittifolium* infected by *Pythium myriotylum*. *South African Journal of Botany* 71(3):432-440.
- Ondobo ML, Effa OP, Djocgoué PF, Boudjeko T, Manga NJ, Djoko KJC, Omokolo ND (2013). Influence of *Phytophthora megakarya* inoculation on necrosis length, phenolic content, peroxidase and polyphenoloxidase activity in cocoa (*Theobroma cacao* L.) plants. *Syllabus Review, Science Series.* 4:8-18.
- Ondobo ML, Effa OP, Djocgoue PF, Manga NJ, Boudjeko T, Omokolo ND (2014). Phenolic content and heritability of resistance in four hybrid populations of *Theobroma cacao* L. after Leaves inoculation with *Phytophthora megakarya* Bras. and Grif. *Int. J. Biol. Chem. Sci.* 8:17-30.
- Partiot M (1975). La résistance horizontale du cacaoyer au *Phytophthora* spp. *Café Cacao Thé.* 19 :123-130.
- Phudenpa A, Jogloy S, Toomsan B, Wongkaew S, Kesmla T, Patanothai A (2004). Heritability and phenotypic correlation of traits related to N₂-fixation and agronomic traits in peanut (*Arachis hypogaea* L.). *Songklanakarin J. Sc. Techn.* 26:317-325.
- Simon MR, Cordo CA (1998). Diallel analysis of the resistance components to *Septoria tritici* in *Triticum aestivum*. *Plant Breed.* 117:123-126.
- Singleton VL, Rossi JA (1965). Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am. J. Enol. Vitic.* 16:144-158.
- Szabados L, Savoure A (2009). Proline: a multifunctional amino acid. *Trends in Plant Science.* 2:89-97.
- Tahi M, Keke I, Eskes AB, Ouattara S, Sangaré A, Mondeil F (2000). Dépistage rapide des génotypes de cacao pour la résistance à *Phytophthora palmivora* en champ en utilisant des brindilles de feuilles et de racines. *Eur. J. Plant Pathol.* 106:87-94.
- Tan GY, Tan WK (1990). Additive inheritance of resistance to pod rot caused by *Phytophthora palmivora* in cocoa. *Theor. Appl. Genet.* 80:258-264.
- Tarjot M (1969) Étude de la résistance des cacaoyers à la pourriture brune des cabosses due à *Phytophthora palmivora* (Butl) Butl en Côte d'Ivoire. 3e partie: inoculations expérimentales sur le terrain. *Café Cacao Thé.* 13:297-309.
- Tijani AA (2005). Profitability of Fungicide Use Decision among Cocoa Farmers in Southwestern Nigeria. *J. Soc. Sci.* 2:165-171.
- Yemm EW, Cocking EC (1955). The determination of amino acid with ninhydrin. *Analyst,* 80: 209-213.
- Zahour A (1992). *Eléments d'amélioration génétique des plantes.* Ed. Actes: Rabat, P 232.