

Short Communication

Crassulacean acid metabolism (CAM) in leaves of a common weed *Commelina nudiflora* L.

R. B. Deshmukh^{1*} and C. V. Murumkar²

¹Department of Botany, Pune University Affiliated Shardabai Pawar Mahila Mahavidyalaya, Shardanagar, Baramati, District Pune, Maharashtra, -413 115, India.

²Department of Botany, Pune University Affiliated Tuljaram Chaturchand College, Baramati, District Pune, Maharashtra - 413 112, India.

Accepted 16 September, 2013

Survey of weed flora of Baramati region has revealed that *Commelina nudiflora* L. is a dominant weed among common weeds in croplands and fallow lands. In order to ascertain the survival potential of *C. nudiflora* some photosynthetic parameters along with water relations of this plant were investigated. Baramati region is a drought prone hot semi-arid area in Pune, Maharashtra. *C. nudiflora*, belonging to the family Commelinaceae is a typical succulent plant and this was confirmed further by water relation studies like moisture content, osmotic potential, leaf water content and succulent index. Besides above mesophyll succulence, surface expansion and succulence index values were within the range of a typical crassulacean acid metabolism (CAM) plant. Further studies on diurnal fluctuations in pH status, stomatal behaviour and acidity status also confirmed its CAM mode of photosynthetic behaviour. All these findings led to the conclusion that CAM is in an operation in this weed.

Key words: Crassulacean acid metabolism (CAM), *Commelina nudiflora* L., weeds.

INTRODUCTION

Baramati is a drought prone hot semi-arid region of Pune, Maharashtra, India is severely affected by rain shadow effect of Western Ghat (Blatter and Hallberg, 1984). The study area lies in between 18° 3'N latitude and 74° 13'-74° 40'E longitude having a total geographic area of 1, 38 and 247 ha, receiving 530.2 mm of average rainfall. Maximum and minimum temperatures are 43 and 19°C respectively, with mean daily temperature of above 22°C (Bhagat et al., 2008).

Preliminary study revealed that *C. nudiflora* grows well even under unfavourable conditions like extreme hot summer as a dominant weed in crop fields and fallow lands (Murumkar and Magdum, 1991). Morphotypical and physiotypical plasticity of crassulacean acid metabolism (CAM) phenotypes allows wide ecophysiological amplitude of niche occupation in the tropics (John, 2001). But

physiological and biochemical plasticity appears more responsive than do morphological adaptations. This makes CAM plants particularly fit for the multi-factor stressor networks of tropical forests (Luttge, 2012).

Plants which have acclimatised to a certain habitat always show morphological, reproductive, phenological and physiological adaptations for successful survival. In the present work we studied the relationship in between survival potential and water relations, mode of photosynthesis in *C. nudiflora* plants.


MATERIALS AND METHODS

Plant populations of *C. nudiflora* in an open field were selected for studies.

*Corresponding author. E-mail: rbd_spmc@rediffmail.com or rbd.spmc@gmail.com.

Table 1. Some leaf characteristics of *Commelina nudiflora* L.

Parameter	Value
Leaf thickness (mm)	1.68 ± 0.012
Moisture content (%)	85.3 ± 3.18
Succulence index (SI) (g.g ⁻¹)	11.02
Mesophyll succulence (Sm) Kg.g ⁻¹ chl.	5.88
Surface expansion (m ² .kg ⁻¹)	1.76

**Figure 1.** Diurnal Variation in Titrable Acidity status pH of leaf tissues and stomatal behaviour (Black bar indicates the night? Day light absent time).

The investigation was conducted during October to November and April to May months. Plant population of same age growing in an open field with a daily average temperature of $38.5/20.0 \pm 2^\circ\text{C}$, relative humidity ($34.7 \pm 5\%$), rate of evapotranspiration (6.18 mm/h) with a natural photoperiod of about 13 h/day length was selected for this study. Healthy fully grown two month old plants were sorted out and the leaves of these plants were subjected to study various leaf parameters like moisture content, succulence index, mesophyll succulence, surface expansion values using standard protocols. Moisture content of leaves was calculated by measuring the difference in weight before and after oven drying for 10 days at 80°C . Succulence was calculated as the ratio of fresh and dry matter, mesophyll succulence and surface expansion (Klug and Ting, 1978).

Study of biochemical and physiological parameters

Diurnal behaviour of titrable acidity status, pH values and stomatal behaviour of the leaves of *C. nudiflora* was studied for successive days using standard protocols. Every three hours fresh leaves were homogenised using glass distilled water and filtrate was used for pH determination using Elico digital Li 120 pH meter. Thomas and Beevers' (1949) method was used to determine titrable acidity for every three hours. Stomatal behaviour was studied every three hours using thin peels of leaf epidermis at $10 \times 45\text{X}$ of compound microscope. Leaf extracts in 85% acetone was estimated for chlorophyll contents at 645 and 663nm using SPECTRONIC 20

spectrophotometer (Arnon, 1949). Standard statistical procedures were followed for analysis of results and values represented means of at least five replicates.

RESULTS

C. nudiflora is a herbaceous annual plant with stem thick, green and creeping while leaves are thick, linear, lanceolate, pubescent with hairy outgrowth and highly succulent. Some characteristic features of the leaves were presented in Table 1. Leaf had an average thickness of 1.68 mm with 85.31% moisture contents, succulence index was 11.02 g.g^{-1} , mesophyll succulence was $5.88 \text{ Kg.g}^{-1} \text{ Chl.}$, with surface expansion of $1.76 \text{ m}^2.\text{Kg}^{-1}$. From the Table 1 it showed that leaf and its succulence parameters were within the range for CAM plants (Klug and Ting, 1978).

Parameters of photosynthetic behaviour are showed in Figure 1. Titrable acidity number (TAN) clearly indicates titrable acidity status in leaf tissues and a typical diurnal fluctuation of about 6 fold increase in TAN values. This was further supported by stomatal behaviour which showed characteristic nocturnal opening of stomata. The CAM

pathway is characterized by the temporal separation of carbon fixation between nocturnal CO₂ fixation by phosphoenolpyruvate carboxylase in the cytosol and daytime decarboxylation of organic acids to release CO₂ which is then refixed by Rubisco in the chloroplast (Ting, 1985). All these findings strongly advocated that CAM is mode of photosynthesis in this plant.

DISCUSSION

In the present study, we noticed that there is nocturnal CO₂ uptake through open stomata at night. Under the dominance of high irradiance and hot temperatures and low availability of water, CAM provides plants to operate photosynthetically with closed stomata (Luttge 2004). Paper chromatography confirmed 4 fold increase in malic acid contents at night. There was also significant diurnal fluctuation in TAN values, pH and osmotic potential in photosynthetic mature leaf tissue. CAM is a taxonomically widespread photosynthetic pathway that has evolved in plants having CO₂ and water-limited environments, including tropical forest canopies with intermittent or seasonal water availability, hot semiarid regions, and some aquatic environments (Katia et al., 2009).

These findings clearly indicated that CAM is also a mode of photosynthesis in this species. In previous studies CAM behaviour was reported in *Thlaspi arvense* of Brassicaceae (Murumkar et al., 1991) and *Aristolochia bracteata* of Aristolochiaceae (Deshmukh and Murumkar, 1996). Thus in present findings, the occurrence of CAM in *C. nudiflora* clearly designated a strong positive function between acidity and CAM mode of photosynthesis in plants growing under extreme drought conditions.

ACKNOWLEDGEMENT

Authors are grateful to authorities of Anekant Education Society, Baramati, Agricultural Development Trust, Baramati, Tuljaram Chaturchand College, Baramati and Shardabai Pawar Mahila Mahavidyalaya, Shardanagar, Baramati for their support and providing facilities during this work.

REFERENCES

- Arnon DI, (1949). Copper enzymes in isolated chloroplasts, Polyphenoloxidase in *Beta vulgaris*. *Plant Physiol.* 24:1-15.
- Bhagat, RB, Shimpale VB, Deshmukh RB (2008). *Flora of Baramati*, ISBN-978-81-8465-009-9, Prakash offset, Pune, India, pp. 1-2.
- Blatter C, Hallberg F (1984). *The flora of Indian Deserts*. Scientific Publishers, Jodhpur, India, pp. 131.
- Deshmukh RB, Murumkar CV (1996). Occurrence of crassulacean acid metabolism in leaves of *Aristolochia bracteata* Ritz. *Acta Societatis Botanicorum Poloniae*, 65(3-4):297-298.
- Kluge M, Ting IP, (1978). *Crassulacean acid metabolism. Analysis of an ecological adaptation*. Springer-Verlag, Berlin-Heidelberg-New York-Tokyo.
- John C Cushman (2001). *Crassulacean Acid Metabolism, A Plastic Photosynthetic Adaptation to Arid Environments*, *Plant Physiol.* 127: 1439-1448.
- Luttge U (2004). *Ecophysiology of Crassulacean acid metabolism (CAM)*, *Ann. Bot. (Lond)* 93: 629-652.
- Luttge U (2012) *Ability of crassulacean acid metabolism plants to overcome interacting stresses in tropical environments*, *AoB Plants*, 2010(0): pp.005
- Murumkar CV, Magdum DK (1991). *Ecophysiological studies in some weeds of Baramati. I. Floristic analysis*, *Biol. Indica.* 2: 73-78.
- Murumkar CV, Magdum DK, Sangar KS, Chavan PD (1991). *CAM-behaviour in a weed, *Thlaspi arvense* L.* *Photosynthetica* 25: 129-131.
- Katia S, Louis S, Santiago, John C. Cushman, Klaus (2009). *Winter Acid Metabolism and Epiphytism Linked to Adaptive Radiations in the Orchidaceae*, *Plant Physiol.*, 149:1838-1847.
- Thomas M, Beevers M (1949) *Physiological studies in acid metabolism in green plants. II Evidence of CO₂ fixation in *Bryophyllum calycinum* and the study of diurnal fluctuations in the genus.* *New Phytol.*, 48:421-447.
- Ting IP (1985). *Crassulacean acid metabolism*, *Ann. Rev. Plant Physiol.* 36:595-622.