

Full Length Research Paper

# Timelike-spacelike Mannheim partner curves in $IR_1^3$

Süleyman ŞENYURT<sup>1\*</sup> and Özcan BEKTAŞ<sup>2</sup>

<sup>1</sup>Department of Mathematics, Faculty of Arts and Sciences, Ordu University, Ordu, Turkey.

<sup>2</sup>Department of Mathematics, Faculty of Arts and Sciences, Yildiz Technical University, Istanbul, Turkey.

Accepted 15 December, 2011

**In this paper, timelike-spacelike Mannheim partner curve couple was defined in Lorentzian space  $IR_1^3$  and the relations were given between the curvatures and torsions of these curves. Furthermore, for a given curve couple, the necessary and sufficient conditions were obtained to become timelike-spacelike Mannheim partner curve couple in  $IR_1^3$ .**

**Key words:** Mannheim curves, Lorentzian space, curvature, torsion.

## INTRODUCTION

As is well-known, a surface is said to be “ruled” if it is generated by moving a straight line continuously in Euclidean space (O’Neill, 1997). Ruled surfaces are one of the simplest objects in geometric modeling. One important fact about ruled surfaces is that they can be generated by straight lines. A practical application of these type of surfaces is that they are used in civil engineering and physics (Guan et al., 1997).

Since building materials, such as wood are straight, they can be considered as straight lines. The result is that if engineers are planning to construct something with curvature, they can use a ruled surface since all the lines are straight (Orbay et al., 2009). In the differential geometry of a regular curve in the Euclidean 3-space  $IE^3$ , it is well-known that one of the important problem is the characterization of a regular curve. The curvature functions  $k_1$  and  $k_2$  of a regular curve play an important role to determine the shape and size of the curve (Kuhnel, 1999; Do Carmo, 1976). For example, If  $k_1 = k_2 = 0$ , the curve is geodesic. If  $k_1 \neq 0$  (constant) and  $k_2 = 0$ , then the curve is a circle with radius  $1/k_1$ . If  $k_1 \neq 0$  (constant) and  $k_2 \neq 0$  (constant), then the curve is a helix in the space.

Another way to the classification and characterization of curves is the relationship between the Frenet vectors of the curves. For example Saint Venant proposed the question whether upon the surfaces generated by the principal normal of a curve, a second curve can exist which has for its principal normal the principal normal of the given curve. This question was answered by Bertrand in 1850; he showed that a necessary and sufficient condition for the existence of such a second curve is that a linear relationship with constant coefficients exists between the first and second curvatures of the given original curve. The pairs of curves of this kind have been called conjugate Bertrand curves, or more commonly Bertrand curves. There are many works related with Bertrand curves in the Euclidean space and Minkowski space. Another kind of associated curves are called Mannheim curve and Mannheim partner curve. If there exists a corresponding relationship between the space curves  $\alpha$  and  $\beta$  such that, at the corresponding points of the curves, principal normal lines of  $\alpha$  coincides with the binormal lines of  $\beta$ , then  $\alpha$  is called a Mannheim curve, and  $\beta$  Mannheim partner curve of  $\alpha$ .

In recent studies, Liu and Wang (2007, 2008) are curious about the Mannheim curves in both Euclidean and Minkowski 3-space and they obtained the necessary and sufficient conditions between the curvature and the torsion for a curve to be the Mannheim partner curves. Meanwhile, the detailed discussion concerned with the Mannheim curves can be found in literature (Wang and

\*Corresponding author. E-mail: [senyurtsuleyman@hotmail.com](mailto:senyurtsuleyman@hotmail.com) or [ssenyurt@odu.edu.tr](mailto:ssenyurt@odu.edu.tr).

Liu, 2007; Liu and Wang, 2008; Orbay and Kasap, 2009; Özkaldi et al., 2009; Azak, 2009) and references therein.

In this paper, we study the timelike-spacelike Mannheim partner curves in Lorentzian space  $IR_1^3$ .

**PRELIMINARY**

The Minkowski 3-space  $IR_1^3$  is the real vector space  $IR^3$  provided with the standart flat metric given by:

$$\langle , \rangle = -dx_1^2 + dx_2^2 + dx_3^2$$

where  $(x_1, x_2, x_3)$  is a standard rectangular coordinate system of  $IR_1^3$ . An arbitrary vector  $\vec{v} = (v_1, v_2, v_3)$  in  $IR_1^3$  can have one of three Lorentzian causal characters; it can be spacelike if  $\langle \vec{v}, \vec{v} \rangle > 0$  or  $\vec{v} = 0$ , timelike if  $\langle \vec{v}, \vec{v} \rangle < 0$  and null (lightlike) if  $\langle \vec{v}, \vec{v} \rangle = 0$  and  $\vec{v} \neq 0$ .

Similarly, an arbitrary curve  $\alpha = \alpha(s)$  can locally be spacelike, timelike or null (lightlike), if all of its velocity vectors  $\alpha'(s)$  are spacelike, timelike or null (lightlike), respectively. We say that a timelike vector is future pointing or past pointing if the first compound of the vector is positive or negative, respectively. The norm of the vector  $\vec{v} = (v_1, v_2, v_3) \in IR_1^3$  is given by:

$$\|\vec{v}\| = \sqrt{|\langle \vec{v}, \vec{v} \rangle|}$$

For any vectors  $\vec{x} = (x_1, x_2, x_3)$  and  $\vec{y} = (y_1, y_2, y_3)$  in  $IR_1^3$  in the meaning of Lorentz vector product of  $\vec{x}$  and  $\vec{y}$  is defined by:

$$\vec{x} \wedge \vec{y} = \begin{vmatrix} e_1 & -e_2 & -e_3 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{vmatrix} = \left( \begin{vmatrix} x_2 & x_3 \\ y_2 & y_3 \end{vmatrix}, \begin{vmatrix} x_1 & x_3 \\ y_1 & y_3 \end{vmatrix}, - \begin{vmatrix} x_1 & x_2 \\ y_1 & y_2 \end{vmatrix} \right)$$

The Lorentzian sphere and hyperbolic sphere of radius  $r$  and center 0 in  $IR_1^3$  are given by:

$$S_1^2 = \{ \vec{x} = (x_1, x_2, x_3) \in IR_1^3 : \langle \vec{x}, \vec{x} \rangle = r^2 \}$$

and

$$H_0^2 = \{ \vec{x} = (x_1, x_2, x_3) \in IR_1^3 : \langle \vec{x}, \vec{x} \rangle = -r^2 \}, \text{ respectively}$$

**Definition 1**

**Hyperbolic angle**

Let  $\vec{a}$  and  $\vec{b}$  be timelike vectors in  $IR_1^3$ . Then, the angle between  $\vec{a}$  and  $\vec{b}$  is defined by  $\langle \vec{a}, \vec{b} \rangle = -\|\vec{a}\| \|\vec{b}\| \cosh \theta$ . The number  $\theta$  is called the hyperbolic angle.

**Central angle**

Let  $\vec{a}$  and  $\vec{b}$  be spacelike vectors in  $IR_1^3$  that span a timelike vector subspace. Then, the angle between  $\vec{a}$  and  $\vec{b}$  is defined by  $\langle \vec{a}, \vec{b} \rangle = \|\vec{a}\| \|\vec{b}\| \cosh \theta$ . The number  $\theta$  is called the central angle.

**Spacelike angle**

Let  $\vec{a}$  and  $\vec{b}$  be spacelike vectors in  $IR_1^3$  that span a spacelike vector subspace. Then, the angle between  $\vec{a}$  and  $\vec{b}$  is defined by  $\langle \vec{a}, \vec{b} \rangle = \|\vec{a}\| \|\vec{b}\| \cos \theta$ . The number  $\theta$  is called the spacelike angle.

**Lorentzian timelike angle**

Let  $\vec{a}$  be a spacelike vector and  $\vec{b}$  be a timelike vector in  $IR_1^3$ . Then, the angle between  $\vec{a}$  and  $\vec{b}$  is defined by  $\langle \vec{a}, \vec{b} \rangle = \|\vec{a}\| \|\vec{b}\| \sinh \theta$ . The number  $\theta$  is called the Lorentzian timelike angle.

Let  $\{t(s), n(s), b(s)\}$  be the moving Frenet frame along the curve  $\alpha(s)$ . Then  $t(s), n(s)$  and  $b(s)$  are tangent, the principal normal and the binormal vector of the curve  $\alpha(s)$ , respectively. Depending on the casual character of the curve  $\alpha$ , we have the following dual Frenet-Serret formulas. If  $\alpha$  is a timelike curve;

$$\begin{pmatrix} t' \\ n' \\ b' \end{pmatrix} = \begin{pmatrix} 0 & k_1 & 0 \\ k_1 & 0 & k_2 \\ 0 & -k_2 & 0 \end{pmatrix} \begin{pmatrix} t \\ n \\ b \end{pmatrix} \tag{1}$$

where

$$\langle t, t \rangle = -1, \langle n, n \rangle = \langle b, b \rangle = 1, \langle t, n \rangle = \langle n, b \rangle = \langle t, b \rangle = 0.$$

We denote by  $\{v_1(s), v_2(s), v_3(s)\}$  the moving Frenet frame along the curve  $\beta(s)$ . Then,  $v_1(s), v_2(s)$  and  $v_3(s)$  are tangent, the principal normal and the binormal vector of the curve  $\beta(s)$ , respectively. Depending on the casual character of the curve  $\beta$ , we have the following Frenet-Serret formulas. If  $\beta$  is a spacelike curve with a timelike binormal  $v_3$ ;

$$\begin{pmatrix} v_1' \\ v_2' \\ v_3' \end{pmatrix} = \begin{pmatrix} 0 & p & 0 \\ -p & 0 & q \\ 0 & q & 0 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} \tag{2}$$

where

$$\langle v_3, v_3 \rangle = -1, \langle v_1, v_1 \rangle = \langle v_2, v_2 \rangle = 1, \langle v_1, v_2 \rangle = \langle v_2, v_3 \rangle = \langle v_1, v_3 \rangle = 0.$$

If the curves are unit speed curve, then curvature and torsion are calculated by:

$$\begin{cases} k_1 = \|t'\|, \\ k_2 = \langle n', b \rangle, \\ p = \|v_1'\|, \\ q = \langle v_2', v_3 \rangle. \end{cases} \tag{3}$$

If the curves are not unit speed curve, then curvature and torsion are calculated by:

$$\begin{cases} k_1 = \frac{\|\alpha' \wedge \alpha''\|}{\|\alpha'\|^3}, \\ k_2 = \frac{\det(\alpha', \alpha'', \alpha''')}{\|\alpha' \wedge \alpha''\|^2}, \\ p = \frac{\|\beta' \wedge \beta''\|}{\|\beta'\|^3}, \\ q = \frac{\det(\beta', \beta'', \beta''')}{\|\beta' \wedge \beta''\|^2}. \end{cases} \tag{4}$$

**TIMELIKE-SPACELIKE MANNHEIM PARTNER CURVE IN  $IR_1^3$**

Here, we define timelike-spacelike Mannheim partner curves in  $IR_1^3$  and we give some characterization for

timelike-spacelike Mannheim partner curves in the same space. Using these relationships, we will comment on Shell's and Mannheim's theorems again.

**Definition 2**

Let  $\alpha: I \rightarrow IR_1^3$  be a timelike curve and  $\beta: I \rightarrow IR_1^3$  be spacelike with timelike binormal. If there exists a corresponding relationship between the timelike curve  $\alpha$  and the spacelike curve with dual timelike binormal  $\beta$  such that, at the corresponding points of the curves, the binormal lines of  $\alpha$  coincides with the principal normal lines of  $\beta$ , then  $\alpha$  is called a timelike Mannheim curve, and  $\beta$  is called a Mannheim partner curve of  $\alpha$ . The pair  $\{\alpha, \beta\}$  is said to be timelike-spacelike Mannheim pair. Let  $\{t, n, b\}$  be the Frenet frame field along  $\alpha = \alpha(s)$  and let  $\{v_1, v_2, v_3\}$  be the Frenet frame field along  $\beta = \beta(s)$ . On the other way  $\theta$  is angle between  $t$  and  $v_1$ , there is a following equation between the Frenet vectors and their derivative;

$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} \sinh \theta & \cosh \theta & 0 \\ 0 & 0 & 1 \\ \cosh \theta & \sinh \theta & 0 \end{pmatrix} \begin{pmatrix} t \\ n \\ b \end{pmatrix} \tag{5}$$

**Theorem 1**

The distance between corresponding points of the timelike-spacelike Mannheim partner curves in  $IR_1^3$  is constant.

**Proof**

From the definition of spacelike Mannheim curve, we can write:

$$\tilde{\beta}(s^*) = \tilde{\alpha}(s) + \lambda(s)B(s). \tag{6}$$

By taking the derivate of this equation with respect to  $s$  and applying the Frenet formulas, we get:

$$v_1 \frac{ds^*}{ds} = t - \lambda k_2 n + \lambda' b \tag{7}$$

where the superscript  $(^*)$  denotes the derivate with respect to the arc length parameters of the dual curve

$\alpha(s)$ . Since the vectors  $b$  and  $v_2$  are linear, we get :

$$\left\langle v_1 \frac{ds^*}{ds}, b \right\rangle = \langle t, b \rangle - \lambda k_2 \langle n, b \rangle + \lambda' \langle b, b \rangle, \quad \lambda' = 0.$$

Then, we get  $\lambda = c$ . On the other hand, from the definition of distance function between  $\alpha(s)$  and  $\beta(s)$  we can write:

$$d(\alpha(s), \beta(s)) = \|\lambda(s)B(s)\| = |\lambda|.$$

This completes the proof.

**Theorem 2**

For a timelike-spacelike curve  $\alpha$  in  $IR_1^3$ , there is a spacelike curve  $\beta$  so that  $\{\alpha, \beta\}$  is a spacelike Mannheim pair.

**Proof**

Since the vectors  $v_2$  and  $b$  are linearly dependent, the Equation 6 can be written as:

$$\alpha = \beta - \lambda v_2. \tag{8}$$

Since  $\lambda$  is a nonzero constant, there is a timelike curve  $\beta$  for all values of  $\lambda$ .

Now, we can give the following theorem related to curvature and torsion of the timelike-spacelike Mannheim partner curves.

**Theorem 3**

Let  $\{\alpha, \beta\}$  be a timelike-spacelike Mannheim pair in  $IR_1^3$ . If  $k_2$  is dual torsion of  $\alpha$  and  $p$  is dual curvature and  $q$  is dual torsion of  $\beta$ , then:

$$k_2 = -\frac{p}{\lambda q}. \tag{9}$$

**Proof**

By taking the derivate of Equation 7 with respect to  $s$  and applying the Frenet formulas, we obtain:

$$v_1 \frac{ds^*}{ds} = t - \lambda k_2 n. \tag{10}$$

Let  $\theta$  be dual angle between the dual tangent vectors  $t$  and  $v_1$ , we can write:

$$\begin{cases} v_1 = \sinh \theta t + \cosh \theta n \\ v_3 = \cosh \theta t + \sinh \theta n. \end{cases} \tag{11}$$

From Equations 10 and 11, we get:

$$\frac{ds^*}{ds} = \frac{1}{\sinh \theta}, \quad -\lambda k_2 = \cosh \theta \frac{ds^*}{ds}. \tag{12}$$

By taking the derivate of Equation 8 with respect to  $s$  and applying the Frenet formulas, we obtain:

$$k_2 = (1 + \lambda p) v_1 \frac{ds^*}{ds} - \lambda q v_3 \frac{ds^*}{ds}. \tag{13}$$

From Equation 11, we can write:

$$\begin{cases} t = -\sinh \theta v_1 + \cosh \theta v_3 \\ n = \cosh \theta v_1 - \sinh \theta v_3, \end{cases} \tag{14}$$

where  $\theta$  is the angle between  $t$  and  $v_1$  at the corresponding points of the curves of  $\alpha$  and  $\beta$ . By taking into consideration Equations 13 and 14, we get:

$$\sinh \theta = -(1 + \lambda p) \frac{ds^*}{ds}, \quad \cosh \theta = -\lambda q \frac{ds^*}{ds}. \tag{15}$$

Substituting  $\frac{ds^*}{ds}$  into Equation 15, we get:

$$\sinh^2 \theta = -(1 + \lambda p), \quad \cosh^2 \theta = \lambda^2 k_2 q. \tag{16}$$

From the Equation 16, we can write:

$$k_2 = -\frac{p}{\lambda q}.$$

**Corollary 1**

Let  $\{\alpha, \beta\}$  be a timelike-spacelike Mannheim pair in  $IR_1^3$ . Then, the product of torsions  $k_2$  and  $q$  at the

corresponding points of the spacelike Mannheim partner curves are not constant.

Namely, Schell's theorem is invalid for the timelike-spacelike Mannheim curves. By considering Theorem 3, we can give the following results.

**Corollary 2**

Let  $\{\alpha, \beta\}$  be a timelike-spacelike Mannheim pair in  $IR_1^3$ . Then, torsions  $k_2$  and  $q$  has a negative sign.

**Theorem 4**

Let  $\{\alpha, \beta\}$  be a timelike-spacelike Mannheim pair in  $IR_1^3$ . Between the curvature and the torsion of the spacelike curve  $\beta$ , there is the relationship:

$$\mu q - \lambda p = 1 \tag{17}$$

where  $\mu$  and  $\lambda$  are nonzero dual numbers.

**Proof**

From Equation 15, we obtain:

$$\frac{\sinh \theta}{1 + \lambda p} = \frac{\cosh \theta}{\lambda q} \tag{18}$$

Arranging this equation, we get:

$$\tanh \theta = \frac{1 + \lambda p}{\lambda q}$$

and if we choose  $\mu = \lambda \tanh \theta$  for brevity, we will see that:  $\mu q - \lambda p = 1$ .

**Theorem 5**

Let  $\{\alpha, \beta\}$  be a timelike-spacelike Mannheim pair in  $IR_1^3$ . The following equations are for the curvatures and the torsions of the curves  $\alpha$  and  $\beta$

$$1. k_1 = -\frac{d\theta}{ds},$$

$$2. k_2 = p \cosh \theta \frac{ds^*}{ds} - q \sinh \theta \frac{ds^*}{ds},$$

$$3. p = k_2 \cosh \theta \frac{ds}{ds^*},$$

$$4. q = k_2 \sinh \theta \frac{ds}{ds^*}.$$

**Proof**

1. By considering Equation 11, we can easily show that  $\langle t, v_1 \rangle = \sinh \theta$ . Differentiating this equality with respect to  $s$  by considering Equation 1, we have:

$$\langle t', v_1 \rangle + \langle t, v_1' \rangle = -\sinh \theta \frac{d\theta}{ds},$$

From Equations 1 and 2, we can write:

$$\langle k_1 n, v_1 \rangle + \langle t, p v_2 \frac{ds^*}{ds} \rangle = -\sinh \theta \frac{d\theta}{ds}$$

From Equation 14, we get:

$$k_1 = -\frac{d\theta}{ds}.$$

2. By considering Equation 11, we can easily show that  $\langle n, v_2 \rangle = 0$ . Differentiating this equality with respect to  $s$  and by considering Equation 1, we have:

$$\langle n', v_2 \rangle + \langle n, v_2' \frac{ds^*}{ds} \rangle = 0,$$

From Equations 1 and 2, we can write:

$$\langle k_1 t + k_2 b, v_2 \rangle + \langle \cosh \theta v_1 - \sinh \theta v_3, (-p v_1 + q v_3) \frac{ds^*}{ds} \rangle = 0,$$

From Equations 14, we get:

$$k_2 = p \cosh \theta \frac{ds^*}{ds} - q \sinh \theta \frac{ds^*}{ds}.$$

3. By considering Equation 14, we can easily show that  $\langle b, v_1 \rangle = 0$ . Differentiating this equality with respect to  $s$  and by considering Equation 1, we have:

$$\langle b', v_1 \rangle + \left\langle b, v_1' \frac{ds^*}{ds} \right\rangle = 0,$$

From Equations 1, 2 and 14, we can write:

$$\langle -k_2 (\cosh \theta v_1 - \sinh \theta v_3), v_1 \rangle + \left\langle b, p v_2 \frac{ds^*}{ds} \right\rangle = 0,$$

$$p = k_2 \cosh \theta \frac{ds}{ds^*}.$$

4. By considering Equation 11, we can easily show that  $\langle b, v_3 \rangle = 0$ . Differentiating this equality with respect to  $s$  and by considering Equation 1, we have:

$$\langle b', v_3 \rangle + \left\langle b, v_3' \frac{ds^*}{ds} \right\rangle = 0.$$

From Equations 1, 2 and 14, we can write:

$$\langle -k_2 (\cosh \theta v_1 - \sinh \theta v_3), v_3 \rangle + \left\langle b, q v_2 \frac{ds^*}{ds} \right\rangle = 0,$$

$$q = k_2 \sinh \theta \frac{ds}{ds^*}.$$

By considering statements 3 and 4 of Theorem 5, we can give the following results.

**Corollary 3**

Let  $\{\alpha, \beta\}$  be a timelike-spacelike Mannheim pair in  $IR_1^3$ . Then, there exist the following relation between curvature and torsion of  $\beta$  and torsion of  $\alpha$ ;

$$p^2 - q^2 = k_2^2 \left( \frac{ds}{ds^*} \right)^2. \tag{19}$$

**Theorem 6**

A timelike space curve in  $IR_1^3$  is a timelike-spacelike Mannheim curve, if and only if its curvature  $p$  and torsion  $q$  satisfy the formula:

$$\lambda(q^2 - p^2) = p \tag{20}$$

where  $\lambda$  is never a pure dual constant.

**Proof**

By taking the derivate of the statement  $\alpha = \beta - \lambda V_2$  with respect to  $s$  and applying the Frenet formulas we obtain:

$$t \frac{ds}{ds^*} = v_1 + \lambda (p v_1 - q v_3),$$

$$k_1 n \left( \frac{ds}{ds^*} \right)^2 + t \frac{d^2 s}{ds^{*2}} = p v_2 + \lambda (p' v_1 - q' v_3 + (p^2 - q^2) v_2).$$

Taking the inner product, which is the last equation with  $b$ , we get:

$$\lambda(q^2 - p^2) = p.$$

**Theorem 7**

Let  $\{\alpha, \beta\}$  be a timelike-spacelike Mannheim partner curves in  $IR_1^3$ . Moreover, the points  $\alpha(s)$  and  $\beta(s)$  be two corresponding points of  $\{\alpha, \beta\}$  and  $M$  and  $M^*$  be the curvature centers at these points, respectively. Then, the ratio:

$$\frac{\|\beta(s)M\|}{\|\alpha(s)M\|} : \frac{\|\beta(s)M^*\|}{\|\alpha(s)M^*\|} = (1 + k_1 p)(1 + \lambda p) \neq \text{constant}. \tag{21}$$

**Proof**

A circle that lies in the osculating plane of the point  $\alpha(s)$  on the timelike curve  $\alpha$  and that has the centre

$$M = \alpha(s) + \frac{1}{k_1} n$$

lying on the principal normal  $n$  of the

point  $\alpha(s)$  and the radius  $\frac{1}{k_1}$  far from  $\alpha(s)$ , is called

osculating circle of the curve  $\alpha$  in the point  $\alpha(s)$ .

Similar definition can be given for curve  $\beta$  too.

Then, we can write:

$$\|\alpha(s)M\| = \left\| \frac{1}{k_1} n \right\| = \frac{1}{k_1},$$

$$\|\alpha(s)M^*\| = \left\| \lambda b + \frac{1}{p}v_2 \right\| = \frac{1}{p} + \lambda,$$

$$\|\beta(s)M^*\| = \left\| \frac{1}{p}v_2 \right\| = \frac{1}{p},$$

$$\|\beta(s)M\| = \left\| \lambda v_3 + \frac{1}{k_1}n \right\| = \frac{1}{k_1} + \lambda$$

Therefore, we obtain:

$$\frac{\|\beta(s)M\| \cdot \|\beta(s)M^*\|}{\|\alpha(s)M\| \cdot \|\alpha(s)M^*\|} = (1 + \lambda p)\sqrt{1 - \lambda^2 k_1^2} \neq \text{constant}.$$

Thus, we can give the following.

**Corollary 4**

Mannheim’s theorem is invalid for the timelike-spacelike Mannheim partner curve  $\{\alpha, \beta\}$  in  $IR_1^3$ .

**REFERENCES**

Azak AZ (2009). On Timelike Mannheim Partner Curves in  $L^3$  Sakarya University Faculty of Arts and Science, The Journal of Arts and Science, (11): 35-45.  
 Do Carmo MP (1976). Differential Geometry of Curves and Surfaces, Pearson Education. New York: Academic Press.  
 Guan Z, Guan, Ling J, Ping X, Rongxi T (1997). Study and Application of Physics-Based Deformable Curves and Surfaces. Computers and Graphics, 21: 305-313.

Kahraman T, Önder M, Kazaz M, Uğurlu HH (2011). Some Characterizations of Mannheim Partner Curves in the Minkowski 3-space  $E_1^3$ . Proceedings of the Estonian Academy of Sciences, 60, 4, 210–220.  
 Kuhnel W (1999). Differential Geometry: Curves-Surfaces-Manifolds. Braunschweig, Wiesbaden.  
 Liu H, Wang F (2007). Mannheim Partner Curves in 3-Space. Proceedings of The Eleventh Int. Workshop on Diff. Geom., 11: 25-31  
 Liu H, Wang F (2008). Mannheim Partner Curves in 3-Space, J. Geom., 88(1-2): 120-126.  
 Orbay K, Kasap E (2009). On Mannheim Partner Curves in  $E^3$ . Int. J. Phy. Sci., 4(5): 261-264.  
 Orbay K, Kasap E, Aydemir İ (2009). Mannheim Offsets of Ruled Surfaces. Mathematical Problems in Engineering. Article Number: 160917.  
 O'Neill B (1997). Elementary Differential Geometry, 2nd ed. Academic Press, New York.  
 Özkaldi S, Ilarslan K, Yaylı Y (2009). On Mannheim Partner Curves In Dual Space. Analele Stiintifice ale Universitatii Ovidius Constanta. Seria Matematica. XVII, fasc. 2.