

Full Length Research Paper

Intestinal helminth infection and anaemia during pregnancy: A community based study in Ghana

S. E. Baidoo^{1, 2}, S. C. K. Tay¹, K. Obiri-Danso³ and H. H. Abruquah^{4*}

¹Department of Clinical Microbiology, School of Medical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

²Department of Laboratory Technology, University of Cape Coast, Cape Coast, Ghana.

³Department of Theoretical and Applied Biology, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

⁴Division of Microbiology and Infectious Diseases, University Hospital, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

Accepted 2 February, 2010

Intestinal helminths are among the most common and widespread of human infections, contributing to poor nutritional status, anaemia and impaired growth. Anaemia and iron deficiency in pregnancy is a major public health problem in developing countries, but their causes are not always known. The objective of this study was to assess the prevalence and severity of anaemia and iron deficiency and their association with helminths, among pregnant women in the Ashanti region of Ghana. A cohort study was carried out in the Sekyere, East district of the Ashanti region of Ghana. One hundred and eight (108) pregnant women were followed until 5-10 weeks postpartum, during the period of (December 2005 - November 2006). Haemoglobin and total serum iron concentrations were evaluated in venous blood samples, and helminths infections evaluated in stool samples in each trimester using standard methods. Most of the 108 pregnant women, 54.9% were found to be anaemic. The highest prevalence of anaemia and low iron stores (57.4 and 32.4%, respectively) were found in the second trimester. Only 17.6% had evidence of helminths infection, with *Necator americanus* (hookworm) being the commonest (13.9%). There was a significant association between hookworm infection and low iron stores. The study concluded that hookworm infection is a strong predictor of iron status. These findings reinforce the need to provide antihelminthic therapy to infected women before conception as a public health strategy in reducing the prevalence of hookworm infection, and in addition to providing nutritional and iron supplements to effectively control anaemia in pregnancy.

Key words: Intestinal helminth, anaemia, paragravids, primigravids.

INTRODUCTION

Intestinal helminthes are among the most common and widespread of human infections. They contribute to poor nutritional status, anaemia and impaired growth in children of school going age (Dickson et al., 2000). Epidemiological surveys have revealed that, poor sanitation and inappropriate environmental conditions

coupled with indiscriminate defaecation, geophagy, and contamination of water bodies are the most important predisposing factors to intestinal worm infestation (Brooker et al., 2008). The prevalence and intensity of infection is especially high in developing countries, particularly among populations with poor environmental sanitation (van Eijk et al., 2009). Other practices such as hand washing, disposal of refuse, personal hygiene, wearing of shoes and others, when not done properly may contribute to the infection or picking of these worms from the environment (Stoltzfus et al., 1997).

*Corresponding author. E-mail: kowabruquah@yahoo.co.uk.
Tel: 051-60320.

Anaemia denotes the complex of signs and symptoms associated with low haemoglobin levels. Globally, the most common cause of anaemia is believed to be iron deficiency due to inadequate dietary iron intake, physiologic demands of pregnancy and rapid growth and iron losses due to parasitic infections (Ayoya et al., 2006; Dreyfuss et al., 2000). However, iron deficiency is not the only cause of anaemia. Other prevalent causes of anaemia include malaria, chronic infections and nutritional deficiencies of vitamin A, folate and vitamin B-12 (Dreyfuss et al., 2000). The relative contributions of these causes of anaemia and iron deficiency varies by sex, age and population and are not well described in many populations (Dreyfuss et al., 2000).

Iron deficiency is responsible for 95% of cases of anaemia during pregnancy (Porter and Kaplan, 1999) which is often due to inadequate dietary intake (especially in teenage girls), to a previous pregnancy or to the normal loss of iron in menstruation (Porter and Kaplan, 1999).

Impact of intestinal helminth infections on anaemia during pregnancy is aggravated by low nutritional status of subjects whose staple foods, such as rice, cassava, and maize are poor sources of folate, and iron (Ayoya et al., 2006; Pasricha et al., 2008). The most important cause of pathological chronic loss of blood and iron in the tropics is hookworm and other soil-transmitted helminthes (Brooker et al., 2008), and malaria in pregnancy (Fleming, 1982). At a hospital in Kathmandu, Nepal, hookworm infection was associated with severe but not moderate anaemia among women receiving antenatal care (Bondevik et al., 2000). Data on the epidemiology of iron deficiency, anaemia in East Africa and elsewhere point to the important contribution of hookworms to this condition (Ayoya et al., 2006; Stoltzfus et al., 1997). Hookworm infection has been established as a strong predictor of iron deficiency and anaemia in other populations (Ayoya et al., 2006; Stoltzfus et al., 1997), but few studies have examined their relationships in pregnant women (WHO, 1993).

The occurrence of helminth infection at high rates among pregnant women is mostly indicative of faecal pollution of soil and domestic water supply around homes due to poor sanitation and improper sewage disposal (Bundy et al., 1995; van Eijk et al., 2009). Pregnant women are at high risk of infection because of their close relationship with children (Bundy et al., 1995; van Eijk et al., 2009). Most of these worms are transmitted through the soil whilst, the practice of soil-eating (geophagy) is common amongst pregnant women in many communities in developing countries (Brooker et al., 2008). It is in this vain that, this study investigates the prevalence and some clinical effects of anaemia in pregnant women infected with intestinal helminthes in the Sekyere East district of the Ashanti region of Ghana. This would help largely in the management of intestinal worm burden interfering with iron stores in pregnant mothers.

MATERIALS AND METHODS

Setting

Subjects were recruited from the Sekyere East district of the Ashanti region with the Effiduase hospital at the base laboratory. The examination center was centrally located in the study area and represented the general characteristics of the district. This analysis includes data collected from December, 2005 - November, 2006 from pregnant women attending antenatal clinics in the clinic sub study area.

Subject selection

The study population involved pregnant women of ages between 12 - 45 years who participated in a randomized community intervention. Three hundred eligible pregnant women who visited the clinic had pregnancies confirmed with a β -human chorionic gonadotropin urine test. Persons were excluded from the study if the initial ANC screening detected that they had any haemoglobinopathy such as G6PD or sickle cell diseases.

Sample collection and analysis

Blood was collected via venipuncture. Anaemia and Iron status were assessed with haemoglobin (Hb) and total serum iron (TSI) or ferritin concentrations. Morphological study of red cells was carried out as a confirmatory test for iron deficiency anaemia (Leishman staining). Hb was measured with a Haemocue hemoglobinometer (Mission Viejo, CA). Serum ferritin (TSI) was assessed with a fluorometric immunoassay (Delfia System; Wallac, Gaithersburg, MD). The formol-ether concentration method was used in the preparation of stool samples for microscopy and detection of helminthes.

Data analysis

Firstly, estimates of the severity of anaemia, defined by Hb concentration, and of iron deficiency, defined by TSI or ferritin concentration were examined as a cause of anaemia. Anaemia was defined as Hb <10 g/dl and differentiated as Hb <10 g/dl - 'severe anaemia'; Hb <11 - 11.5 g/dL - 'moderate anaemia' and Hb 12 - 15.2 g/dL - 'normal'. Persons with TSI or serum ferritin <60 μ g/dL were also described as 'anaemic' (Porter and Kaplan, 1999). In the morphological examination of iron deficiency anaemia, the morphology of the red cells should be microcytic, hypochromic with anisocytosis and poikilocytosis (Porter and Kaplan, 1999). Gestation age and gravidity were also retained in all models because iron status and haemoglobin concentration is known to be strongly associated with gestation age and gravidity (Porter and Kaplan, 1999). Data was analyzed using SPSS (version 11.0; SPSS Inc, Chicago), while statistical significance was defined as a p-value of <0.05. The Pearson Chi - square test was used to examine the association between both total serum iron and haemoglobin concentrations and helminth infection as indicators of anaemia.

RESULTS

Most of the 108 women recruited into the study, 66 (61.1%) were paragravids and 42 (39.9%) primigravids. Clinical data obtained within the first month of recruitment show that 27.8% of these women were in their first

Table 1. Total serum iron (TSI) concentration of study subjects as distributed over the trimesters of pregnancy. Using TSI 60 persons were categorized as being anaemic as compared with 59 using Hb levels

Gestation	Serum iron concentration		Total
	< 60 ug/dl (Anaemic, %)	> 60 ug/dl (Normal)	
1 st Trimester	17 (15.7)	13	30
2 nd Trimester	35 (32.4)	27	62
3 rd Trimester	8 (7.4)	8	16
Total	60 (55.6)	48	108

trimester, while 57.4 and 14.8% were in their second and third trimesters respectively. Of the paragravids 15 (13.8%), 39 (36.1%), and 12 (11.1%) were in their first, second and third trimester, respectively; and 15 (13.8%), 23 (21.3%) and 4 (3.7%) of primigravids were also in their first, second and third trimesters, respectively.

Prevalence of anaemia and iron deficiency

Anaemia was present in more than 50% of the pregnant women studied (Table 1). Iron deficiency appeared to be the dominant cause of anaemia, especially, moderate to severe anaemia in these communities with 60 subjects having TSI concentrations of <60 µg/dl while 59 persons had Hb levels < 12 g/dl (55.6 and 54.6%, respectively, Figure 1).

Helminth infection

Most of the 108 pregnant women, only 19 or 17.6% were infected with helminthes; the commonest being *Necator americanus* (hookworm) infection (15 or 13.9%) followed by *Strongloides stercoralis* (2 or 1.9%), *Ascaris lumbricoides* (1 or 0.9%) and *T. trichiura* (1 or 0.9%).

All 19 women infected with at least one species of pathogenic intestinal helminthes were found to be moderate to severe anaemia. All 15 pregnant women who were infected with hookworm were found to be anaemic (Hb <10.0 g/dL and/or TSI < 20 µg/dL), with 13 of them being severely anaemic (Figure 2). Hookworm infection had a strong association with all two indicators of anaemia and iron deficiency ($p= 0.001$ and 0.00 for TSI and Hb respectively).

DISCUSSION

In developing countries, most pregnant women generally become anaemic and this is presumed to be primarily, as a result of iron deficiency (WHO, 1996). However, the definition and identification of iron deficiency has been a problem especially in situations where there are multifactorial causes of anaemia. Nevertheless,

hookworm has been established as a major predictor for iron deficiency anaemia (van den Broek, 1996; van den Broek et al., 1998). It is also known that poverty, ignorance, geophagy, promiscuous defaecation and poor personal hygiene and environmental sanitation also predispose humans to hookworm infection (Larry and Janovy Jnr, 1996).

The 108 pregnant women studied, 55.6% of them were classified as anaemic, which although is on the high side, it is not uncharacteristic of the community setting of which the study was undertaken (Jans et al., 2008; Pasricha et al., 2008). In a recent study in Netherlands, the prevalence of anaemia in pregnancy was found to be 3.4% (Jans et al., 2008). Despite the discrepancies in the number of study subjects and the study areas of the two studies, ethnicity played a major role in defining the risk of being anaemic in pregnancy, with women of non-northern European descent, there are more at risk than their northern European counterparts (Jans et al., 2008). In similar studies in India and Nepal, the percentages of pregnant women who were anaemic were 88 and 81, respectively, (Agarwal et al., 1987; WHO, 1998). This affirms the predominance of predisposing factors of anaemia within rural settings of developing countries. In our part of the world, anaemia may also be caused by malaria, HIV and poor nutrition (Ayoya et al., 2006; Muhangi et al., 2007).

Iron deficiency appeared to be the dominant cause of anaemia in these communities as defined by TSI concentration and Hb levels (55.6 and 54.6%, respectively) and although only 19 of the 60 anaemic women (31.7%) had helminthes infection, iron deficiency was found to be significantly associated with hookworm infection ($p = 0.001$).

It was encouraging to note that, only 17.6% were infected with helminths in this study, the commonest hookworm (13.9%), compared with a similar study in western Kenya (van Eijk et al., 2009). However, since this study is an epidemiological one and the subjects were asymptomatic, any helminth ova or larvae present would be in very low intensity and possibly undetectable (Ayoya et al., 2006; Bundy et al., 1995; van Eijk et al., 2009).

Hookworm infection has been established as a strong predictor of iron deficiency and anaemia in other populations (Bundy et al., 1995; Hopkins et al., 1997;




Figure 1. Different species of helminthes identified among the pregnant women studied with *Necator americanus* (hookworm) being the most common.




Figure 2. Haemoglobin and TSI levels in 15 pregnant women who had hookworm infection. All 15 were found to have Hb levels and TSI concentrations below normal, with 13 of them being severely anaemic.

Stoltzfus et al., 1997), and few studies have examined these relationships in pregnant women. In this study, hookworm infection was associated with severe anaemia (95% CI 0.2-0.8) which conformed to published data (Ayoya et al., 2006; Bondevik et al., 2000; Brooker et al., 2008; van Eijk et al., 2009).

A single course of anthelminthic therapy in addition to iron-folate supplementation significantly increased hemoglobin concentrations and improved iron status (serum ferritin and EP) in pregnant Sri Lankan plantation workers, suggesting that hookworm infection caused iron deficiency anaemia in that population (Atukorala et al., 1994). However, allocation to anthelminthic therapy was nonrandom and the prevalence and intensity of hookworm infection were not assessed.

The pathogenicity of hookworm infection shows that the disease manifests in three main phases, with the intestinal phase representing the most important period.

A moderate hookworm infection according to studies, will gradually produce iron deficiency anaemia as the body reserves of iron are used up, with the severity depending on the worm load and the dietary intake of iron (Larry and Janovy Jnr, 1996).

In sub-Saharan Africa, up to 24 million women may become pregnant each year (Bundy et al., 1995). This high rate is often related to an increase in the susceptibility of pregnant women to infections because pregnancy is a time of high hormone activity which may exert immuno-suppressive effects on the child bearing woman (Beer and Billingham, 1978; McGregor et al., 1983). Information also shows that, iron deficiency is responsible for about 95% cases of anaemia during pregnancy usually due to inadequate dietary intake (especially in teenage girls), to previous pregnancies or to normal loss of iron in blood with menses to the interference of iron stores by parasites (Porter and

Kaplan, 1999).

Findings in this study provide a population based picture of iron status during pregnancy among women in the Sekyere East district, who live in conditions of poverty, malnutrition and endemic infections. After evaluating all the commonly available variables for measuring iron status, van der Broek and Letsky, 1998 estimated TSI (ferritin) as the best predictor of iron deficiency (van den Broek et al., 1998).

Given that women in these communities are low income generating people, with majority of study subjects between the ages of 20 - 29, malnutrition is a possibility which could account for the high endemicity of anaemia.

Conclusion

The study concludes that anaemia is prevalent within rural communities of the Sekyere East district of the Ashanti region and that hookworm infection is a strong predictor of iron status.

Recommendation

Anthelmintic therapy is inexpensive and safe during pregnancy after the first trimester, therefore, it should be part of the antenatal programme since malaria diagnosis and treatment is also part of the antenatal programme (Larocque and Gyorkos, 2006). The world health organization recommends anthelmintic therapy for women to control hookworm infection in areas, in which the prevalence of infection is high and anaemia is prevalent (WHO, 1996).

Anthelmintic therapy and improvement in the amount and quality of dietary iron intake among pregnant women should be encouraged as an important long-term goal, and if well established, it can meet the iron needs of pregnant women with hookworm and other intestinal helminth infections, particularly among populations who consume diets of low iron bioavailability.

Also, other causes of anaemia should be looked at in the Sekyere East district to arrive at a comprehensive approach in solving the problem of anaemia.

REFERENCES

- Agarwal DK, Agarwal KN, Tripathi AM (1987). Nutritional status in rural pregnant women of Bihar and Uttar Pradesh. Indian Pediatr. 24: 119-125.
- Atukorala TM, Silva LD, Dechering WH, Dassenaeke TS, Perera RS (1994). Evaluation of effectiveness of iron-folate supplementation and anthelmintic therapy against anemia in pregnancy--a study in the plantation sector of Sri Lanka. Am. J. Clin. Nutr. 60: 286-292.
- Ayoya MA, Spiekermann-Brouwer GM, Traore AK, Stoltzfus RJ, Garza C (2006). Determinants of anemia among pregnant women in Mali. Food Nutr. Bull. 27: 3-11.
- Beer AE, Billingham RE (1978). Maternal immunological recognition mechanisms during pregnancy. Ciba Found Symp. pp. 293-322.
- Bondevik GT, Eskeland B, Ulvik RJ, Ulstein M, Lie RT, Schneede J, Kvale G (2000). Anaemia in pregnancy: possible causes and risk factors in Nepali women. Eur. J. Clin. Nutr. 54: 3-8.
- Brooker S, Hotez PJ, Bundy DA (2008). Hookworm-Related Anaemia among Pregnant Women: A Systematic Review. PLoS Negl Trop. Dis. 2: e291.
- Bundy DA, Chan MS, Savioli L (1995). Hookworm infection in pregnancy. Trans. R Soc. Trop. Med. Hyg. 89: 521-522.
- Dickson R, Awasthi S, Demelweek C, Williamson P (2000). Anthelmintic drugs for treating worms in children: effects on growth and cognitive performance. Cochrane Database Syst. Rev. CD000371.
- Dreyfuss ML, Stoltzfus RJ, Shrestha JB (2000). Hookworms, malaria and vitamin A deficiency contribute to anemia and iron deficiency among pregnant women in the plains of Nepal. J. Nutr. 130: 2527-2536.
- Fleming AF (1982). Iron deficiency in the tropics. Clin. Haematol. 11: 365-388.
- Hopkins RM, Gracey MS, Hobbs RP, Spargo RM, Yates M, Thompson RC (1997). The prevalence of hookworm infection, iron deficiency and anaemia in an aboriginal community in North-west Australia. Med. J. Aust. 166: 241-244.
- Jans SM, Daemers DO, Vos R, Lagro-Jansen AL (2008). Are pregnant women of non-Northern European descent more anaemic than women of Northern European descent? A study into the prevalence of anaemia in pregnant women in Amsterdam. Midwifery.
- Larocque R, Gyorkos TW (2006). Should deworming be included in antenatal packages in hookworm-endemic areas of developing countries? Can. J. Public Health 97: 222-224.
- Larry SR, Janovy Jnr J (1996). Introduction to Parasitology. In Foundations Parasitology, pp. 1 - 7, 410-415.: The McGraw Hill Companies, Inc.
- McGregor IA, Wilson ME, Billewicz WZ (1983). Malaria infection of the placenta in The Gambia, West Africa; its incidence and relationship to stillbirth, birthweight and placental weight. Trans. R Soc. Trop. Med. Hyg. 77: 232-244.
- Muhangi L, Woodburn P, Omara M (2007). Associations between mild-to-moderate anaemia in pregnancy and helminth, malaria and HIV infection in Entebbe, Uganda. Trans. R Soc. Trop. Med. Hyg. 101: 899-907.
- Pasricha SR, Caruana SR, Phuc TQ (2008). Anemia, iron deficiency, meat consumption, and hookworm infection in women of reproductive age in Northwest Vietnam. Am. J. Trop. Med. Hyg. 78: 375-381.
- Porter RS, Kaplan JL (1999). The Merck Manual of diagnosis and therapy, 17th Ed edn. White house station, N.J.
- Stoltzfus RJ, Chwya HM, Tielsch JM, Schulze KJ, Albonico M, Savioli L (1997). Epidemiology of iron deficiency anemia in Zanzibari schoolchildren: the importance of hookworms. Am. J. Clin. Nutr. 65: 153-159.
- Van den Broek N (1996). The aetiology of anaemia in pregnancy in West Africa. Trop. Doct. 26: 5-7.
- Van den Broek NR, Letsky EA, White SA, Shenkin A (1998). Iron status in pregnant women: which measurements are valid? Br. J. Haematol. 103: 817-824.
- Van Eijk AM, Lindblade KA, Odhiambo F (2009). Geohelminth Infections among Pregnant Women in Rural Western Kenya; a Cross-Sectional Study. PLoS Negl. Trop. Dis. 3: e370.
- WHO (1993). Prevention and management of severe anaemia in pregnancy: report of a technical working group. Geneva. WHO/FHE/MSM/ 93.
- WHO (1996). Report of the WHO Informal Consultation on Hookworm Infection and Anaemia in Girls and Women World Health Organization Geneva. WHO/CTD/SIP/ 96.
- WHO (1998). UNICEF and UNU Iron Deficiency: Indicators for Assessment and Strategies for Prevention: Geneva.