

Full Length Research Paper

# The effect of gibberellins on russeting in golden delicious apples

F. Nejat-zadeh Barandoozi<sup>1\*</sup> and A. Talaie<sup>2</sup>

<sup>1</sup>Department of agricultural engineering, Islamic Azad University of Khoy, Iran.

<sup>2</sup>Department of agricultural engineering, Islamic Azad University of Tehran, Iran.

Accepted 12 June, 2009

**Russet on apples, *Malus domestica* Borkh and other fruits results in lowered fruit quality and substantial economic losses to growers. In this study we investigated the effect of GA<sub>3</sub> and GA<sub>4+7</sub> on russeting. Russeting index of fruits were estimated at harvest. Apple trees, twenty years old Golden Delicious was sprayed with GA<sub>3</sub> at the concentrations 10-20 mgL<sup>-1</sup> and GA<sub>4+7</sub> at 5-10 mgL<sup>-1</sup> during petal-fall at 7-10 days 3 time again. All applied concentration of GA<sub>3</sub> and GA<sub>4+7</sub> reduced russeting in Golden Delicious cultivar significantly and significant difference was found between GA<sub>3</sub> and GA<sub>4+7</sub> actions. The application of GA<sub>3</sub> or GA<sub>4+7</sub> caused fruits to shift to elongated shape due to increase of L/D ratio. The application of GA<sub>3</sub> or GA<sub>4+7</sub> caused to increase the weight of Golden Delicious fruit significantly. When GA<sub>3</sub> or GA<sub>4+7</sub> were separately sprayed on Golden Delicious, it resulted in considerable decrease in russeting neither return bloom nor were fruit set influenced by treatment.**

**Key words:** Apple russeting, golden delicious, gibberellins, GA<sub>3</sub> and GA<sub>4+7</sub>.

## INTRODUCTION

Russet on apples, *Malus domestica* Borkh and other fruits results in lowered fruit quality and substantial economic losses to growers (Cummins et al., 1977). Thorough reviews have been written on the etiology of fruit russet (Eccher and Boffelli, 1981; Fuast and Shear, 1972). Factors associated with apple russet include infection by the powdery mildew pathogen (*podosphaera Leucotricha*) (Gildemacher et al., 2004; Matteson-Heidenreich et al., 1997; Spotts, 2002) insects (Easterbrook and Fuller, 1986), cultural and environmental factors (Michailides, 1991), mechanical injuries (Walter, 1967) and pesticides (Taylor, 1978).

The use of some fungicides (Creasy and Swartz, 1981) silicon dioxides (Edgerton, 1976; Matteson-Heidenreich et al., 1997), antagonistic bacteria (Lindow, 1992) and gibberellic acid (Fouk and Hoover, 1994; Greene, 1993; Looney et al., 1992) was shown to reduce the incidence of apple russet. Microscopic observation of russeted areas of fruit shows breaks in the cuticle layer with underlying layers of dead epidermal cells (Ashizawa et al., 2005; Skene and Green, 1989).

Fruit cultivars differ greatly in their susceptibility to russet (Kirby and Bennett, 1967). In this study we investigated the effect of application of GA<sub>4+7</sub> and GA<sub>3</sub> on russeting of apple. We also investigated if the concentration of GA<sub>4+7</sub> and GA<sub>3</sub> is sufficient to reduce russeting of apple. The aim of this study was to compare the effect of GA<sub>4+7</sub> and GA<sub>3</sub> in control of russeting.

## MATERIALS AND METHODS

The experiments were conducted in the experimental Orchard of Agricultural Institute of Iran, kahriz Orumia in 2007-2008. Apple trees of cultivar "Golden Delicious" were used as experimental materials. A complete randomized blocks design with 13 trees of same size, homogenous and medium growth were used. The distance between trees was 6 x 6 m and age of the trees was 20 years. The concentrations of GA<sub>4+7</sub> and GA<sub>3</sub> used were 5, 10, 10 and 20 mgL<sup>-1</sup>. The concentrations were sprayed 3 times starting from petal-fall (P.F) stage and at intervals of 7-10 days. Single, whole selected tree were sprayed at petal-fall (22 may 2007) with a motorized air blast sprayer with GA<sub>3</sub> at a rate of 10 to 20 mgL<sup>-1</sup> and GA<sub>4+7</sub> at a rate of 5 to 10 mgL<sup>-1</sup> separately. The GA<sub>4+7</sub> and GA<sub>3</sub> treatment did not include a witting agent. Russeting of fruit was estimated at harvest using a scale of grading described by (Buban, 2001):

- (1) No russet.
- (2) Less than 30%.
- (3) 31 to 60%.

\*Corresponding author. E-mail: [fnejat-zadeh@yahoo.com](mailto:fnejat-zadeh@yahoo.com). Tel.: 0989126759651.




Figure 1. Russet index of GA<sub>4+7</sub> and GA<sub>3</sub> treated apples.




Figure 2. The effect of GA<sub>4+7</sub> and GA<sub>3</sub> on fruit shape. Indicate error bars.

- (4) More than 61% of the fruit surface with network-like russetting.  
 (5) Confluent severe russetting.

Mean fruit weight, seed number and fruit length: diameter ratio (L/D) was estimated on a random sample of 30 fruits/trees at harvest. The experiments were repeated 2 times. Data were subjected to statistical analysis of variance and LSD for mean comparisons at  $P=0.01$  was used.

## RESULTS

### The effect of GA<sub>4+7</sub> and GA<sub>3</sub> on russetting

This study has shown that GA<sub>4+7</sub> and GA<sub>3</sub> reduced russet in apple but did not have the same effect. Figure 1 shows the effect of GA<sub>4+7</sub> and GA<sub>3</sub> on russetting. Russetting Index indicated significance between control and case.

Although reduce russetting with increasing concentration and number of spray. The concentration of 10 mgL<sup>-1</sup> of GA<sub>4+7</sub> was significantly different with control.

### The effect of GA<sub>4+7</sub> and GA<sub>3</sub> on fruit shape

Both of GA<sub>4+7</sub> and GA<sub>3</sub> elongated the fruit shape resulting


in higher L/D ratios measured at harvest. There was a significant positive effect of GA<sub>4+7</sub> and GA<sub>3</sub> concentration on the fruit L/D ratio ( $P<0.01$ ). There was a markedly higher of elongated fruit (with L/D ratio of 0.91 or more). Figure 2 shows the effect of GA<sub>4+7</sub> and GA<sub>3</sub> fruit shape.

### The effect of GA<sub>3</sub> or GA<sub>4+7</sub> treatments on fruit weight

Result of this study showed that GA<sub>3</sub> and GA<sub>4+7</sub> treatments can increase fruit weight. The highest fruit weight was produced with concentration of 10 mgL<sup>-1</sup> GA<sub>4+7</sub> and lowest with the control (Figure 3). There was no relationship between number of seeds, firmness, color and GA<sub>3</sub> and GA<sub>4+7</sub> treatments.

## DISCUSSION

Treatments by GA<sub>3</sub> and GA<sub>4+7</sub> proved to be useful to increase the market value of fruits likewise to earlier reports on improving fruit shape and decrease in russetting. Non-treatment trees of "Golden Delicious" in this study were russeted. Therefore, treated trees were free of rus-



**Figure 3.** The effect of GA<sub>3</sub> and GA<sub>4+7</sub> treatments on fruit weight.

setting. 10 mgL<sup>-1</sup> GA<sub>3</sub> concentration has a few effect on reduction of russetting but caused increase L/D ratio, while 20 mgL<sup>-1</sup> GA<sub>3</sub> concentration caused increasing of inhabitation of fruit set in the year after treatment and increased fruit size and weight not significantly. Both GA<sub>4+7</sub> concentrations caused reduction of russetting and increase fruit size and weight significantly. When higher concentration of GA<sub>3</sub> or GA<sub>4+7</sub> was applied on the trees, there was increasing of inhabitation of flower bud formation and reduction of fruit number in the year after treatment and increased of cost (Taylor and Knight, 1986). There was no influence on return bloom. However, GA<sub>3</sub> and GA<sub>4+7</sub> treatment increased fruit weight because of thinning effect and decrease of fruit number.

In similar experiment in Australia on "Golden Delicious" influence on flower bud formation was found following the application of 24 mgL<sup>-1</sup> of GA<sub>4+7</sub> this effect while with application 10 mgL<sup>-1</sup> of GA<sub>4+7</sub> was not found this effect. When the higher concentration of GA<sub>3</sub> and GA<sub>4+7</sub> was used, the weight of "Golden Delicious" fruit was higher. These results confirm the thesis of additional fruit growth after GA application (Looney et al., 1992).

Similar experiments on "Golden Delicious" and "COX Orange Pippin" with rate of GA<sub>4+7</sub> 2.5-10 mg/l (depending on cultivar) applied 3 or 4 times at 10 day intervals starting during the period of first-flower to petal-fall to achieved satisfactory control (Skene and Green, 1989).

While some authors found a positive influence of GA<sub>3</sub> and GA<sub>4+7</sub> on return bloom (Jones et al., 2004) in other trial the application of GA<sub>4+7</sub> 12 or 24 mgL<sup>-1</sup> did not increase the return bloom of "Golden Delicious" trees (Taylor and Knight, 1986). In this study, best treatment was 10 mgL<sup>-1</sup> GA<sub>4+7</sub> in control of russetting on "Golden Delicious" and improved fruit shape, weight and increase quality of fruits and reduce economic losses and has not effect on return bloom.

## REFERENCE

Ashizawa M, Horigome Y, Chujo T (2005). Histological studies on the cause of russet in "Golden Delicious" apple, Hort. Sci. 30: 47-52.  
 Buban T (2001). Improved fruit shape and Less Russetting of apples by

using Gibberellins Acta. Hort. 20: 14-17.  
 Creasy LL, Swartz HJ (1981). Agents influencing russet on apple (*Malus domestica* cultivar Golden Delicious) J. Am. Soc. Hort. Sci., pp: 735-738.  
 Cummins JN, Forsline PL, Way RD (1977). A comparison of russetting among "Golden Delicious" Sub clones Hort. Sci. 12: 241-242.  
 Easterbrook MA, Fuller MM (1986). Russetting of apples caused by apple rust mite *Aculus schlechtendali* Acarina Eriophidae Annu. Appl. Biol., 109: 1-10.  
 Eccher T, Boffelli G (1981). Effects of dose and time application of gibberellin A4 plus A7 on russetting, fruit set and shape of "Golden Delicious" apples Hort. Sci., 14: 307-314.  
 Edgerton LNJ (1976). Foliage sprays of Silicon dioxide containing compound reduce russetting in "Golden Delicious" apple fruits J. Hort. Sci 12: 28-34.  
 Foulk DS, Hoover EE (1994). Wax platelet arrangement GA4+7 applications, and their relationships to russet in: Haralsonapples. Hort. Sci. 29: 472-477.  
 Fuast M, Shear CB (1972). Russetting of apples, an interpretive review. Hort. Sci. 7: 233-235.  
 Gildemacher P, Heijne B, Houbraken J, Vromans T, Hoekstra E, Boekhout T (2004). Can phyllosphere Yeasts explain the effect of scab fungicides on russetting of elstar apples? Eur. J. Plant Path. 110: 929-937.  
 Greene DW (1993). Effects of GA7 on flower bud formation and russet development on apple. J. Hort. Sci. 68: 171-176.  
 Jones K, Bound S, Oakford M, Wilson D (2004). A strategy for reducing russet in Red Fuji apples while maintaining control of black spot (*Venturia inaequalis*). Aust. J. Exp. Agric. 34: 127-130.  
 Kirby AHM, Bennett M (1967). Susceptibility of apple and pear varieties to damage by certain organic fungicides. J. Hort. Sci. 42: 117-131.  
 Lindow SE (1992). Integrated control of frost injury, fire blight, and fruit russet of pear with a blossom application of an antagonistic bacterium. (Abstr.). Phytopathol. 82: 11-29.  
 Looney NE, Granger RL, Haris RP (1992). Influences of gibberellins A-4 gibberellins A4 plus gibberellin A-7 and gibberellin A-4 plus Iso-minus gibberellin A-7 on apple fruit quality and tree productivity. Effects of fruit russet and tree yield components. J. Hort. Sci. 67: 613-618.  
 Matteson HMC, Corral GEA, Momol EA, Burr TJ (1997). Russet of apple fruit caused by *Aureobasidium pullulans* and *Rhodotorula glutinis*. plant Dis. 81: 337-342.  
 Michailides TJ (1991). Russetting and russet scab of prune, an environmentally induced fruit disorder: symptomatology, induction, and control. Plant Dis. 75: 1114-1123.  
 Skene DS, Green DW (1989). The development of russet, rough russet and cracks on the fruit of the apple "Cox's Orange Pippin" during the course of the season. J. Hort. Sci. 56: 165-174.  
 Spotts RC (2002). Involvement of *Aureobasidium Pullulans* and *Rhodotorula glutinis* in russet of d' Anjou pear fruit. *Plant Dis.* 86: 625-628.  
 Taylor BL (1978). Effects of gibberellin sprays on fruit russet and tree performance of Golden Delicious apple. J. Hort. Sci. 53: 167-169.  
 Taylor DR, Knight JN (1986). Russetting and cracking of apple fruit and

their control with plant Growth Regulators. (Abstr.). Acta. Hort. 179: 819-820.

Walter TE (1967). Russeting and cracking in apples. A review of world literature. Annu. Rep. Malling Res. pp: 83-95.