

*Full Length Research Paper*

# Indigenous knowledge of plant uses by the community of Batiaghata, Khulna, Bangladesh

Tama Ray<sup>1\*</sup>, Sharif Hasan Limon<sup>2</sup>, Sajjad Hossain Tuhin<sup>2</sup> and Arifa Sharmin<sup>2</sup>

<sup>1</sup>Faculty of Forest Sciences and Forest Ecology, University of Göttingen, Goettingen-37077, Germany.

<sup>2</sup>Forestry and Wood Technology Discipline, Khulna University, Khulna-9208, Bangladesh.

Received 15 December, 2020; Accepted 6 May, 2021

Southwestern region of Bangladesh is very rich in floral diversity, and their diversified uses. An extensive survey was conducted to investigate ethnobotanical applications of botanical species by the community of Khulna, Bangladesh. We focused on plants and community relationships, identify the most important species used, determine the relative importance of the species surveyed and calculated the Fidelity level (FI) and Cultural Significance Index (CSI) concerning individual species. In total, we have listed 136 species of 114 genera under 52 families, of which 41% (63) species were utilized as food, followed by 30% (45) medicine, 14% (21) constructional timber, 11% (17) ornamental, and 4% (6) other uses. Inheritance of traditional knowledge of medicinal plants was the primary source of knowledge acquisition through oral transmission over the generations. However, only 34% of the informants were traditional herbal practitioners. Most of the medicinal uses are primly associated with anti-inflammatory, anti-microbial, antiseptic, expectorant, antidote, fever reduction, and pain relief.

**Key words:** Plants, ethnobotany, homestead, Khulna, Bangladesh.


## INTRODUCTION

Geographically, most of the territories of Bangladesh are formed by a delta plain with a tropical monsoon climate. It is vibrant with a vast biological diversity (Chowdhury and Koike, 2010) and lies under the Indo-Burma biological hotspot area (Mukul et al., 2008). In total, 7,000 floral species were listed from this area with several endemic plants, and 50% of them are herb, 35% shrub, and woody climber and 15% tree (Rahman, 2004). Meanwhile, angiosperms are dominated in checklists with 5,700 species followed by 1,700 pteridophytes, 500 medicinal plant species, 130 fiber yielding plants species, 68 woody legume species, 29 orchid species and three species of

gymnosperm (Mukul et al., 2008). However, Ahamed et al. (2007-2009), has enlisted 3611 angiosperm, 195 pteridophyte species, and seven gymnosperm species, excluding Bacteria and fungi. In addition to that, there are 750-800 more tree species found in these areas, including indigenous, exotic, and naturalized ones (Irfanullah, 2011). Among all different regions, the hilly region is the richest one in consideration of floristic diversity and richness with 2,260 plant species (Mukul et al., 2008).

In Bangladesh, humans and plants share the natural habitats with traditional bonding and influence each other

\*Corresponding author. E-mail : [tuhin.taxon@gmail.com](mailto:tuhin.taxon@gmail.com); Tel.+8801722240090.



**Figure 1.** Map showing the study area.

(Partha, 2014). Still, at most of the aspects of biological and economic needs, people depend on plants for food, shelter, construction materials, clothing, medicines, rituals, fuelwood, household implements, musical instruments, pesticides and so on (Gemedo-Dalle et al., 2005). This dependency became the grass-root basis of species conservation for humans (Singh et al., 2002). But at present, people are using trees in an exploiting manner for the economy (Idu, 2009; Kargioğlu et al., 2008). As a result, number of plants species is decreasing at faster rate, along with the knowledge of traditional use of those species (Balick and Cox, 1996; Avocèvou-Ayisso et al., 2012; Asafo-Agyei et al., 2019; Abdala and Carlos, 2020; Korkmaz and Karaku, 2015). Several studies have identified the scarcity of ethnobotanical information and the lack of transmission of ethnobotanical knowledge from generation to generation as the crucial factors in disappearance of species from a locality (Khan et al., 2018). Currently, ethnobotanical knowledge about medicinal plants found exposure to the scientific community and several studies have been conducted regarding this issue (Yusuf et al., 2009; Partha and Hossain, 2007; Roy et al., 2008; Rahmatullah et al., 2012; Uddin et al., 2012). But most of the research missing the other usages bear comparatively the same

importance in conservation and site-specific parameters. So, we considered in this study to identify species composition and diversity, analyze the uses of species and their mode of use, and evaluate the value or importance of species within the culture.


## METHODOLOGY

### Study area

The study was conducted in Batiaghata Upazila under Khulna district. It is located between 22°46'07'' N to 22°37'50'' N and 89°24'14'' E to 89°31'47'' E (Figure 1). Batiaghata experiences a subtropical climatic condition with a mild winter from October to March, hot and humid summer in March to June and moist, warm rainy monsoon in June to October. In December-January, the temperature fell to the lowest at 12-15°C, and it reached highest in April-June at 41 -45°C. Most of the rainfall during June to October. July is the month of maximum precipitations with 20-25 days of rain. Average wind speed is over 8 km/h during April-August, which is the highest value for this area (BBS, 2014).

### Demography

Like other Upazilas of Bangladesh, Batiaghata is densely populated with a total population of 140,574 in which 72,717 are males and



**Figure 2.** Composition of identified species, (a) different uses, (b) life forms.

67,857 females with 40,779 units of households. Among the total population, Muslims dominate with 79,301 along with 60,894 Hindu, 285 Buddhist, 6 Christian, and 85 others. Administratively, Batiaghata Upazila has seven unions named Amirpur, Baliadanga, Batiaghata, Bhandarkote, Gangarampur, Jalma, and Surkhali. Almost 91% of people are engaged in agriculture, followed by service (7%) and industry (2%) (BBS, 2011, 2013).

### Sampling design

A reconnaissance survey was conducted in the study area before questionnaire preparation to obtain general information about the villages and the villagers. Depending on this survey, a semi-structured questionnaire was prepared for ethnobotanical information collection. Five unions (Batiaghata, Baliadanga, Gangarampur, Amirpur and Surkhali) have been selected and one village from each union was chosen to survey by random selection. In total, 150 households were studied in this study, where 30 houses were chosen randomly from every village for data collection. Interviewees were divided into five age groups (20-34, 35-49, 50-64, 65-79, 80 and above) as age plays a distinctive role in ethnobotanical knowledge (Nawash et al., 2014). Cited plant species were checked physically, photographed, and voucher specimens were collected for further identification and conservation. Collected specimens were analyzed and identified based on the key provided by Hooker (1872-1890), Prain (1903-04), Kanjilal et al. (1934-1940), Deb (1983), Matthew (1999) and Ahmed et al. (2007-09).

### Calculations

After collecting data, they were categorized according to their specific use such as food, medicine, construction, fuel, ornamental, and others and analyzed according to the following indices:

(a) Fidelity level,  $FL = I_p - I_u * 100\%$  (Friedman, 1986; Hoffman and Gallaher, 2007)

Here,  $I_p$  = Number of informants who cited the species for the particular use.

$I_u$  = Total number of informants that mentioned the plant for any use.

(b) Cultural significance index,  $CSI = \sum (i * e * c) * CF$  (Turner, 1988; Stoffl et al., 1990; Hoffman and Gallaher, 2007)

Here,  $I$  = species management where, 1 indicates non- managed

and 2 indicates managed.

$E$  = Use preference where 1 indicates non- preferred and 2 indicates preferred.


$C$  = Use frequency where 1 indicates rarely used, and 2 indicates frequently used.

$CF$  (Correction factor) = Number of citations for a given species divided by the number of citations for the most mentioned species.

## RESULTS AND DISCUSSION

In total, 136 species of 114 genera under 52 families have been identified throughout this study. Among the counted species, 41% (63) species were utilized as food, followed by 30% (45) medicine, 14% (21) constructional timber, 11% (17) ornamental, and 4% (6) other uses (Figure 2a). Forty-four species have been cited by the informants to be used in the treatment of various human diseases, including respiratory, digestive, liver, skin, rheumatism, diabetes, cancer, and other disorders. Among the 150 informants interviewed, 72.6% were males, and 27.4% were females, with 74.3% above 50 years of age. Inheritance of traditional knowledge of medicinal plants was the primary source of knowledge acquisition through oral transmission over the generations. However, only 34% of the informants were traditional herbal practitioners, with the remaining majority (66%) of informants having no professional practice of herbal medicine.

One hundred thirty-six floral species have been cited in the homestead of the study area belonging to 52 families and 114 genera. Fabaceae (17) found the most dominant family followed by Anacardiaceae (6), Myrtaceae (6), Apocynaceae (5), Arecaceae (5), Malvaceae (5), Moraceae (5), Solanaceae (5) (Figure 3). Horticultural species like *Syzygium* (5), *Terminalia* (4) *Artocarpus* (2) found most dominant over other species. The life forms and growth habits of plants were distributed into 49.3% trees, 19.9% herb, 1.5% grass, 1.5% palm, and 0.7% vine (Figure 2b). Most of the medicinal uses are undoubtedly associated with anti-inflammatory, anti-microbial and antiseptic, antibacterial, expectorant,



**Figure 3.** Dominant families of the study area.

antidote, fever reduction, and pain relief.

*Mangifera indica* found most cited species over the study area with a high informant consensus (IC) of 86, followed by *Areca catechu* (79), *Cocos nucifera* (76), *Ocimum tenuiflorum* (73), *Swietenia macrophylla* (73), *Albizia lebbek* (69) and so on. In terms of Fidelity Index (FI %), 114 species scored 100%, which indicates the dedicated use of those species without any alternatives. Cultural Significance Index ranges from 13.58 to 0.08. Six species lie above ten, namely, *Ocimum tenuiflorum* (13.58), *M. indica* (13), *Areca catechu* (11.94), *Cocos nucifera* (11.49), *Swietenia macrophylla* (11.03), *Albizia lebbek* (10.43) (Table 1).

A considerable amount of floral species has been used traditionally by the local community with diversified implications in the study area. Very few studies have been done in this region to preserve the ethnobotanical knowledge inherited by generations. Nawaz et al. (2009), listed 26 plant species from 22 families used in ethnomedicine from Khulna et al. (2009) identified 33 species in folk medicine from Khulna division and Ray and Mondal (2018) describe 25 species from Shyamnagar, Satkhira near Sundarbans. However, at present studies, 136 species of 114 genera under 52 families were identified, of which 45 species have been reported to be

used in folk medicine. The use-value indicates the total number of uses of a specific species and two types of tally have been used for calculations, Uses Total (Researcher Tally) indicated specific applications and Use- Value indicated individual allocation. However, in researcher tally, the uses were recorded, ranked, and summed, which showed a similar contingency to previous studies (Rahman, 2013; Ray and Mondal, 2018; Faruque et al., 2018). In this study, food, fuel, medicine, construction, ornamental, and other categories were used to investigate multiple uses of a single species and numerous species for individual use.

We have found the highest species use-value for *M. indica* (0.95), and total use is also very high for it (86), which means the highest number of people engage with this species by its meaningful use at their daily life. Meanwhile, *Cuscuta reflexa* remains at the lowest level (0.01) for its minimal use by the lowest number of people from the participants. In terms of fidelity level, it describes the importance of the species for any specific purpose. It is used to identify the preferable species used by key informants for one particular treatment. Species having high fidelity levels is generally widely used for dedicated purposes. It also illustrates the number of informants in the percentage who state the use of certain species for

**Table 1.** Ethnobotanical indices of Batiaghata, Khulna (IC=Informant Citation; UT=Used Total; FI=Fidelity Level; CSI=Cultural Significance Index; F=Food; M=Medicine; F=Fuel; C=Construction; Or=Ornamental and O=Other).

Species	Family	Habit	Use	UT	FI%	CSI
<i>Abroma augusta</i> (L.) L.fil.	Malvaceae	Tree	M	1	100	0.1
<i>Acalypha hispida</i> Burm.f.	Euphorbiaceae	Herb	M	1	100	0.08
<i>Acalypha wilkesiana</i> Müll.Arg.	Euphorbiaceae	Shrub	O	14	100	1.47
<i>Aegle marmelos</i> (L.) Correa	Rutaceae	Tree	F	35	100	3.66
<i>Albizia lebbbeck</i> (L.) Benth.	Fabaceae	Tree	C	69	100	10.43
<i>Albizia richardiana</i> (Voigt) King & Prain	Fabaceae	Tree	C	55	100	7.67
<i>Allium cepa</i> L.	Liliaceae	Herb	F	3	66.67	0.31
<i>Alocasia macrorrhizos</i> (L.) G.Don.	Araceae	Herb	F	4	100	0.46
<i>Aloe vera</i> (L.) Burm.f.	Liliaceae	Herb	M	1	100	0.1
<i>Alstonia scholaris</i> (L.) R. Br.	Apocynaceae	Tree	C	7	100	0.73
<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	Anacardiaceae	Herb	F	4	100	1.54
<i>Amaranthus tricolor</i> L.	Anacardiaceae	Herb	F	45	100	4.7
<i>Amaranthus viridis</i> L.	Anacardiaceae	Herb	F	48	100	3.9
<i>Amorphophallus bulbifer</i> (Roxb.) Blume	Araceae	Herb	F	13	92.31	1.26
<i>Andrographis paniculata</i> (Burm.f.) Nees	Acanthaceae	Herb	M	1	100	0.31
<i>Anisoptera scaphula</i> (Roxb.) Kurz	Dipterocarpaceae	Tree	M	1	100	0.1
<i>Annona squamosa</i> L.	Annonaceae	Tree	F	13	100	1.36
<i>Aphanamixis polystachya</i> (Wall.) R.Parker	Meliaceae	Tree	M	1	100	0.16
<i>Areca catechu</i> L.	Arecaceae	Plam	F	79	100	11.94
<i>Artocarpus heterophyllus</i> Lam.	Moraceae	Tree	F	51	100	7.71
<i>Artocarpus lacucha</i> Buch-Ham.	Moraceae	Tree	C	39	94.87	4.3
<i>Arundina graminifolia</i> (D.Don) Hochr.	Orchidaceae	Herb	Or	7	100	0.57
<i>Asparagus racemosus</i> Willd.	Asparagaceae	Shrub	M	1	100	0.1
<i>Averrhoa carambola</i> L.	Oxalidaceae	Tree	F	41	100	4.29
<i>Azadirachta indica</i> A.Juss.	Meliaceae	Tree	M	59	100	6.17
<i>Bacopa monnieri</i> (L.) Wettst.	Plantaginaceae	Herb	F, M	5	100	0.52
<i>Bambusa tuldoidea</i> Munro	Poaceae	Grass	C	18	100	1.67
<i>Basella alba</i> L.	Basellaceae	Herb	F	36	100	3.77
<i>Berberis asiatica</i> Roxb. ex DC.	Berberidaceae	Shrub	M	1	100	0.08
<i>Boerhavia diffusa</i> L.	Nyctaginaceae	Herb	M	1	100	0.1
<i>Bombax ceiba</i> L.	Malvaceae	Tree	C, O	60	100	3.45
<i>Borassus flabellifer</i> L.	Arecaceae	Plam	F	52	100	7.86
<i>Bryophyllum pinnatum</i> (Lam.) Oken	Crassulaceae	Herb	M	2	100	0.16
<i>Butea monosperma</i> (Lam.) Taub.	Fabaceae	Tree	C	16	94.11	1.67
<i>Caesalpinia pulcherrima</i> (L.) Sw.	Fabaceae	Shrub	M	1	100	1.78
<i>Calotropis gigantea</i> (L.) Dryand.	Apocynaceae	Shrub	M	1	100	0.1
<i>Capsicum annuum</i> L.	Solanaceae	Herb	F	57	100	5.97
<i>Carica papaya</i> L.	Caricaceae	Tree	F	53	100	8.01
<i>Carissa carandas</i> L.	Apocynaceae	Shrub	F	11	100	1.15
<i>Caryota urens</i> L.	Arecaceae	Plam	M	1	100	0.1
<i>Cassia fistula</i> L.	Fabaceae	Tree	C	56	62.5	5.44
<i>Catharanthus roseus</i> (L.) G.Don	Apocynaceae	Shrub	M	1	100	0.21
<i>Celosia cristata</i> L.	Anacardiaceae	Herb	O	7	100	0.73
<i>Centella asiatica</i> (L.) Urb.	Apiaceae	Herb	M	34	100	2.78
<i>Cestrum nocturnum</i> L.	Solanaceae	Shrub	O	2	100	0.21
<i>Chromolaena odorata</i> (L.) R.M.King & H.Rob.	Asteraceae	Herb	M	18	100	1.88
<i>Cinnamomum verum</i> J.Presl	Lauraceae	Tree	FM	2	100	0.21
<i>Citrus aurantiifolia</i> (Christm.) Swingle	Rutaceae	Shrub	F	83	68.37	8.62
<i>Citrus grandis</i> (L.) Osbeck	Rutaceae	Tree	F	39	100	4.08
<i>Clerodendrum chinense</i> (Osbeck) Mabb.	Lamiaceae	Shrub	O	4	100	0.42

Table 1. Cont'd

<i>Clitoria ternatea</i> L.	Fabaceae	Herb	O	3	60	0.52
<i>Cocos nucifera</i> L.	Arecaceae	Plam	F	76	100	11.49
<i>Cucurbita maxima</i> Duchesne	Cucurbitaceae	Herb	F	17	100	1.78
<i>Cucurbita moschata</i> Duchesne	Cucurbitaceae	Herb	F	9	100	0.94
<i>Curcuma domestica</i> Valetou	Zingiberaceae	Herb	F, M	8	100	0.52
<i>Cuscuta reflexa</i> Roxb.	Convolvulaceae	Herb	M	1	100	1.15
<i>Cynodon dactylon</i> (L.) Pers.	Poaceae	Grass	M	98	56.12	5.76
<i>Dalbergia sissooides</i> Wight & Arn.	Fabaceae	Tree	C	58	100	6.07
<i>Datura metel</i> L.	Solanaceae	Shrub	M	1	100	0.15
<i>Delonix regia</i> (Hook.) Raf.	Fabaceae	Tree	C	17	100	1.78
<i>Diospyros malabarica</i> (Desr.) Kostel.	Ebenaceae	Tree	O	97	50.52	6.84
<i>Diospyros montana</i> Roxb.	Ebenaceae	Tree	C	1	100	0.24
<i>Elaeocarpus tectorius</i> (Lour.) Poir.	Elaeocarpaceae	Tree	F	36	100	3.77
<i>Elettaria cardamomum</i> (L.) Maton	Zingiberaceae	Herb	F	2	50	0.1
<i>Erythrina ovalifolia</i> Roxb.	Fabaceae	Tree	C	29	100	2.36
<i>Excoecaria agallocha</i> L.	Euphorbiaceae	Tree	C	20	85	2.37
<i>Feronia limonia</i> (L.) Swingle	Rutaceae	Tree	F	8	100	0.84
<i>Ficus auriculata</i> Lour.	Moraceae	Tree	F	53	100	5.55
<i>Ficus benghalensis</i> L.	Moraceae	Tree	C	12	100	1.26
<i>Ficus religiosa</i> L.	Moraceae	Tree	M	4	100	0.41
<i>Gardenia jasminoides</i> J.Ellis	Rubiaceae	Shrub	O	5	100	0.52
<i>Glebionis coronaria</i> (L.) Cass. ex Spach	Asteraceae	Herb	M	1	100	0.57
<i>Gossypium arboreum</i> L.	Malvaceae	Shrub	O	49	100	6.84
<i>Helianthus annuus</i> L.	Asteraceae	Herb	F, O	3	100	0.21
<i>Heritiera fomes</i> Buch.-Ham.	Sterculiaceae	Tree	C	3	100	0.31
<i>Hibiscus esculentus</i> L.	Malvaceae	Shrub	F	4	100	0.56
<i>Hibiscus rosa-sinensis</i> L.	Malvaceae	Shrub	O	55	94.54	5.76
<i>Ipomoea aquatica</i> Forssk.	Convolvulaceae	Herb	F	8	87.5	0.73
<i>Juglans regia</i> L.	Juglandaceae	Tree	M	1	100	0.1
<i>Justicia adhatoda</i> L.	Acanthaceae	Shrub	M	1	100	0.2
<i>Lablab purpureus</i> (L.) Sweet	Fabaceae	Herb	F	49	100	5.13
<i>Lawsonia inermis</i> L.	Lythraceae	Shrub	Or	24	100	3.35
<i>Leucaena leucocephala</i> (Lam.) de Wit	Fabaceae	Tree	C	16	100	1.49
<i>Litchi chinensis</i> Sonn.	Sapindaceae	Tree	F	45	100	6.8
<i>Mangifera indica</i> L.	Anacardiaceae	Tree	F	86	100	13
<i>Manilkara zapota</i> (L.) P.Royen	Sapotaceae	Tree	F	63	100	9.52
<i>Mentha spicata</i> L.	Lamiaceae	Herb	F	2	50	0.16
<i>Mesua ferrea</i> L.	Calophyllaceae	Tree	M	1	100	0.31
<i>Mimosa pudica</i> L.	Fabaceae	Herb	M	1	100	2.41
<i>Mimusops elengi</i> L.	Sapotaceae	Tree	O	9	100	0.94
<i>Momordica charantia</i> L.	Cucurbitaceae	Herb	F	21	100	1.71
<i>Moringa oleifera</i> Lam.	Moringaceae	Tree	F	64	100	9.67
<i>Musa paradisiaca</i> cv. Awak	Musaceae	Tree	F	17	100	1.78
<i>Neolamarckia cadamba</i> (Roxb.) Bosser	Rubiaceae	Tree	C	3	66.67	0.3
<i>Ocimum tenuiflorum</i> L.	Lamiaceae	Shrub	M, O	120	100	13.58
<i>Opuntia cylindrica</i> (Lam.) DC.	Cactaceae	Herb	O, M	29	100	2.36
<i>Oxalis corniculata</i> L.	Oxalidaceae	Herb	F	46	95.65	4.6
<i>Phoenix sylvestris</i> (L.) Roxb.	Arecaceae	Plam	F	43	100	6.5
<i>Phragmites australis</i> (Cav.) Trin. ex Steud.	Poaceae	Grass	M	4	100	0.42
<i>Phyllanthus emblica</i> L.	Phyllanthaceae	Tree	F, M	90	54.44	9.12
<i>Piper betle</i> L.	Piperaceae	Herb	F, M	3	100	0.16
<i>Piper longum</i> L.	Piperaceae	Herb	M	1	100	0.16

Table 1. Cont'd

<i>Piper nigrum</i> L.	Piperaceae	Shrub	F, M	6	100	0.24
<i>Piper retrofractum</i> Vahl	Piperaceae	Shrub	F	56	100	5.86
<i>Pithecellobium dulce</i> (Roxb.) Benth.	Fabaceae	Tree	F, U	68	61.76	4.39
<i>Polyalthia longifolia</i> (Sonn.) Thwaites	Annonaceae	Tree	O	6	100	0.56
<i>Prunus bokhariensis</i> Royle ex C.K.Schneid.	Rosaceae	Shrub	M	1	100	0.24
<i>Psidium guajava</i> L.	Myrtaceae	Tree	F	66	100	9.98
<i>Punica granatum</i> L.	Lythraceae	Shrub	F	31	100	3.24
<i>Rauvolfia serpentina</i> (L.) Benth. ex Kurz	Apocynaceae	Shrub	M	1	100	0.16
<i>Rosa x damascena</i> Herrm.	Rosaceae	Shrub	Or	13	100	1.36
<i>Santalum album</i> L.	Santalaceae	Tree	M	2	100	0.84
<i>Senna siamea</i> (Lam.) H.S.Irwin & Barneby	Fabaceae	Tree	C	9	75	1.15
<i>Sesbania grandiflora</i> (L.) Pers.	Fabaceae	Tree	O	1	33.33	0.31
<i>Smilax zeylanica</i> L.	Smilacaceae	Vine	M	1	100	0.31
<i>Solanum melongena</i> L.	Solanaceae	Shrub	F	47	100	4.92
<i>Solanum tuberosum</i> L.	Solanaceae	Shrub	F	26	100	2.72
<i>Sonneratia apetala</i> Buch.-Ham.	Lythraceae	Tree	F	7	100	0.73
<i>Spondias dulcis</i> Parkinson	Anacardiaceae	Tree	F	37	100	3.87
<i>Swietenia macrophylla</i> King	Meliaceae	Tree	C	73	100	11.03
<i>Syzygium aromaticum</i> (L.) Merr. & L.M.Perry	Myrtaceae	Tree	F, M	3	100	0.28
<i>Syzygium cumini</i> (L.) Skeels	Myrtaceae	Tree	F	63	100	9.52
<i>Syzygium fruticosum</i> DC.	Myrtaceae	Tree	M	1	100	0.57
<i>Syzygium jambos</i> (L.) Alston	Myrtaceae	Tree	F	1	100	0.1
<i>Syzygium samarangense</i> (Blume) Merr. & L.M.Perry	Myrtaceae	Tree	F	51	100	5.34
<i>Tamarindus indica</i> L.	Fabaceae	Tree	F, C	102	54	8.95
<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Arn.	Combretaceae	Tree	M	25	100	3.78
<i>Terminalia bellirica</i> (Gaertn.) Roxb.	Combretaceae	Tree	F, M	62	100	4.74
<i>Terminalia catappa</i> L.	Combretaceae	Tree	F	43	100	4.5
<i>Terminalia chebula</i> Retz.	Combretaceae	Tree	M	33	100	3.45
<i>Trichosanthes dioica</i> Roxb.	Cucurbitaceae	Tree	F	3	100	0.31
<i>Vigna unguiculata</i> (L.) Walp.	Fabaceae	Herb	F	16	100	1.67
<i>Vitex negundo</i> L.	Lamiaceae	Shrub	M	1	100	0.08
<i>Xylocarpus mekongensis</i> Pierre	Meliaceae	Tree	C	12	100	1.26
<i>Zingiber officinale</i> Roscoe	Zingiberaceae	Herb	F	8	87.5	0.73
<i>Ziziphus jujuba</i> Mill.	Rhamnaceae	Tree	F	47	100	7.1

the same purpose (Khan et al., 2014). Following that, fidelity levels were calculated highest (100%) for most of the species (114), which represents the single specific use of those species. But lower fidelity level indicates multiple uses of a species, and we found 24 species have various applications with lower fidelity levels. *Alocasia indica* has the fidelity level 100% having one primary purpose of use as food. 114 (83.33%) species have the highest (100%) fidelity level, which indicates single use of those species. *Sesbania grandiflora* found lowest FI% (33.33), indicating multiple uses (Table 1). Cultural Significant index (CSI) indicates the versatility of the application of a species along with the number of informants it uses, which means the spread of the use of the species (Parthiban et al., 2016). We have calculated the highest CSI for *O. tenuiflorum* (13.58), which indicates

different uses like medicinal and worship purposes.

The identified species composition showed higher diversity in use and practice. Religion, social structure, and economic conditions plays pivotal role to regulate the level of applications and inheritance of ethnobotanical knowledge. The current study found that most of the information (66%) about ethnobotany was inherited through oral communications. The aged personals of the community have a significant role in transmitting this knowledge among the population. In addition to that, several professionals, locally named as Kabiraj or Gunim (traditional healer), also have a strong influence over folk medicine use at the community level. Local people also cited that the use and practice of folk medicine is diminishing. Limited use of this folk medicine also threatened the conservation of the species in the region.

Currently, most of the medicinal purposes are limited to treat anti-inflammatory, anti-microbial and antiseptic, antibacterial, expectorant, antidote, fever reduction, and pain relief.

## CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

## ACKNOWLEDGEMENT

The authors appreciate the local personals participated in the study and also grateful to the Forestry and Wood Technology Discipline for technical support, Mr. Kingson Mondal for his comments on the initial version of the manuscript and Late. Md. Shirajul Islam Sheikh to collect field data.

## REFERENCES

- Abdala K, Carlos S (2020). Bioprospective potential in the traditional use of herbal medicines in an institution in Central Brazil. *Journal of Medicinal Plants Research* 14(8):389-405.
- Ahamed ZU, Begum ZNT, Khondokar M, Kabir SMH, Ahmed M, Ahmed ATA, Rahaman AKA, Haque EU (eds) (2007-2009). *Encyclopedia of Flora and Fauna of Bangladesh*, Vol 6-12. Asiatic Society of Bangladesh, Dhaka, Bangladesh.
- Asafo-Agyei T, Blagogee HR, Mintah SO, Archer MA, Ayertey F, Sapaty AC, Jnr PA, Boamah D, Asiedu-Larbi J, Appiah AA (2019). Ethnobotanical studies of medicinal plants used in traditional treatment of malaria by some herbalists in Ghana. *Journal of Medicinal Plants Research* 13(16):370-383.
- Avocèvou-Ayisso C, Avohou TH, Omorou M, Dessou G, Sinsin B (2012). Ethnobotany of *Pentadesma butyracea* in Benin: A quantitative approach. *Ethnobotany Research and Applications* 10:151-166.
- Balick MJ, Cox A (1996). *Plants that heal; people and culture: the science of ethnobotany*. Scientific American Library, Bronx, New York, USA.
- Bangladesh Bureau of Statistics (BBS) (2011). *Population and housing census 2011 Dhaka, Bangladesh*. Statistics Division, Dhaka, Bangladesh.
- Bangladesh Bureau of Statistics (BBS) (2013). *District Statistics 2011 Khulna*. Dhaka, Bangladesh. Statistics Division, Dhaka, Bangladesh.
- Bangladesh Bureau of Statistics (BBS) (2014). *Statistical Pocketbook of Bangladesh – 2013*. Ministry of planning. People's Republic of Bangladesh, Dhaka, Bangladesh.
- Chowdhury MSH, Koike M (2010). An overview on the protected area system for forest conservation in Bangladesh. *Journal of Forestry Research* 21(1):111-118. Available at: <http://dx.doi.org/10.1007/s11676-010-0019-x>
- Deb DB (1983). *The flora of Tripura*, Vol (1-3). Today and tomorrow printers and publishers, New Delhi, India.
- Faruque MO, Uddin SB, Barlow JW, Hu S, Dong S, Cai Q, Li X, Hu X (2018). Quantitative ethnobotany of medicinal plants used by indigenous communities in the Bandarban District of Bangladesh. *Frontiers in Pharmacology* 9:40. Available at: <http://dx.doi.org/10.3389/fphar.2018.00040>
- Friedman J, Yaniv Z, Dafni A, Palewitch D (1986). A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among Bedouins in the Negev Desert, Israel. *Journal of Ethnopharmacology* 16:275-287. Available at: [http://dx.doi.org/10.1016/0378-8741\(86\)90094-2](http://dx.doi.org/10.1016/0378-8741(86)90094-2)
- Gemedo-Dalle T, Maass BL, Isselstein J (2005). Plant biodiversity and ethnobotany of Borana pastoralists in southern Oromia, Ethiopia. *Economic Botany* 59(1):43-65.
- Hoffman B, Gallaher T (2007). Importance indices in ethnobotany. *Ethnobotany Research and Applications* 5:201-218.
- Hooker JD (1872-1897). *The Flora of British India*, Vols. (1-7) (Reprint. 1973). L. Reeve & Co., Dehradun, India.
- Idu M (2009). Ethnobotany today-who are the players? *Tropical Journal of Pharmaceutical Research* 8(4):295-296.
- Irfanullah HM (2011). Conserving threatened plants of Bangladesh: Miles to go before we start? *Bangladesh Journal of Plant Taxonomy* 18(1):81-91
- Kanjilal UN, Kanjilal PC, De RN, Das A (1934-1940). *Flora of Assam*, Vol 1-4. (Reprint. 1982). A Von Book Company, Delhi, India.
- Kargioğlu M, Ceneci S, Serteser A, Evliyaoğlu N, Konuk M, Kök MS, Bağcı Y (2008). An ethnobotanical survey of inner-West Anatolia, Turkey. *Human Ecology* 36(5):763-777.
- Khan M, Shinwari ZK, Shah M, Musharaf S (2014). An ethnobotanical survey of medicinal and other useful plants of Khattak Tribe in Tehsil Karak, Khyber Pakhtunkhwa, Pakistan. *Medicinal Plant Research* 4(8):61-66.
- Khan MN, Razzaq A, Hadi F, Khan N, Basit A, Jan F, Khan N (2018). Ethnobotanical profile of weed flora of district Charsadda, Khyber Pakhtunkhwa. *RADS Journal of Biological Research & Applied Sciences* 9(1):14-23.
- Korkmaz M, Karakuş SİNEM (2015). Traditional uses of medicinal plants of Üzümlü district, Erzincan, Turkey. *Pakistan Journal of Botany* 47(1):125-134.
- Matthew KM (1999). *The flora of the Palni Hills, South India*, Vol (1-4). The Rapinat Herbarium, Tiruchirappalli, India.
- Mukul SA, Uddin MB, Uddin MS, Khan MASA, Marzan B (2008). Protected areas of Bangladesh: current status and efficacy for biodiversity conservation. *Proceedings of the Pakistan Academy of Sciences* 45(2):59-68.
- Nawash O, Al-Assaf A, El-oqlah A, Omari M (2014). Floristic features, distribution, and ethnobotany of plants gathered and used by local people from the Mediterranean forest in Northern Jordan. *Ethnobotany Research and Applications* 12:385-396.
- Nawaz A, Hossain MM, Karim M, Khan M, Jahan M, Rahmatullah M (2009). An ethnobotanical survey of Jessore district in Khulna division, Bangladesh. *American-Eurasian Journal of Sustainable Agriculture* 3(2):238-243.
- Partha P (2014). Ethnobotany of the Laleng (Partha) Community in Bangladesh. *Journal of Pharmacognosy and Phytochemistry* 2(6):173-184.
- Partha P, Hossain AE (2007). Ethnobotanical investigation into the Mandi ethnic community in Bangladesh. *Bangladesh Journal of Plant Taxonomy* 14(2):129-145.
- Parthiban R, Vijayakumar S, Prabhu S, Yabesh JMEM (2016). Quantitative traditional knowledge of medicinal plants used to treat livestock diseases from Kudavasal taluk of Thiruvavur district, Tamil Nadu, India. *Revista Brasileira de Farmacognosia* 26(1):109-121. Available at: <http://dx.doi.org/10.1016/j.bjp.2015.07.016>
- Prain D (1903-04). *Bengal Plants*. Vols. (1- 2). Indian Reprint Edition 1963, M/s Bishen Singh Mahendra Pal Singh, Calcutta, India.
- Rahman AHM (2013). *Medico-Ethnobotany: A study on the tribal people of Rajshahi Division, Bangladesh*. Peak Journal of Medicinal Plants Research 1(1):1-8.
- Rahman MM (2004). Forest resources of Bangladesh with reference to conservation of biodiversity and wildlife in particular for poverty alleviation. In *Proceedings of the workshop on Forests for poverty reduction: Opportunities with clean development mechanism, environmental services and biodiversity*. Bangkok: FAO Regional Office for Asia and the Pacific (FAO-RAP), pp. 139-148.
- Rahmatullah M, Azam MNK, Malek I, Nasrin D, Jamal F, Rahman MA, Khatun Z, Jahan S, Seraj S, Jahan R (2012). An ethnomedicinal survey among the Marakh sect of the Garo tribe of Mymensingh district, Bangladesh. *International Journal of PharmTech Research* 4:141-149.
- Ray T, Mondal K (2018). Ethnobotanical Study of Munshiganj in Shyamnagar Upazilla, Satkhira, Bangladesh. *American-Eurasian Journal of Sustainable Agriculture* 12(1):22-31. Available at:



- <http://dx.doi.org/10.22587/aejsa.2018.12.1.4>
- Roy S, Uddin MZ, Hassan MA, Rahman MM (2008). Medico-botanical report on the Chakma community of Bangladesh. *Bangladesh Journal of Plant Taxonomy*, 15(1):67-72.
- Singh AK, Raghubanshi AS, Singh JS (2002). Medical ethnobotany of the tribals of Sonaghati of Sonbhadra district, Uttar Pradesh, India. *Journal of Ethnopharmacology* 81(1):31-41. Available at: [http://dx.doi.org/10.1016/S0378-8741\(02\)00028-4](http://dx.doi.org/10.1016/S0378-8741(02)00028-4)
- Stoffl RW, Halmo DB, Evans MJ, Olmsted JE (1990). Calculating the cultural significance of American Indian plants: Paiute and Shoshone ethnobotany at Yucca Mountain, Nevada. *American Anthropologist* 92:416-432. Available at: <http://dx.doi.org/10.1525/aa.1990.92.2.02a00100>
- Turner NJ (1988). The importance of a rose": Evaluating the cultural significance of plants in Thompson and Lillooet Interior Salish. *American Anthropologist* 90(2):272-290.
- Uddin MZ, Hassan MA, Rahman M, Arefin K (2012). Ethno-medico-botanical study in Lawachara National park, Bangladesh. *Bangladesh Journal of Botany* 41(1):97-104. Available at: <http://dx.doi.org/10.3329/bjb.v41i1.11087>
- Yusuf M, Begum J, Hoque MN, Chowdhury JU (2009). Medicinal Plants of Bangladesh. BCSIR, Chittagong, Bangladesh.