

## Review

## The genus *Eupatorium* L. (Asteraceae): A review of their antimicrobial activity

Antonio Carlos Nogueira Sobrinho<sup>1\*</sup>, Selene Maia de Morais<sup>1</sup>, Elnatan Bezerra de Souza<sup>2</sup> and Raquel Oliveira dos Santos Fontenelle<sup>2</sup>

<sup>1</sup>Postgraduate Program in Biotechnology-Renorbio, Center for Science and Technology, State University of Ceará, Fortaleza, Brazil.

<sup>2</sup>Course of Biological Sciences, Center for Agricultural Sciences and Biological Sciences, State University of Vale do Acaraú, Sobral, Brazil.

Received 30 November, 2016; Accepted 6 January, 2017

In recent years, the number of infectious diseases linked to the occurrence of bacterial and fungal resistance has increased, leading to extensive search for new drugs to treat these infections. Species of the Asteraceae family and the genus *Eupatorium*, have high biological potential and are used in folk medicine to treat various diseases. This review article presents the main phytochemical and biological characteristics of the Asteraceae family and the genus *Eupatorium s.l.*, whose antimicrobial activity is promising, especially antibacterial and antifungal activity. The current review was achieved using an organized search of the scientific data published on antimicrobial activity and phytochemical of the species of the genus *Eupatorium* using various databases, including PubMed, ScienceDirect, Scopus, Scielo, SciFinder and Google Scholar. The species of *Eupatorium* are rich in terpenes, phytosterols and sesquiterpene lactones, the latter being chemotaxonomic markers of the group, with broad anticancer, antiplasmodial and antimicrobial activity, making them promising for the development of new drugs. Various species of *Eupatorium* seems to hold great potential for in-depth investigation for antimicrobial activities. Many species have broad folk use, with scientific confirmation of its antimicrobial properties making these plants potential sources of safer and more effective treatments.

**Key words:** Compositae, *Eupatorium s.l.*, antimicrobial potential, ethnopharmacology, ethnobotany.

### INTRODUCTION

The family Asteraceae, belonging to the class of eudicotyledons, is one of the largest families of angiosperm plants. Its species are known for their therapeutic, cosmetic and aromatic properties and have great importance both from the environmental and economic standpoints. They have been widely studied for

their anti-inflammatory, antimicrobial, anticancer, antisyphilitic, antigonorrheal and insecticidal properties (Carrillo-Hormaza et al., 2015). Biological studies of its extracts, oils and constituents for production of phytochemicals have attracting growing interdisciplinary interest from the scientific community (Garcia-Sanchez et

\*Corresponding author. E-mail: caiosobrinho@yahoo.com.br.

al., 2011; Yang et al., 2004; Souza et al., 2007).

Among several genera of Asteraceae studied, *Eupatorium* L. *sensu lato* stands out. Numerous studies of its species have been performed and the majority showed the presence of terpenes, phytosterols and sesquiterpene lactones (Liu et al., 2015), the latter identified as promising sources for development of new drugs due to the antimicrobial, anticancer and antiplasmodial action (Albuquerque et al., 2010; Lang et al., 2002).

In a broader context, ethnobotanical and ethnopharmacological studies are very useful for investigation of new bioactive plant sources (Silambarasan and Ayyanar, 2015). Many traditional indigenous communities have longstanding contact with plant species having medicinal properties, with information being transmitted from generation to generation. This information is useful to the scientific community in studies of natural resources, as occurred in studies of *Eupatorium* *s.l.* (Silva et al., 2012; Paredes-Flores et al., 2007).

Chemical compounds with biological activity isolated from plants and used by the population through extracts are promising sources to prospect for new antimicrobial drugs. These substances may also act synergistically with other drugs, potentially improving the antibiotic action (Al-Fatimi et al., 2007). However, it is necessary to establish whether the traditional use of plants for the treatment of infectious diseases is supported by pharmacological action or merely based on folklore (Pessini et al., 2003).

One obstacle to the treatment of bacterial infections is the phenomena of resistance to antibacterial drugs, caused by the selection of resistant bacteria from exposure to chemotherapeutic agents, causing transfer of resistant gene fragments between bacterial strains and clonal spread of resistant strains among hospitalized patients (Spellberg et al., 2013). The exposure of antimicrobial compound can increase bacterial resistance also by selecting for mutation in genes that help microbes detoxify antibiotics. As for antibiotics, inadequate knowledge of the pharmacokinetic and pharmacodynamic properties, in addition to factors inherent to the patient, such as immune status and non-adherence to treatment, affect the efficiency of chemotherapy against pathogenic bacteria (Muller et al., 2015; Udy et al., 2013).

The use of natural extracts in the treatment of infectious diseases is promising, but studies on antimicrobial activity of different plant extracts should be expanded, including analysis of essential oils and their constituents in order to maximize the effect studied and to discover which chemical compound plays the main role in antimicrobial action (Danielli et al., 2013). In relation to antifungal drugs, natural products are widely used and the results are promising (Cragg and Newman, 2013). Due to the resistance of many pathogenic microorganisms, the search for new antimicrobial agents from plants is intense

(Alviano and Alviano, 2009). The relevance of these studies is based on the fact that infections of bacterial etiology, and in particular fungal infections, are aggravated in immunosuppressed individuals, due to therapy against cancer and AIDS and suppression of the immune response to prevent rejection of transplanted organs (Miceli et al., 2011).

For a comprehensive literature overview, the current review was achieved by using an organized search of the scientific data published on antimicrobial activity and phytochemical, focusing on the antibacterial and antifungal activities of the species of the genus *Eupatorium*. The search was conducted using the keyword search term "Eupatorium antimicrobial activity". The searches were carried out using various databases, including PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>), Science Direct (<http://www.sciencedirect.com/>), Scopus (<http://www.scopus.com/>), Scielo (<http://www.scielo.org/>), SciFinder (<https://scifinder.cas.org/>) and Google Scholar (<http://www.scholar.google.com/>).

Based on these considerations, in the present paper, the antibacterial and antifungal potential of species of the genus *Eupatorium* *s.l.*, were analyzed highlighting also biological properties of the Asteraceae family.

## ASTERACEAE FAMILY

Asteraceae (Compositae) includes about 1600 genera and 25000 species (Petacci et al., 2012) belonging to 17 tribes and three subfamilies. It is considered by some authors as the largest family of eudicotyledonous angiosperm plants (Nakajima and Semir, 2001; Hattori and Nakajima, 2008). The Asterales order is formed by the Calyceraceae, Menyanthaceae, Goodeniaceae, Campanulaceae and Asteraceae (Judd et al., 2009). This family has been studied intensely over the past 25 years regarding morphology, anatomy, ontogeny and ecology as well as phytochemical, cytogenetic and macromolecular aspects (Hind and Beentje, 1996).

Asteraceae are for the most part herbs, subshrubs or shrubs, trees or vines (Roque and Bautista, 2008). Its representatives of greatest importance are *Baccharis trimera* (Less) DC. (gorse), *Matricaria chamomilla* L. (camomile), *Cynara scolimus* L. (artichoke), *Vernonia condensata* Baker ("boldo-da-bahia") and *Arnica montana* L. (true arnica). These species are widely used and marketed as herbal products, suggesting species of the family Asteraceae have significant potential as a source of bioactive compounds (Maia et al., 2010; Mello et al., 2008). They are also sources of edible oils (*Helianthus annuus* L., sunflower), pesticides and latex, as well as being eaten as leafy vegetables (e.g., *Lactuca sativa* L., lettuce) and used as ornamental plants (Wu et al., 2006). Phytochemical studies of *Eupatorium* have identified the presence of sesquiterpene lactones, flavonoids, triterpenes, benzofuran compounds,

pyrrolizidine alkaloids, chromene and steroids (Liu et al., 2015; Zhang et al., 2008; Albuquerque et al., 2004; Lang et al., 2001).

## GENUS *EUPATORIUM*

The genus *Eupatorium* s.l. (tribe Eupatorieae, subtribe Eupatoriinae) was one of the most representative of the family Asteraceae, containing around 1200 species, however a detailed survey conducted by King and Robinson (1987) found that about 1010 species would be reclassified to different new genera. When interpreted in its traditional concept, this genus form a polyphyletic group. *Eupatorium* s.l. have been a highly artificial concept, tending to include all members of the tribe with a pappus of numerous capillary bristles, five ribs on the achene, and another appendage as long as or longer than wide (King and Robinson, 1987). This genus has been restructured and was fragmented into 80 genera. Currently, the genus *Eupatorium sensu stricto* containing about 45 species, distributed primarily by arctic-tertiary region (King and Robinson, 1987; Bremer, 1994).

The species of this genus are used in folk medicine in different parts of the world (Albuquerque et al., 2010), like Mexico, where over 1000 medicinal plants have been used for over 400 years in folk medicine, many are in the genus *Eupatorium* (Garcia-Sanchez et al., 2011).

Ethnopharmacological studies of medicinal plants are important for the development of new drugs (Silambarasan and Ayyanar, 2015). The main goal is to find new bioactive substances for disease control with minimal side effects (Rodríguez and Carlini, 2003). In this context, studies have been conducted of biological activities such as toxic activity of essential oils (Albuquerque et al., 2004) and plant extracts (Rozto et al., 2008) against the larvae of *Aedes aegypti* and antiplasmodial activity against *Plasmodium falciparum*, a protozoan that causes malaria (Lang et al., 2002).

In particular, the literature contains numerous studies of the genus *Eupatorium* with use of plant extracts and essential oils against strains of bacteria and fungi that infect animals and plants, for development of new drugs for veterinary and human use and new herbicides. Studies were found from 1948 to 2016 with *Eupatorium* species. *E. adenophorum*, *E. odoratum* and *E. triplinerve* are the species with the most work reporting their antimicrobial potential (Table 1).

### Antimicrobial activity of *Eupatorium* s.l.

Most studies of the antimicrobial activity involved species of Asia and Europe, with fewer studies of South American plants. This is counterintuitive, given that although countries like China, Malaysia, India, Zaire, Indonesia and Madagascar can be considered megadiverse (Sandes and Di Blasi, 2000), species of *Eupatorium* are particularly

diverse in tropical and subtropical areas of the Americas (Zhang et al., 2008).

For antibacterial action, tests have been conducted on activity against a wide range of microorganisms, both Gram-positive and negative bacteria. For investigations of plant extracts, the most commonly used solvents are methanol, ethanol, ethyl acetate and n-hexane, used in studies the aerial parts, branches, twigs and leaves. Nevertheless, there are reports of the use of roots to obtain the essential oil of *E. adenophorum* Spreng. (Ahluwalia et al., 2014) and the aqueous extract of *E. purpureum* L. (Carlson et al., 1948; Pates Madsen, 1955), in both cases having antibacterial action.

Referring to antifungal bioassays, the studies focused on *Eupatorium* generally also have involved tests against bacteria. The tested fungi include filamentous fungi, dermatophytes and yeasts. Tests with dermatophytes, filamentous fungi that cause dermatophytosis are promising for the genus, with reports of activity for *Eupatorium arnottianum* Griseb. (Muschiatti et al., 2005), *Eupatorium aschenbornianum* S. Schauer (García et al., 2003 Rios et al., 2003), *Eupatorium buniifolium* Hook. and Arn. (Muschiatti et al., 2005), *Eupatorium japonicum* Thunb. (Inouye et al., 2006), *Eupatorium laciniatum* Kitam. (Inouye et al., 2006) and *Eupatorium odoratum* L. (Taylor et al., 1996). In all these studies, extracts of plant material were used. There are few reports in the literature on the use of essential oils for susceptibility testing. As for the fungi tested, of the three genera of dermatophytes (*Trichophyton*, *Microsporum* and *Epidermophyton*), only two showed sensitivity and growth inhibition (*Trichophyton* and *Microsporum*), highlighting species *T. rubrum* and *T. mentagrophytes* in the genus *Trichophyton*.

### *Eupatorium adenophorum* Spreng.

The species *E. adenophorum* [syn. *Ageratina adenophora* (Spreng.) R.M.King & H.Rob.], native to Mexico, is now widely distributed in the world (King and Robinson, 1970) and is used in folk medicine as an antimicrobial, antiseptic, analgesic and antipyretic (Bhattarai and Shrestha, 2009). It has also been shown to have molluscicide potential against *Oncomelania hupensis*, the intermediate host of *Schistosoma japonicum* (Zou et al., 2009). The essential oil extracted from the flowers, and roots was tested and showed inhibitory activity for the strains of *Xanthomonas oryzae*, *Erwinia chrysanthemi*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*. Assays were performed using the disk diffusion and agar dilution methods (Ahluwalia et al., 2014). The essential oil extracted from the aerial parts was also investigated against strains of *Arthrobacter protophormiae*, *Escherichia coli*, *Micrococcus luteus*, *Rhodococcus rhodochrous* and *Staphylococcus aureus*. Assays were performed by broth dilution and the results showed growth inhibition for all strains tested (Kurade et

**Table 1.** Species of the genus *Eupatorium* s.l. with antimicrobial activity.

Scientific name	Part used	Biological preparation	Bioactive compounds	Biological activity	Reference
<i>Eupatorium adenophorum</i> Spreng.	I, L, St, R	EO, OsE	Terpenoids	Antibacterial activity: Gram-positive and Gram-negative bacteria Antifungal activity: filamentous fungi	Bhattarai and Shrestha, 2009; Kurade et al., 2010; Kundu et al., 2013; Ahluwalia et al., 2014; Chauhan et al., 2015
<i>Eupatorium altissimum</i> L.	AP	DE	-	Antibacterial activity: Gram-positive and Gram-negative bacteria	Carlson et al., 1948; Cantrell et al., 1998
<i>Eupatorium arnottianum</i> Griseb.	AP	ME	-	Antifungal activity: filamentous fungi	MuschietTI et al., 2005
<i>Eupatorium aromaticum</i> L.	L	AqE	-	Antibacterial activity: Gram-positive bacteria	Pates and Madsen, 1955
<i>Eupatorium aschenbornianum</i> S. Schauer.	AP	HE, ME	Benzofurans compounds and phytosterols	Antifungal activity: yeast and filamentous fungi	García et al., 2003, Rios et al., 2003
<i>Eupatorium ayapana</i> Vent.	L	PE	Steroids, coumarins, tannins and saponins	Antibacterial activity: Gram-positive and Gram-negative bacteria	Gupta et al., 2002
<i>Eupatorium ballotifolium</i> Kunth	AP	EO	Terpenoids	Antifungal activity: filamentous fungi	Sobrinho et al., 2016
<i>Eupatorium buniifolium</i> Hook. ex Hook. & Arn.	AP	ME	-	Antifungal activity: filamentous fungi	MuschietTI et al., 2005
<i>Eupatorium cannabinum</i> L.	AP	OsE	-	Antibacterial activity: Gram-positive and Gram-negative bacteria	Freerksen and Bönicke, 1951; Senatore et al., 2001
<i>Eupatorium capillifolium</i> (Lam.) Small.	AP	EO	Terpenoids	Antifungal activity: filamentous fungi	Senatore et al., 2001; Tabanca et al., 2010
<i>Eupatorium fortunei</i> Turcz.	AP	AqE	-	Antibacterial activity: Gram-negative bacteria	Li et al., 2010
<i>Eupatorium glutinosum</i> Lam.	AP	HE, ME	Terpenoids	Antibacterial activity: Gram-positive and Gram-negative bacteria	Lopez et al., 2001; El-Seedi et al., 2002
<i>Eupatorium havanense</i> Kunth	AP	DE	-	Antibacterial activity: Gram-positive bacteria	Cantrell et al., 1998
<i>Eupatorium intermedium</i>	F	EO	Terpenoids	Antibacterial activity: Gram-positive and Gram-negative bacteria	Czaikoski et al., 2015
<i>Eupatorium inulaefolium</i> Kunth.	L	AqE, EE	-	Antibacterial activity: Gram-positive and Gram-negative bacteria	Pérez and Anesini, 1994; Sanabria-Galindo et al., 1998; Álvarez et al., 2005
<i>Eupatorium japonicum</i> Thunb.	L, F	EA	Terpenoids	Antifungal activity: filamentous fungi	Inouye et al., 2006
<i>Eupatorium laciniatum</i> Kitam.	L, F	EA	Terpenoids and coumarin	Antifungal activity: filamentous fungi	Inouye et al., 2006

Table 1. Cont'd

<i>Eupatorium laevigatum</i> Lam.	L	HE, EE, ME	Alkaloids, steroids, phenols, tannins and flavonoids	Antibacterial activity: Gram-positive and Gram-negative bacteria	Schmidt et al., 2009; Fabri et al., 2011
<i>Eupatorium lindleyanum</i> DC	AP	EE, ME, EA	-	Antibacterial activity: Gram-positive and Gram-negative bacteria	Ji et al., 2008
<i>Eupatorium maculatum</i> L.	AP, S	EE	-	Antibacterial activity: Gram-positive bacteria	Bishop and Macdonald, 1951; Borchardt et al., 2008
<i>Eupatorium odoratum</i> L.	L, AP, F	EE, ME	Polyphenols	Antibacterial activity: Gram-positive and Gram-negative bacteria Antifungal activity: filamentous fungi	Cáceres et al., 1994; Taylor et al., 1996; Suksamrarn et al., 2004; Chomnawang et al., 2005; Molina-Salinas et al., 2007; Gautam et al., 2007; Chomnawang et al., 2009; Sharma et al., 2013; Patil and Kamble, 2015
<i>Eupatorium patens</i> D. Don ex Hook. & Arn.	L	EO	Terpenoids	Antibacterial activity: Gram-positive bacteria	Bailac et al., 2000
<i>Eupatorium perfoliatum</i> L.	L, F	EE, PE	-	Antibacterial activity: Gram-positive and Gram-negative bacteria Antifungal activity: yeast	Carlson et al., 1948; Bishop and Macdonald, 1951; Madsen and Pates, 1952; Liegey, 1953; Frisby et al., 1953; Borchardt et al., 2008
<i>Eupatorium purpureum</i> L.	R, St, L	AqE, EE	-	Antibacterial activity: Gram-positive and Gram-negative bacteria	Carlson et al., 1948; Pates and Madsen, 1955
<i>Eupatorium rugosum</i> Houtt.	L, S	AqE	-	Antibacterial activity: Gram-positive and Gram-negative bacteria	Frisby et al., 1954
<i>Eupatorium salvia</i> Colla	AP	RE	Resinous exudate	Antibacterial activity: Gram-positive bacteria	Urzua et al., 1998
<i>Eupatorium serratum</i> Spreng.	L	ME, EA	Tannins, flavonoids, steroids and alkaloids	Antibacterial activity: Gram-positive bacteria	Desoti et al., 2011
<i>Eupatorium tashiroi</i> Hayata.	AP	GE	-	Antibacterial activity: Gram-positive bacteria	Chen et al., 1989
<i>Eupatorium triplinerve</i> Vahl	St, L, AP	EO, CE	Terpenoids	Antibacterial activity: Gram-positive and Gram-negative bacteria Antifungal activity: filamentous fungi	Rahman and Junaid, 2008; Begum et al., 2010; Unnikrishnan et al., 2014; Sugumar et al., 2015
<i>Eupatorium urticaefolium</i> Reichard	AP	AE	-	Antibacterial activity: Gram-positive and Gram-negative bacteria	Carlson et al., 1948

I- inflorescence; L- leaves; R- root; AP- aerial part; F- flower; St- stems; S- seed; ME- methanol extract; CE- chloroform extract; HE- hexane extract; EE- ethanol extract; PE- petroleum ether extract; DE- dichloromethane extract; EA- ethyl acetate extract; AqE- aqueous extract; AE- acid extract; GE- gross extract; OsE- organic solvents extract; RE- resinous exudate; EO- essential oil.

al., 2010).

Extracts from fresh leaves using organic

solvents were tested against 15 bacterial strains by the agar diffusion method. The organic extracts

showed antibacterial activity against *Salmonella paratyphi*, *Salmonella* spp., *Staphylococcus aureus*,

*Staphylococcus* spp., *Bacillus cereus*, *Bacillus subtilis*, *Bacillus thuringiensis*, *Enterobacter aerogenes*, *Proteus mirabilis* and *Proteus* spp. while the aqueous extract showed activity against *P. aeruginosa*, *E. coli*, *S. aureus*, *Staphylococcus* spp., *Citrobacter freundii*, *Proteus* spp., *B. subtilis*, *E. aerogenes*, *Salmonella* spp., *S. paratyphi* and *B. thuringiensis* (Bhattarai and Shrestha, 2009).

Cadinene derivatives extracted from leaves of *E. adenophorum* showed antifungal activity against four plant pathogenic fungi strains: *Rhizoctonia solani*, *Sclerotium rolfsii*, *Fusarium oxysporum* and *Macrophomina phaseolina*. Cadinene sesquiterpenes were isolated by column chromatography and preparative thin layer chromatography. Less polar cadinene sesquiterpenes were active against *S. rolfsii* and polar cadinene derivatives with mono or dihydroxy groups were more inhibitory towards *R. solani* (Kundu et al., 2013).

Ahluwalia et al. (2014) analyzed the antifungal action of the essential oil of *E. adenophorum* against five pathogenic fungi: *Sclerotium rolfsii*, *Macrophomina phaseolina*, *Rhizoctonia solani*, *Pythium debaryanum* and *Fusarium oxysporum*. They found the strongest inhibitory activity against *S. rolfsii*. Regarding toxicity, they evaluated the phytotoxic action of the essential oil, which inhibited the growth and seed germination of *Phalaris minor* Retz. and *Triticum aestivum* L., used for phytotoxicity assays in a dose-dependent concentration.

In another study, the essential oil from aerial parts exhibited moderated antifungal activity against *Fusarium oxysporum* by disk diffusion method (Chauhan et al., 2015). Twenty-two components were identified in the composition with an abundance of monoterpenes p-cymene (14.56%) followed by phellandrene (12.25%), camphene (10.42%), bornyl acetate (9.76%) and  $\delta$ -2-carene (5.39%).

### ***Eupatorium altissimum* L.**

In a study on the species collected in the state of Louisiana in the United States, a dichloromethane extract was tested against *Mycobacterium tuberculosis* bacterium, causative agent of most cases of tuberculosis (Cantrell et al., 1998). In a bioprospecting study of 550 plants collected in the states of Ohio and Oregon in the United States, action was reported on the extract of aerial parts of *E. altissimum* against strains of *S. aureus* and *E. coli* (Carlson et al., 1948).

### ***Eupatorium arnottianum* Griseb.**

*E. arnottianum* [syn. *Chromolaena arnottiana* (Griseb.) R.M. King & H. Rob.] is an herb native to Latin America, common in the North-East and Centre of Argentina and South of Bolivia, traditionally used to treat stomach ailments (Clavin et al., 2007), against asthma, bronchitis,

colds and topically in plasters for bone fractures and dislocations by healers from the bolivian altiplano (Giraoult, 1987), in addition to having anti-inflammatory and analgesic properties (Muschiatti et al., 2005).

A methanol extract of the aerial parts showed antifungal activity against the dermatophytes *Microsporum gypseum*, *Trichophyton rubrum* and *Trichophyton mentagrophytes* according to broth microdilution tests (Muschiatti et al., 2005).

### ***Eupatorium aromaticum* L.**

*E. aromaticum* [syn. *Ageratina aromatica* (L.) Spach.] is a perennial herb, medium-sized, native to America, common in the southeastern United States, known as "Small-leaved white snakeroot" or "Lesser snakeroot" (Blackwell and Mcmillan, 2013). The ethyl extract of the leaves of *E. aromaticum* showed antibacterial activity against Gram-positive bacteria (Pates and Madsen, 1955).


### ***Eupatorium aschenbornianum* S. Schauer.**

*E. aschenbornianum* [syn. *Ageratina pichinchensis* (Kunth) R.M. King & H. Rob.] is a plant traditionally used in Mexico to treat skin lesions, mouth ulcers and tumors (Rios et al., 2003). Studies of hexane and methanol extracts obtained from the aerial parts point to the antifungal potential of *E. aschenbornianum* against the dermatophytes, *T. mentagrophytes*, *T. rubrum*, the yeast *C. albicans* and the filamentous fungus *Aspergillus niger* (Rios et al., 2003; García et al., 2003).

Yeasts belonging to the genus *Candida* are involved in the etiology of candidiasis (Netea et al., 2015). Studies on the species *E. aschenbornianum* showed activity against *C. albicans*, using hexane and methanol extracts of the aerial parts (Figure 1). Benzofurans compounds (5-acetyl-3 $\beta$ -angeloyloxy-2 $\beta$ -(1-hydroxyisopropyl)-2,3-dihydrobenzofurane and 5-acetyl-3 $\beta$ -angeloyloxy-2 $\beta$ -(1-hydroxyisopropyl)-6-methoxy-2,3-dihydrobenzofurane) and phytosterols were isolated (Rios et al., 2003; García et al., 2003).

### ***Eupatorium ayapana* Vent.**

The leaves of *E. ayapana* [syn. *Ayapana triplinervis* (Vahl) R.M. King & H. Rob.] are used in folk medicine as a heart stimulant, laxative and anticoagulant (Gupta et al., 2002). In rainforests of South America, its leaves are used in decoctions, infusions, tea and baths as tonic, stimulant, sedative, febrifuge and anti-inflammatory, especially in Brazilian folk medicine (Melo et al., 2013). The first preliminary studies of this species were performed by Oswaldo Gonçalves de Lima, then director



**Figure 1.** Structures of two benzofurans compounds from aerial parts of *Eupatorium aschenbornianum*. **A.** 5-acetyl-3 $\beta$ -angeloyloxy-2 $\beta$ -(1-hydroxyisopropyl)-2,3-dihydrobenzofurane. **B.** 5-acetyl-3 $\beta$ -angeloyloxy-2 $\beta$ -(1-hydroxyisopropyl)-6-methoxy-2,3-dihydrobenzofurane.

of the Antibiotics Institute of the University at Recife, Brazil (Lima, 1963).

The phytochemical study of petroleum ether extract showed presence of steroids, coumarins, tannins and saponins. For the species *E. ayapana*, the petroleum ether extract inhibited growth of bacteria *B. subtilis*, *Staphylococcus epidermidis*, *S. aureus*, *M. luteus*, *E. coli*, *P. aeruginosa*, *Salmonella typhi*, *Shigella* sp., *Vibrio cholerae* and *Vibrio parahaemolyticus*. The extract also inhibited the growth of *A. niger*, *A. flavus*, *Alternaria solani* and *Fusarium solani*, pathogenic fungi causing black mold in fruits and vegetables, deterioration by mycotoxins in grains, alternaria spot in solanaceous plants and root and stem rot, respectively (Gupta et al., 2002).

### *Eupatorium ballotifolium* Kunth

*Eupatorium ballotifolium* Kunth (syn. *Lourteigia ballotaefolia* (Kunth) R.M. King & H. Rob.) popularly known in Brazil by many names, such as “maria-preta”, “maria-preta-verdadeira” and “picão-roxo” is a perennial herbaceous species whose vertical growth ranges from 40 to 80 cm (Cardoso et al., 2013).

The essential oil extracted from the aerial parts, was tested by broth microdilution method and showed antifungal activity against strains fungal dermatophytes, *Trichophyton rubrum*. The chromatographic analysis showed 25 components, accounting for 93.84%, among them mono- and sesquiterpenes, with prevalence of sesquiterpenes (Sobrinho et al., 2016).

The modulatory activity assay was performed to determine the combined effect of the essential oil with the ketoconazole, standard antifungal drug, against strains of *T. rubrum* by the checkerboard technique. The

combination of ketoconazole with essential oil reduced the MICs for both strains of *T. rubrum* indicating a synergistic effect (Sobrinho et al., 2016).

### *Eupatorium buniifolium* Hook. ex Hook. & Arn.


*E. buniifolium* [syn. *Acanthostyles buniifolius* (Hook. ex Hook. & Arn.) R.M. King & H. Rob.] is a shrubby species widely used in folk medicine, such as to treat rheumatic pains and liver problems with tea obtained by decoction. Preparations from this plant are also used as disinfectants (Ríos et al., 1993).

The methanol extract obtained from the aerial parts was used to investigate antifungal activity against strains of the dermatophytes, *M. gypseum* (MIC = 250  $\mu$ g/mL), *T. rubrum* (MIC = 100  $\mu$ g/mL) and *T. mentagrophytes* (MIC = 250  $\mu$ g/mL). The assays were performed by the broth microdilution method (Muschiatti et al., 2005).

### *Eupatorium cannabinum* L.

*E. cannabinum* is a perennial herbaceous species distributed in the northern hemisphere, especially in Europe and North America (Senatore et al., 2001). A study of the essential oil of *E. cannabinum* subsp. *cannabinum* L. obtained from the aerial parts identified high percentage of terpenoids, such as germacrene, monocyclic sesquiterpene, found in large amounts. The oil showed antibacterial activity against strains of Gram-positive bacteria (Senatore et al., 2001).

In another investigation, the extract of the aerial parts showed antimicrobial activity against strains of Gram-positive and negative bacteria, the main ones tested being *S. typhi*, *Micrococcus aureus*, *B. subtilis* and



**Figure 2.** Structures of two isolated compounds from leaves of *E. glutinosum*. A. 15-hydroxy-7-labden-17-oic acid: R = H; B. 15-acetoxy-7-labden-17-oic acid: R = Ac.

*Proteus* spp. (Freerksen and Bönicke, 1951).

### ***Eupatorium capillifolium* (Lam.) Small. ex Porter & Britton**

*E. capillifolium* is a perennial herbaceous species found in North America. It is used as food for livestock in the southeastern United States when other fodder is lacking, but it is generally considered a weed (Sellers et al., 2009).

Studies of the effect of plant extracts against pathogenic fungi are important to find new agents against fungi causing plant diseases in agricultural crops. Tabanca et al. (2010) investigated, by contact bioautography, the antifungal activity of the essential oil of *E. capillifolium*, which inhibited growth of the fungus *Colletotrichum acutatum*, the cause of strawberry anthracnose. The main constituents were thymol methyl ether (36.3%), 2,5-dimethoxy-p-cymene (20.8%) and myrcene (15.7%).

### ***Eupatorium fortunei* Turcz.**

*E. fortunei* is a traditional Chinese medicinal herb, common in the northeastern of China, used in folk medicine to treat many diseases (Liu et al., 1992). In a study of bacteria that produce volatile sulfur compounds (VSCs) that cause halitosis, Li et al. (2010) investigated extracts of 40 plants found in China, of which 14 inhibited bacterial growth and production of VSCs. The main compounds found were hydrogen sulfide, dimethyl sulfide and methylmercaptan. Among the 14 plants that were active in the screening was *E. fortunei*, which was used to obtain the aqueous extract.

### ***Eupatorium glutinosum* Lam.**

This species *E. glutinosum* [syn. *Aristeguietia glutinosa*

(Lam.) R.M. King & H. Rob.] is widely used in folk medicine, with preparations obtained by decoction used as antimicrobial, antirheumatic and antidiarrheal agents and to heal stomach ulcers (El-Seedi et al., 2002).

The methanol extract of leaves of *E. glutinosum* presented antibacterial activity against strains of *Mycobacterium phlei*, *B. subtilis* and *S. aureus*, by the disk diffusion method (Lopez et al., 2001). The antibacterial activity of the hexane extract of leaves and stems and of two isolated compounds (15-hydroxy-7-labden-17-oic acid and 15-acetoxy-7-labden-17-oic acid) were investigated by diffusion in agar (Figure 2). The tests showed activity against *S. aureus*, *B. cereus*, *E. coli* and *P. aeruginosa* (El-Seedi et al., 2002).

### ***Eupatorium havanense* Kunth**

*E. havanense* [syn. *Ageratina havanensis* (Kunth) R.M. King & H. Rob.] is a native shrub found in North America (Zamudio and Villanueva, 2011) and Central America (Córdoba et al., 1995) known as "Havana snakeroot". Previous study identified the flavone sakuranetin (Wollenweber and Dietz, 1981) and flavonols, flavones and flavonoid glucosides (Yu et al., 1987).

In an antimicrobial prospecting study of plants of the Americas, the dichloromethane extract of leaves of *E. havanense*, collected in the state of Tamaulipas, Mexico, inhibited the growth of the pathogenic microorganisms *M. tuberculosis* and *M. avium*, the latter an opportunistic pathogen related to infections in immunocompromised individuals (Cantrell et al., 1998).

### ***Eupatorium intermedium* DC.**

*E. intermedium* [syn. *Grazielia intermedia* (DC.) R.M. King and H. Rob.] is a branchy shrub, native to southern Brazil with upright growth between 1.0 to 1.5 m high and it is densely leafy until close to the inflorescences, which are composed of white flowers (Czaikoski et al., 2015).

The extracts obtained by supercritical extraction from the flowers with scCO<sub>2</sub>, propane and petroleum ether as solvents were tested and showed antibacterial activity against two Gram-positive bacterial strains, *S. aureus* and *L. monocytogenes*, and two Gram-negative bacterial strains, *E. coli* and *S. typhimurium*. The assays were performed using the disk diffusion method (Czai Koski et al., 2015). The extracts were effective against all bacteria strains, particularly the extract obtained with scCO<sub>2</sub> and compressed propane.

The essential oil was tested against Gram-negative bacterial strains using the agar well method. The main constituents of essential oil were  $\alpha$ -pinene, sabinene,  $\beta$ -pinene, limonene, (*E*)-caryophyllene, germacrene D, spathulenol and caryophyllene oxide (Czai Koski et al., 2016).

### ***Eupatorium inulaefolium* Kunth.**

*E. inulaefolium* [syn. *Austroeupatorium inulaefolium* (Kunth) R.M. King & H. Rob.] is a species found in Latin America, whose common name in Argentina is "sanalotodo" or "yerba de Santa María". This species is used in folk medicine for treatment of skin infections due to its antimicrobial properties (Lancelle et al., 2009) and throat diseases (Álvarez et al., 2005).

In a study of 122 plant species from 54 families, Pérez and Anesini (1994) investigated the antibacterial activity against the microorganism *S. typhi*, causative agent of typhoid fever. Of the plants tested in the initial screening, *E. inulaefolium* showed inhibitory activity against the growth of *S. typhi*. In an investigation of the *in vitro* antimicrobial activity of Colombian angiosperms, the inhibitory activity against the microorganisms *S. aureus* and *B. subtilis*, was investigated at a concentration of 3 mg/mL, by the agar diffusion method (Sanabria-Galindo et al., 1998). In a third study, the ethyl extract of the leaves of *E. inulaefolium* were tested against the strains of *S. aureus*, *E. coli* and *P. aeruginosa*. The authors only observed growth inhibition against *S. aureus*, at a concentration of 50 mg/mL (Álvarez et al., 2005).

### ***Eupatorium japonicum* Thunb.**

*E. japonicum* [syn. *Eupatorium chinense* L.] is a perennial herb, native to Japan, Korea and northeastern area of China. It is a medicinal herb used in folk medicine as a decoction for the treatment of measles, rheumatic bone pains, colds and cough (Roeder, 2000).

The ethyl acetate extract of the leaves and flowers of *E. japonicum* was used to investigate the antifungal activity against the dermatophyte *T. mentagrophytes*, the most common cause of *Tinea pedis*. At the concentration of 0.25 mg/mL, the authors observed an inhibition zone of 11 mm (Inouye et al., 2006).

### ***Eupatorium laciniatum* Kitam.**

*E. laciniatum* [syn. *Eupatorium chinense* L.] an annual herbal plant, native to Asia is distributed in the south of China, used in Chinese folk medicine for the treatment of cold, snakebite and inflammation (Yang et al., 2005).

The antidermatophytic activity of this plant species was tested against the fungus, *T. mentagrophytes*, using the ethyl acetate extract of the leaves and flowers at a concentration of 0.25 mg/mL. The extract caused an inhibition zone of 12 mm (Inouye et al., 2006).

### ***Eupatorium laevigatum* Lam.**

*E. laevigatum* [syn. *Chromolaena laevigata* (Lam.) R.M. King & H. Rob.], native to South America is widely distributed in the South part of Brazil and Argentina, whose common name in Brazil is "erva-de-santa-maria" or "camara", is used in folk medicine for wound healing and as an antifungal (Maia et al., 2002). In Argentina, it is known as "caá-hú" or "doctorcito" and used in folk medicine as an anticefalalgic, analgesic, cathartic, emenagogue and purgative agent (Clavin et al., 2000).

The ethanol and hexane extracts of the leaves were tested against *B. subtilis*, by the agar diffusion method, with moderate results for inhibition of bacterial growth by both solvents (Schmidt et al., 2009). In a chemical and antimicrobial prospecting study, the methanol extract of the leaves of *E. laevigatum* showed the presence of alkaloids, steroids, phenols, tannins and flavonoids. The antimicrobial activity was performed by the broth microdilution method against strains of Gram-positive and negative bacteria, showing inhibitory activity against *P. aeruginosa*, *B. cereus*, *Shigella sonnei* and *Salmonella enterica* sorovar *typhimurium* (Fabri et al., 2011).

### ***Eupatorium lindleyanum* DC.**

*E. lindleyanum* is a perennial herbaceous plant, native to China, known as "Yemazhui" and used in folk medicine to relieve fever, remove toxic substances, treat coughs, promote urination, and lower blood pressure (Xia et al., 2004). In traditional Chinese medicine, its aerial part is used as an antipyretic and detoxicant (Yang et al., 2010).

The extracts of the aerial parts obtained by use of ethanol, methanol and ethyl acetate were tested by the broth dilution method against eight bacterial strains: four Gram-positive (*B. subtilis*, *S. aureus*, *B. cereus* and *E. faecium*) and four Gram-negative (*E. coli*, *S. typhimurium*, *P. vulgaris* and *Pseudomonas fluorescens*). The extracts inhibited growth of all strains tested, with the best results for *B. subtilis*, *S. aureus* and *B. cereus* (Ji et al., 2008).

### ***Eupatorium maculatum* L.**

Native to United States, *E. maculatum* known as "Joe-

pye weed" is widely found in moist places, especially in calcareous soils (Goebel et al., 2012). The first report of antimicrobial activity of *E. maculatum* dates back to the work of Bishop and MacDonald (1951), who reported that the ethanol extract of the aerial parts showed activity against strains of Gram-positive bacteria.

In prospective study with 336 native species of the states of Minnesota and Wisconsin in the United States, ethanol extracts of leaves and stems of *E. maculatum* were tested against strains of *S. aureus*, *E. coli*, *P. aeruginosa* and *C. albicans*, presenting significant action only against *S. aureus*. The assays were performed using the disk diffusion method (Borchardt et al., 2008).

### ***Eupatorium odoratum* L.**

*E. odoratum* [syn. *Chromolaena odorata* (L.) R.M. King & H. Rob.] is a free standing shrub which is mostly distributed in America, tropical Asia and West Africa. It is widely used in traditional medicine as immunomodulator, antispasmodic, hepatoprotective, antiprotozoal, antidiabetic, antihypertensive, anti-inflammatory (Umukoro and Ashorobi, 2006). *E. odoratum* also can be used in treatment of skin diseases, dysentery, malaria fever, tooth ache and diabetes (Omoregie et al., 2014). In Mexican folk medicine, it is used for chest complaints and pulmonary affections, known as "crucita" (Molina-Salinas et al., 2007).

Various studies of *E. odoratum* have investigated the plant's antimicrobial activity against bacteria that cause various infectious diseases. Cáceres et al. (1995) observed activity against the bacterium *Neisseria gonorrhoeae*, which causes gonorrhea, with an excellent growth inhibition zone.

Four flavanones extracted from flowers of *E. odoratum* exhibited antibacterial activity against *Mycobacterium tuberculosis*. Flavanones were isolated by silica gel chromatography and analyzed with infrared spectroscopy and nuclear magnetic resonance (Suksamrarn et al., 2004). Chomnawang et al. (2005) tested the antiacne activity (crude extracts with chloroform and ethylacetate) against strains of *Propionibacterium acnes*, *S. epidermidis*, while Molina-Salinas et al. (2007) investigated the antibacterial activity (methanol extract; leaves, flowers and roots) against *Streptococcus pneumoniae* and Chomnawang et al. (2009) tested it against methicillin-resistant *S. aureus*.

Methanol extract of the aerial parts of *E. odoratum* inhibited the growth of *M. phlei* by disk diffusion method, with inhibitory activity at concentrations of 300 mg/mL (Gautam et al., 2007). In another study using the aerial parts, the methanol extract inhibited microbial growth of *B. subtilis*, methicillin-resistant *S. aureus*, methicillin-sensitive *S. aureus*, *T. mentagrophytes* and *M. phlei*, in the last case only when exposed to UV radiation (Taylor et al., 1996). The essential oil from the dried leaves of *E. odoratum* showed *in vitro* antibacterial activity against

both Gram-positive and negative bacterial strains (Sharma et al., 2013). The major components analyzed by gas chromatography-mass spectrometry (GC-MS) were geijerene (25.10%), germacrene D (20.27%), trans- $\beta$ -caryophyllene (10.04%),  $\alpha$ -pinene (9.64%) and  $\beta$ -pinene, (4.85%).

In a screening study, the aqueous leaf extract of *E. odoratum* was tested for their fungitoxicity against spores of *Uromyces vignae* and showed the better result for the inhibition of spore germination (Patil and Kamble, 2015).

### ***Eupatorium patens* D. Don ex Hook. & Arn.**

*E. patens* [syn. *Austrobrickellia patens* (D. Don ex Hook. & Arn.) R.M. King & H. Rob.] is a densely branched shrub that is distributed from Bolivia and Paraguay to Patagonia in Argentina, known by the common names of "acancio" or "bejuco" (Cabrera and Correa, 1971).

The essential oil of the leaves was tested against the microorganisms *S. aureus*, *E. coli*, *B. subtilis*, *C. albicans* and *A. niger* by the agar diffusion method. Only the strains of *B. subtilis* showed significant growth inhibition zones after incubation for 24 h (Bailac et al., 2000).

### ***Eupatorium perfoliatum* L.**

Native to North America, *E. perfoliatum* is a medicinal herb, known as "boneset" or "thoroughwort", used in folk medicine by the native inhabitants for the treatment of fever and flu (Maas et al., 2008). The first record in the literature on the antimicrobial potential of *E. perfoliatum* was performed by Carlson et al. (1948). The authors found that the ether extract obtained from the aerial parts inhibited the growth of *S. aureus*. Later studies investigated extracts from leaves and stems using organic solvents (ethanol, ether and acetone) and water, demonstrating antimicrobial activity against *C. albicans*, Gram-positive and Gram-negative bacteria and *Mycobacterium* spp. (Bishop and MacDonald, 1951; Madsen and Pates, 1952; Liegey, 1953; Frisby et al., 1953).

In prospective study, the ethanol extract of the leaves and flowers of *E. perfoliatum* was tested against strains of *S. aureus*, *E. coli*, *P. aeruginosa* and *C. albicans*. The extract only had significant action against *S. aureus*, with inhibition zones of 11 and 7 mm for extracts of the leaves and inflorescences, respectively. The assays were performed using the disk diffusion method (Borchardt et al., 2008).

### ***Eupatorium purpureum* L.**

*E. purpureum* is a wildflower perennial plant, native to the eastern United States, known as "joe-pye weed", "sweet joe-pye weed", and "sweetscented joe-pye weed"

(Sabanadzovic et al., 2010). It is used in traditional medicine for rheumatoid arthritis and several other disease conditions (Habtemariam, 1998).

The saline extract of the stems of *E. purpureum* L. showed antibacterial activity against *S. aureus* and *E. coli*, by the agar diffusion method (Carlson et al., 1948). The saline extract of the leaves was also investigated against *S. aureus* and *E. coli*, with positive results by the agar diffusion method (Pates and Madsen, 1955).

### ***Eupatorium rugosum* Houtt.**

*E. rugosum* [syn. *Ageratina altissima* (L.) R.M. King & H. Rob.] is a perennial herb commonly found in the midwestern and eastern United States, known as “white snakeroot” (Lee et al., 2012). The saline extract of the stems of *E. rugosum* showed antibacterial activity against strains of Gram-positive bacteria by the agar diffusion method (Frisby et al., 1954).

### ***Eupatorium salvia* Colla**

*E. salvia* [syn. *Aristeguietia salvia* (Colla) R.M. King & H. Rob.] is a species used in folk Chilean medicine, known as “sálvia macho”, “pegajosa” or “pega-pega” (Hoffmann, 1995). The resinous exudate, 7-hydroxy-8(17)-labden-15-oic acid (salvic acid) and 7-acetoxy-8(17)-labden-15-oic acid (salvic acid acetate), isolated from *E. salvia*, were tested against five bacterial strains, with activity against all tested microorganisms: *S. aureus*, *B. cereus*, *B. subtilis*, *M. luteus* and *Clavibacter michiganensis* subsp. *Michiganensis*. These diterpenoids do not affect Gram-negative bacteria and the activity of these compounds on filamentous fungi has not been reported (Urzua et al., 1998).

### ***Eupatorium serratum* Spreng.**

*E. serratum* [syn. *Grazielia serrata* (Spreng.) R.M. King & H. Rob.], popularly known as “erva-milagrosa” or (“miraculous herb”) in Brazil, is used in folk medicine to treat snake bites, as an analgesic, healing agent, antimicrobial agent and for the treatment of stomach problems, liver disorders, diabetes, cancer and bronchial asthma (Desoti et al., 2011).

The hexane, methanol and ethyl acetate extracts obtained from fresh leaves of *E. serratum* were tested against *S. aureus*, *M. luteus*, *E. coli* and *S. typhi*. The best inhibitory halo results were produced by the methanol extract against *M. luteus* (Desoti et al., 2011).

### ***Eupatorium tashiroi* Hayata**

*E. tashiroi*, native to Asia is a wild herb which has been

used in a folk medicine for treating edema and hemoptysis in Taiwan (Wu et al., 1985). The aqueous extract of the entire plant was tested by the agar diffusion method against two serotypes of *S. mutans*, bacteria associated with the development of dental caries. The extracts inhibited bacterial growth of both serotypes (Chen et al., 1989).

### ***Eupatorium triplinerve* Vahl**

*E. triplinerve* [syn. *Ayapana triplinervis* (Vahl) R.M. King & H. Rob.] commonly found in Asia is an erect perennial herb with narrow lanceolate leaves and large number of pedicelled flower-heads at the top of the branch, known as “ayapana” (Selvamangai and Bhaskar, 2012). The species was introduced in Indian as ornamental species. In Indian, folk medicine is used as a stimulant, tonic, laxative and for the treatment of yellow fever (Unnikrishnan et al., 2014).

The antimicrobial activities of the essential oil obtained from the leaves and the stem and thymohydroquinone dimethylether were evaluated using disk diffusion assay against strains of Gram-positive and negative bacteria, showing inhibitory activity against *P. aeruginosa*, *S. aureus*, *K. pneumoniae* and *E. coli*. The essential oil showed antifungal activity against fungal strains of *Penicillium chrysogenum* and *C. albicans*; however thymohydroquinone dimethylether only inhibited the growth of *P. chrysogenum* (Unnikrishnan et al., 2014).

The chloroform extract of leaves exhibited antibacterial activity by the broth microdilution and disk diffusion methods against strains of Gram-positive and negative bacteria *Bacillus megaterium*, *B. subtilis*, *B. cereus*, *S. aureus*, *E. coli*, *S. dysenteriae*, *S. sonnei*, *S. typhi*, *S. paratyphi* and *P. aeruginosa*, that exhibited the largest zone of inhibition against *Vibrio* (22 mm in diameter with 1000 µg/disc extract). The antifungal activity was performed with filamentous fungi *Alternaria alternata*, *Curvularia lunata*, *Colletotrichum corchori*, *Fusarium equiseti*, *Macrophomina phaseolina* and *Botryodiplodia theobromae*, with highest inhibition of fungal radial mycelial growth against *C. corchori*, 73.5% with 100 µg extract/ml medium (Rahman and Junaid, 2008).

In others studies, the essential oil from aerial parts and fresh leaves exhibited moderate antibacterial activity against *B. megaterium*, *B. subtilis*, *B. cereus*, *S. aureus*, *E. coli*, *V. cholerae*, *S. dysenteriae*, *S. sonnei*, *S. typhi*, *Pseudomonas* sp. and *S. paratyphi*. The essential oil demonstrated strong antifungal activities against *A. alternata*, *C. lunata*, *B. theobromae*, *C. corchori*, *F. equiseti* and *M. phaseolina* (Begum et al., 2010; Sugumar et al., 2015).

### ***Eupatorium urticaefolium* Reichard**

*E. urticaefolium* [syn. *Ageratina altissima* (L.) R.M. King &

H. Rob.] is a species native to North America, it is commonly found throughout the eastern half of continent. Ingestion of the plant causes a disease called trembles in livestock (Lee et al., 2010). The ethyl acetate extract of the aerial parts of *E. urticaefolium* inhibited the growth of *S. aureus* and *E. coli*, by the agar diffusion method (Carlson et al., 1948).

## Conclusion

Thirty *Eupatorium* species were identified, some already reclassified to other genera. The aerial parts are the part of plant with more studies and so far the best source for anti-microbial product, being the essential oils and fixed compounds the main natural products. Most of the studies were carried by the broth microdilution and disk diffusion methods, according to Clinical and Laboratory Standards Institute (CLSI).

Microbial infections caused by bacteria and fungi are a growing public health problem, and increasingly common in routine medicine. Outbreaks of infections by methicillin-resistant *S. aureus* in hospital and social environments have been reported in several regions of the world. Regarding fungal infections, the search for new more efficient therapies is particularly urgent considering that the available antifungal chemicals are much less than antibacterial drugs and also typically have more side effects, besides the possible development of fungal resistance.

This review shows that the Asteraceae family, and especially the genus *Eupatorium* s.l., contains many species with antimicrobial activity, making these plants potential sources of safer and more effective treatments. In short, *Eupatorium* s.l. contains a diverse array of species with antimicrobial potential. New studies involving chemical and biological bioprospecting are necessary to develop effective and less toxic herbal products.

## Conflict of interest

The authors declare that there is no conflict of interest.

## ACKNOWLEDGEMENTS

The authors are grateful to the Postgraduate Program in Biotechnology-Renorbio of State University of Ceará for the financial support of CAPES, the Microbiology Laboratory of Vale do Acaraú State University and the Herbarium Prof. Francisco José de Abreu Matos (HUVA) of Vale do Acaraú State University.

## REFERENCES

Ahluwalia V, Sisodia R, Walia S, Sati OP, Kumar J, Kundu A (2014). Chemical analysis of essential oils of *Eupatorium adenophorum* and

- their antimicrobial, antioxidant and phytotoxic properties. *J. Pest. Sci.* 87:341-349.
- Albuquerque MRJR, Silveira ER, Uchôa DEA, Lemos TLG, Souza EB, Santiago GMP, Pessoa ODL (2004). Chemical Composition and Larvicidal Activity of the Essential Oils from *Eupatorium betonicaeforme* (DC.) Baker (Asteraceae). *J. Agric. Food Chem.* 52:6708-6711.
- Albuquerque MRJR, Santos HS, Souza EB, Silva RM, Pessoa ODL, Braz-Filho R, Costa SMO (2010). Composição química volátil e não-volátil de *Eupatorium ballotifolium* Kunth, Asteraceae. *Rev. Bras. Farmacogn.* 20:615-620.
- Al-fatimi M, Wurster M, Schröder G, Lindequist U (2007). Antioxidant, antimicrobial and cytotoxic activities of selected medicinal plants from Yemen. *J. Ethnopharmacol.* 111:657-666.
- Álvarez M, Isaza G, Echeverry HM (2005). Efecto antibacteriano *in vitro* de *Austro eupatorium inulaefolium* HBK (Salvia amarga) y *Ludwigia polygonoides* HBK (Clavo de laguna). *Biosalud*, 14:46-55.
- Alviano DS, Alviano CS (2009). Plant extracts: search for new alternatives to treat microbial diseases. *Curr. Pharm. Biotechnol.* 10:106-121.
- Bailac PN, Dellacasa AD, Bernasconi HO, Firpo NH, Ponzi MI (2000). Composicion del aceite esencial y actividad antimicrobiana de *Eupatorium patens*. *Bol. Soc. Chil. Quím.* 45:207-211.
- Begum J, Bhuiyan MNJ, Taznin T (2010). Chemical composition and antimicrobial activity of essential oil from *Eupatorium triplinerve* Vahl. aerial parts. *Asian J. Microbiol. Biotech. Environ. Sci.* 12:543-547.
- Bhattarai N, Shrestha G (2009). Antibacterial and antifungal effect of *Eupatorium adenophorum* Spreng against bacterial and fungal isolates. *Nepal J. Sci. Technol.* 10:91-95.
- Bishop CJ, Macdonald RE (1951). A survey of higher plants for antibacterial substances. *Can. J. Bot.* 29:260-269.
- Blackwell AH, Mcmillan PD (2013). Collected in South Carolina 1704-1707: The plants of Joseph Lord. *Phytoneuron* 59:1-15.
- Borchardt JR, Wyse DL, Sheaffer CC, Kauppi KL, Fulcher RG, Ehlke NJ, Biesboer DD, Bey RF (2008). Antimicrobial activity of native and naturalized plants of Minnesota and Wisconsin. *J. Med. Plants Res.* 2:098-110.
- Bremer K (1994). Asteraceae, cladistics and classification. Portland, United States: Timber Press.
- Cabrera AL, Correa MN (1971). Flora Patagónica: parte 7. Compositae. *Col. Cient. INTA*, 8:17.
- Cáceres A, Menéndez H, Méndez E, Cohobón E, Samayoa BE, Jauregui E, Peralta E, Carrillo G (1995). Antigonorrhoeal activity of plants used in Guatemala for the treatment of sexually transmitted diseases. *J. Ethnopharmacol.* 48:85-88.
- Cantrell CL, Fischer NH, Urbatsch L, Mcguire MS, Franzblau SG (1998). Antimycobacterial crude plant extracts from South, Central, and North America. *Phytomedicine* 5:137-145.
- Cardoso AD, Viana AES, Barbosa RP, Teixeira PRG, Júnior NSC, Fogaça JJNL (2013). Levantamento fitossociológico de plantas daninhas na cultura da mandioca em Vitória da Conquista, Bahia. *Biosci. J.* 29:1130-1140.
- Carlson HJ, Douglas HG, Robertson J (1948). Antibacterial substances separated from plants. *J. Bacteriol.* 5:241-248.
- Carrillo-Hormaza L, Mora C, Alvarez R, Alzate F, Osorio E (2015). Chemical composition and antibacterial activity against *Enterobacter cloacae* of essential oils from Asteraceae species growing in the Páramos of Colombia. *Ind. Crops Prod.* 77:108-115.
- Chauhan N, Haider SZ, Lohani H, Godbole S, Gwari G, Bhandari U (2015). Chemical Composition and Antifungal Activity of Essential Oil of *Cymbopogon distans* (Nees ex Steud.) W. Watson, *Eupatorium adenophorum* Spreng and *Lantana camara* L. grown in Uttarakhand (India). *J. Biol. Active Prod. Nat.* 5:234-240.
- Chen CP, Lin CC, Namba T (1989). Screening of Taiwanese crude drugs for antibacterial activity against *Streptococcus mutans*. *J. Ethnopharmacol.* 27:285-295.
- Chomnawang MT, Surassmo S, Nukoolkarn VS, Gritsanapan W (2005). Antimicrobial effects of Thai medicinal plants against acne-inducing bacteria. *J. Ethnopharmacol.* 101:330-333.
- Chomnawang MT, Surassmo S, Wongsariya K, Bunyapraphatsara N (2009). Antibacterial Activity of Thai Medicinal Plants against Methicillin-resistant *Staphylococcus aureus*. *Fitoterapia* 80:102-104.

- Clavin ML, Gorzalczy S, Miño J, Kadarian C, Martino V, Ferraro G, Acevedo C (2000). Antinociceptive effect of some Argentine medicinal species of *Eupatorium*. *Phytother. Res.* 14:275-277.
- Clavin M, Gorzalczy S, Macho A, Munoz E, Ferraro G, Acevedo C, Martino V (2007). Anti-inflammatory activity of flavonoids from *Eupatorium arnotianum*. *J. Ethnopharmacol.* 112:585-589.
- Córdoba HF, Báez MB, Domínguez RS (1995). Tamizaje de alcaloides y saponinas en plantas que crecen en Cuba. III: Sierra del Rosario. *Rev. Cubana Enfermer.* 11:23-24.
- Cragg GM, Newman DJ (2013). Natural products: a continuing source of novel drug leads. *Biochim. Biophys. Acta-Gen. Subjects* 1830:3670-3695.
- Czaikoski K, Mesomo MC, Scheer AP, Santa ORD, Queiroga CL, Corazza ML (2015). Kinetics, composition and biological activity of *Eupatorium intermedium* flower extracts obtained from scCO<sub>2</sub> and compressed propane. *J. Supercrit Fluids* 97:145-153.
- Czaikoski K, Mesomo MC, Scheer ADP, Queiroga CL, Deschamps C, Corazza ML (2016). Chemical composition and biological activity of *Eupatorium intermedium* essential oil. *J. Essent. Oil Res.* pp.1-8.
- Danielli LJ, Reis M, Bianchini M, Camargo GS, Bordignon SAL, Guerreiro IK, Fuentesria A, Apel MA (2013). Antidermatophytic activity of volatile oil and nanoemulsion of *Stenachaenium megapotamicum* (Spreng.) Baker. *Ind. Crops Prod.* 50:23-28.
- Desoti VC, Maldaner CL, Carletto MS, Heinz AA, Coelho MS, Piati D, Tiunan TS (2011). Triagem fitoquímica e avaliação das atividades antimicrobiana e citotóxica de plantas medicinais nativas da região oeste do estado do Paraná. *Arq. Ciênc. Saúde Unipar* 15:3-13.
- El-seedi HR, Ohara T, Sata N, Nishiyama S (2002). Antimicrobial diterpenoids from *Eupatorium glutinosum* (Asteraceae). *J. Ethnopharmacol.* 81:293-296.
- Fabri RL, Nogueira MS, Dutra LB, Bouzada MLM, Scio E (2011). Potencial antioxidante e antimicrobiano de espécies da família Asteraceae. *Rev. Bras. Plantas Med.* 13:183-189.
- Freerksen E, Bönicke R (1951). Über antibakterielle Prinzipien in höheren Pflanzen I. *Med. Microbiol. Immunol.* 132:417-449.
- Frisby A, Gottshall RY, Jennings JC, Lucas EH (1953). The occurrence of antibacterial substances in seed plants with special reference to *Mycobacterium tuberculosis* (Third Report). *Mich. State Univ. Agric. Exp. Stn. Q. Bull.* 35:392-404.
- Frisby A, Gottshall RY, Jennings JC, Lucas EH (1954). The occurrence of antibacterial substances in seed plants with special reference to *Mycobacterium tuberculosis* (Fourth Report). *Mich. State Univ. Agric. Exp. Stn. Q. Bull.* 36:477-488.
- García-Sánchez E, López CBR, Torres RENR, Pacheco MMM (2011). A revision de *Eupatorium* (Compositae: Eupatorieae) from Michoacan. *Phyton.* 80:139-146.
- García VMN, Gonzales A, Fuentes M, Aviles M, Rios MY, Zepeba G, Rojas MG (2003). Antifungal activities of nine traditional Mexican medicinal plants. *J. Ethnopharmacol.* 87:85-88.
- Gautam R, Saklani A, Jachak SM (2007). Indian medicinal plants as a source of antimycobacterial agents. *J. Ethnopharmacol.* 110:200-234.
- Giraoult L (1987). Kallawaya. Curanderos itinerantes de los Andes. UNICEF. OPS. OMS, La Paz, Bolivia P. 226.
- Goebel PC, Palik BJ, Pregitzer KS (2012). Structure and composition of riparian forests in an old-growth northern hardwood-hemlock watershed. *Forest Ecol. Manag.* 280:52-61.
- Gupta M, Mazumder UK, Chaudhuri I, Chaudhuri RK, Bose P, Bhattacharya S, Manikandan L, Patra S (2002). Antimicrobial activity of *Eupatorium ayapana*. *Fitoterapia* 73:168-170.
- Habtemariam S (1998). Extract of gravel root (rhizome of *Eupatorium purpureum*) inhibits integrin-dependent U937 cell adhesion. *Phytother. Res.* 12:422-426.
- Hattori EKO, Nakajima JN (2008). A família Asteraceae na Estação de Pesquisa e Desenvolvimento Ambiental Galheiro, Perdizes, Minas Gerais, Brasil. *Rodriguésia* 59:687-749.
- Hind DJN, Beentje HJ (1996). Compositae: Systematics, Proceedings of the International Compositae Conference, Kew, 1994. *R. Bot. Gard.* 1:621-626.
- Hoffmann A (1995). Flora silvestre de Chile, zona central. Ed. Fund. Claudio Gay.
- Inouye S, Uchida K, Abe S (2006). Volatile composition and vapour activity against *Trichophyton mentagrophytes* of 36 aromatic herbs cultivated in Chichibu district in Japan. *Int. J. Aromather.* 16:159-168.
- Ji LL, Luo YM, Yan GL (2008). Studies on the antimicrobial activities of extracts from *Eupatorium lindleyanum* DC against food spoilage and food-borne pathogens. *Food Control* 19:995-1001.
- Judd WS, Campbell CS, Kellogg EA, Stevens PF, Donoghue MJ (2009). Sistemática Vegetal: um enfoque filogenético. Porto Alegre, Brazil: Artmed. P 612.
- King RM, Robinson H (1970). Studies in the Eupatorieae (Compositae). XXXII. A new genus Neocuatrecasia. *Phytologia* 19:208.
- King RM, Robinson H (1987). The genera of the Eupatorieae (Asteraceae). Monographs in Systematic Botany. Missouri Bot. Gard. 22 St. Louis. Missouri Bot. Gard. 22: 581 p.
- Kundu A, Saha S, Walia S, Shakil NA, Kumar J, Annapurna K (2013). Cadinene sesquiterpenes from *Eupatorium adenophorum* and their antifungal activity. *J. Environ. Sci. Health. B.* 48:516-522.
- Kurade NP, Jaitak V, Kaul VK, Sharma OP (2010). Chemical composition and antibacterial activity of essential oils of *Lantana camara*, *Ageratum houstonianum* and *Eupatorium adenophorum*. *Pharm. Biol.* 48:539-544.
- Lancelle HG, Giordano OS, Sosa ME, Tonn CE (2009). Chemical composition of four essential oils from *Eupatorium* spp. biological activities toward *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Rev. Soc. Entomol. Argent.* 68:329-338.
- Lang G, Passreiter CM, Medinilla BE, Castillo JJ, Witte L (2001). Non-toxic pyrrolizidine alkaloids from *Eupatorium semialatum*. *Biochem. Syst. Ecol.* 29:143-147.
- Lang G, Passreiter CM, Wright CW, Filipowicz NH, Addae-Kyereme J, Medinilla BE, Castillo JJ (2002). Antiplasmodial activities of sesquiterpene lactones from *Eupatorium semialatum*. *Z Naturforsch C. Bio. Sci.* 57:282-286.
- Lee ST, Davis TZ, Gardner DR, Colegate SM, Cook D, Green BT, Meyerholtz KA, Wilson CR, Stegelmeier BL, Evans TJ (2010). Tremetone and structurally related compounds in white snakeroot (*Ageratina altissima*): a plant associated with trembles and milk sickness. *J. Agric. Food Chem.* 58:8560-8565.
- Lee ST, Davis TZ, Cook D, Stegelmeier BL (2012). Evaluation of drying methods and toxicity of rayless goldenrod (*Isocoma pluriflora*) and white snakeroot (*Ageratina altissima*) in goats. *J. Agric. Food Chem.* 60:4849-4853.
- Li M-Y, Wang J, Xu ZT (2010). Effect of a variety of Chinese herbs and an herb-containing dentifrice on volatile sulfur compounds associated with halitosis: An in vitro analysis. *Curr. Ther. Res.* 71:129-140.
- Liegey FW (1953). Antibiotic properties of plants common to Cattaraugus County. *St. Bonav. Univ. Sci. Stud.* 15:40-62.
- Lima GO (1963). Antibiotic richness of Latin-American plants particularly from Brazil and Mexico. *Plant Foods Hum. Nutr.* 9:257-285.
- Liu K, Roeder E, Chen THL, Xiu XJ (1992). Pyrrolizidine alkaloids from *Eupatorium fortunei*. *Phytochemistry* 31:2573-2574.
- Liu PY, Liu D, Li WH, Zhao T, Sauriol F, Gu YC, Shi QW, Zhang ML (2015). Chemical Constituents of Plants from the Genus *Eupatorium* (1904-2014). *Chem. Biodivers.* 12:1481-1515.
- Lopez A, Hudson JB, Towers GHN (2001). Antiviral and antimicrobial activities of Colombian medicinal plants. *J. Ethnopharmacol.* 77:189-196.
- Madsen GC, Pates AL (1952). Occurrence of antimicrobial substances in chlorophyllous plants growing in Florida. *Bot. Gaz.* 113:293-300.
- Maia JGS, Zoghbi MDG, Andrade EHA, Silva MHL, Luz AIR, Silva JD (2002). Essential oils composition of *Eupatorium* species growing wild in the Amazon. *Biochem. Syst. Ecol.* 30:1071-1077.
- Maia AIV, Torres MCM, Pessoa ODL, Menezes JESA, Costa SMO, Nogueira VLR, Melo VMM, Souza EB, Cavalcante MGB, Albuquerque MRJR (2010). Óleos essenciais das folhas de *Vernonia remotiflora* e *Vernonia brasiliensis*: composição química e atividade biológica. *Quim. Nova* 33:584-586.
- Maas M, Peteret F, Hensel A (2008). Caffeic acid derivatives from *Eupatorium perfoliatum* L. *Molecules* 14:36-45.
- Melo AS, Monteiro MC, Silva JB, Oliveira FR, Vieira JLF, Andrade MA, Baetas AC, Sakai JT, Ferreira FA, Sousa PJC, Maia CFS (2013). Antinociceptive, neurobehavioral and antioxidant effects of *Eupatorium triplinerve* Vahl on rats. *J. Ethnopharmacol.* 147:293-301.
- Mello JRB, Mello FB, Langeloh A (2008). Toxicidade Pré-Clinica de

- Fitoterápico Contendo *Gentiana lutea*, *Rheum palmatum*, *Aloe ferox*, *Cynara scolymus*, *Atropa belladonna*, *Peumus boldus* e *Baccharis trimera* em Coelhos Nova Zelândia. *Lat. Am. J. Pharm.* 27:752-756.
- Miceli MH, Díaz JA, Lee SA (2011). Emerging opportunistic yeast infections. *Lancet. Infect. Dis.* 11:142-151.
- Molina-Salinas GM, Pérez-López A, Becerril-Montes P, Salazar-Aranda R, Said-Fernández S, Torres NW (2007). Evaluation of the flora of Northern Mexico for in vitro antimicrobial and antituberculosis activity. *J. Ethnopharmacol.* 109:435-441.
- Muller AE, Theuretzbacher U, Mouton JW (2015). Use of old antibiotics now and in the future from a pharmacokinetic/pharmacodynamic perspective. *Clin. Microbiol. Infect.* 21:881-885.
- Muschiatti L, Derita M, Sülsen V, Muñoz JD, Ferraro G, Zacchino S, Martino V (2005). In vitro antifungal assay of traditional Argentine medicinal plants. *J. Ethnopharmacol.* 102:233-238.
- Nakajima JN, Semir J (2001). Asteraceae do Parque Nacional da Serra da Canastra, Minas Gerais, Brasil. *Rev. Bra. Bot.* 24:471-478.
- Netea MG, Joosten LA, Van der Meer JW, Kullberg BJ, Van de Veerdonk FL (2015). Immune defence against *Candida* fungal infections. *Nat. Rev. Immunol.* 15:630-642.
- Omorieg ES, Oriakhi K, Oikeh EI, Okugbo OT, Akpobire D (2014). Comparative Study of Phenolic Content and Antioxidant Activity of Leaf Extracts of *Alstonia boonei* and *Eupatorium odoratum*. *Niger. J. Basic. Appl. Sci.* 22:91-97.
- Paredes-Flores M, Saade RL, Aranda PDD (2007). Estudio etnobotánico de Zapotillán Salinas, Puebla. *Acta Bot. Mex.* 79: 13-61.
- Pates AL, Madsen GC (1955). Occurrence of antimicrobial substances in chlorophyllous plants growing in Florida. *II Bot. Gaz.* 116: 250-261.
- Patil BJ, Kamble SK (2015). Antifungal activity of some plant extracts on Spore germination of *Uromyces vignae* Barclay. *Bionano. Frontier.* 8: 110-112.
- Pérez C, Anesini C (1994). In vitro antibacterial activity of Argentine folk medicinal plants against *Salmonella typhi*. *J. Ethnopharmacol.* 44: 41-46.
- Pessini GL, Holetz FB, Sanches NR, Cortez DAG, Dias-Filho BP, Nakamura CV (2003). Avaliação da atividade antibacteriana e antifúngica de extratos de plantas utilizados na medicina popular. *Rev. Bras. Farmacogn.* 13: 21-24.
- Petacci F, Tavares WS, Freitas SS, Teles AM, Serrão JE, Zanuncio JC (2012). Phytochemistry and quantification of polyphenols in extracts of the Asteraceae weeds from Diamantina, Minas Gerais State, Brazil. *Planta Daninha* 30:9-15.
- Rahman MS, Junaid M (2008). Antimicrobial activity of leaf extracts of *Eupatorium triplinerve* Vahl. against some human pathogenic bacteria and phytopathogenic fungi. *Bangladesh. J. Bot.* 37:89-92.
- Rios MY, Aguilar-Guadarrama AB, Navarro V (2003). Two new benzofuranes from *Eupatorium aschenbornianum* and their antimicrobial activity. *Planta Med.* 69: 967-970.
- Ríos J, Condori L, Muñoz S, Loayza I, Soriano J (1993). The extraction of the essential oil of romerillo. In: Valorisation de la biomasse végétale par les produits naturels. *Actes Coll. Chicoutimi*, 22(25):111-117.
- Rodríguez E, Carlini ELA (2003). Levantamento etnofarmacológico realizado entre um grupo de quilombolas do Brasil. *Arq. Bras. Fitomed. Cientif.* 1:80-87.
- Roeder E (2000). Medicinal plants in China containing pyrrolizidine alkaloids. *Pharmazie* 55:711-726.
- Roque N, Bautista H (2008). Asteraceae: caracterização e morfologia floral. Salvador, Brazil: EDUFBA.
- Rozo A, Zapata C, Bello FJ (2008). Evaluación del efecto tóxico de extractos de *Eupatorium microphyllum* L.F. (Asteraceae) sobre larvas de *Aedes aegypti* (Diptera: Culicidae) en condiciones de laboratorio. *Rev. Cienc. Salud (Bogotá)* 6:64-73.
- Sabanadzovic S, Ingram DM, Lawrence AM (2010). First Report of Tobacco ringspot virus in Joe-Pye Weed (*Eupatorium purpureum*) in Mississippi. *Plant Dis.* 94:126-126.
- Sanabria-Galindo A, Ruiz AM, Moreno AL (1998). Actividad antimicrobiana in vitro de angiospermas colombianas. *Rev. Colomb. Cienc. Quim. Farm.* 27:47-51.
- Sandes ARR, Di Blasi G (2000). Biodiversidade e diversidade química e genética. *Biotecnolog. Cienc. Desenvolv.* 13:28-32.
- Schmidt C, Fronza M, Goettert M, Geller F, Luik S, Flores EMM, Bittencourt CF, Zanetti GD, Heinzmann BM, Laufer S, Merfort I (2009). Biological studies on Brazilian plants used in wound healing. *J. Ethnopharmacol.* 122:523-532.
- Sellers BA, Ferrell JA, Macdonald GE, Kline WN (2009). Dogfennel (*Eupatorium capillifolium*) size at application affects herbicide efficacy. *Weed Technol.* 23:247-250.
- Selvamangai G, Bhaskar A (2012). GC-MS analysis of phytochemicals in the methanolic extract of *Eupatorium triplinerve*. *Asian Pac. J. Trop. Biomed.* 2:1329-1332.
- Senatore F, De Fusco R, Napolitano F (2001). *Eupatorium cannabinum* L. ssp. *cannabinum* (Asteraceae) essential oil: chemical composition and antibacterial activity. *J. Essent. Oil. Res.* 13:463-466.
- Sharma K, Saikia AK, Sharma H, Sahariah BJ, Deka S, Das B (2013). Chemical Composition and Antimicrobial Study of Essential Oil from the Leaves of *Eupatorium odoratum* Linn. From Upper Assam region. *J. Essent. Oil. Bear. PI* 16:482-488.
- Silambarasan R, Ayyanar M (2015). An ethnobotanical study of medicinal plants in Palamalai region of Eastern Ghats, India. *J. Ethnopharmacol.* 172:162-178.
- Silva SMFQ, Pinheiro SMB, Queiroz MVF, Pranchevicius MC, Castro JGD, Perim MC, Carreiro SC (2012). Atividade in vitro de extratos brutos de duas espécies vegetais do cerrado sobre leveduras do gênero *Candida*. *Ciênc. Saúde Coletiva* 17:1649-1656.
- Sobrinho ACN, Souza EB, Rocha MFG, Albuquerque MRJR, Bandeira PN, Santos HS, Pereira R, Morais SM, Fontenelle ROS, Cavalcante CSP (2016). Cytotoxicity, antifungal and antioxidant activities of the essential oil from *Eupatorium ballotifolium* Kunth (Asteraceae). *Afr. J. Pharm. Pharmacol.* 10:346-355.
- Souza TJT, Apel MA, Bordignon S, Matzenbacher NI, Zuanazzi AS, Henriques AT (2007). Composição química e atividade antioxidante do óleo volátil de *Eupatorium polystachyum* DC. *Rev. Bras. Farmacogn.* 17:368-372.
- Spellberg B, Bartlett JG, Gilbert DN (2013). The future of antibiotics and resistance. *N. Engl. J. Med.* 368:299-302.
- Sugumar N, Karthikeyan S, Gowdhami T (2015). Chemical composition and antimicrobial activity of essential oil from *Eupatorium triplinerve* Vahl. aerial parts. *Int. Lett. Nat. Sci.* 4:14-21.
- Suksamram A, Chotipong A, Suavansri T, Boongird S, Timsuksai P, Vimuttipong S, Chuaynugul A (2004). Antimicrobial activity and cytotoxicity of flavonoids from the flowers of *Chromolaena odorata*. *Arch. Pharm. Res.* 27:507-511.
- Tabanca N, Bernier UR, Tsikolia M, Becnel JJ, Sampson B, Werle C, Demirci B, Baser KHC, Blythe EK, Pounders C, Wedge DE (2010). *Eupatorium capillifolium* essential oil: chemical composition, antifungal activity, and insecticidal activity. *Nat. Prod. Commun.* 5:1409-1415.
- Taylor RSL, Edel F, Manandhar NP, Towers GHN (1996). Antimicrobial activities of southern Nepalese medicinal plants. *J. Ethnopharmacol.* 50:97-102.
- Udy AA, Roberts JA, Lipman J (2013). Clinical implications of antibiotic pharmacokinetic principles in the critically ill. *Intensive Care Med.* 39:2070-2082.
- Umukoro S, Ashorobi RB (2006). Evaluation of the anti-inflammatory and membrane-stabilizing effects of *Eupatorium odoratum*. *Int. J. Pharmacol.* 2:509-512.
- Unnikrishnan PK, Varughese T, Sreedhar S, Balan N, Balachandran I, Shree ABR (2014). Study on *Eupatorium triplinerve* Vahl from South India, A Rich Source for Thymohydroquinone dimethylether and its Antimicrobial Activity. *J. Essent. Oil. Bear. PI* 17:652-657.
- Urzua A, Caroli M, Vasquez L, Mendonza L, Wilkens M, Tojo E (1998). Antimicrobial study of the resinous exudate and of diterpenoids isolated from *Eupatorium salvia* (Asteraceae). *J. Ethnopharmacol.* 62:251-254.
- Wollenweber E, Dietz VH (1981). Occurrence and distribution of free flavonoid aglycones in plants. *Phytochemistry* 205:869-932.
- Wu TS, Niwa M, Furukawa H, Kuoh GS (1985). Eupatriol, a new monoterpene from *Eupatorium tashiroi* HAYATA. *Chem. Pharm. Bull.* 33: 4005-4006.
- Wu QX, Shi YP, Jia ZJ (2006). Eudesmane sesquiterpenoids from the Asteraceae family. *Nat. Prod. Rep.* 23:699-734.
- Xia J, Wang GL, Wei F, Lin RC (2004). Chemical constituents of *Eupatorium lindleyanum*. *Chin. Tradit. Herb Drug* 35:855-856.
- Yang SP, Huo J, Wang Y, Lou LG, Yue JM (2004). Cytotoxic

- Sesquiterpenoids from *Eupatorium chinense*. J. Nat. Prod. 67:638-643.
- Yang SP, Cheng JG, Huo J, Jiang HL, Chen KX, Yue JM (2005). Seven new sesquiterpene lactones from *Eupatorium chinense*. Chin. J. Chem. 23:1530-1536.
- Yang NY, Duan JA, Shang EX, Tian LJ (2010). Analysis of sesquiterpene lactones in *Eupatorium lindleyanum* by HPLC-PDA-ESI-MS/MS. Phytochem. Anal. 21:144-149.
- Yu S, Fang N, Mabry TJ (1987). Flavonoids from *Ageratina havanensis*. Rev. Latinoam. Quím. 18:20-22.
- Zamudio S, Villanueva RG (2011). La diversidad vegetal del estado de Guanajuato. In Flora del Bajío y de Regiones Adyacentes. P 101.
- Zhang ML, Wu M, Zhang JJ, Irwin D, Gu YC, Shi QW (2008). Chemical Constituents of Plants from the Genus *Eupatorium*. Chem. Biodivers. 5:40-55.
- Zou FC, Duan G, Xie YJ, Zhou Y, Dong GD, Lin RQ, Zhu XQ (2009). Molluscicidal activity of the plant *Eupatorium adenophorum* against *Oncomelania hupensis*, the intermediate host snail of *Schistosoma japonicum*. Ann. Trop. Med. Parasitol. 103:549-553.