

*Full Length Research Paper*

# Comparative study of wild chamomile plants from the north-west of Morocco: Bioactive components and total antioxidant activity

Amina EL MIHYAOUI<sup>1,2\*</sup>, María Emilia Candela CASTILLO<sup>1</sup>, Antonio CANO<sup>1</sup>, Josefa HERNANDEZ-RUIZ<sup>1</sup>, Ahmed LAMARTI<sup>2</sup> and Marino Bañon ARNAO<sup>1</sup>

<sup>1</sup>Department of Plant Biology (Plant Physiology), Faculty of Biology, University of Murcia, 30100-Murcia, Spain.

<sup>2</sup>Laboratory of Plant Biotechnology, Department of Biology, Faculty of Sciences, Abdelmalek Essaadi University, Tetouan, Morocco.

Received 23 June, 2021; Accepted 26 July, 2021

Morocco presents numerous plants due its localization between the Mediterranean Sea and Atlantic Ocean and the variability of its mountainous terrain, with a rich range of medicinal plants. *Cladanthus mixtus* (L.) Oberpr. & Vogt and *Matricaria chamomilla* L., two Asteraceae (Compositae) from Beni Hassane region in north-western Morocco are both used in traditional medicine for the treatment of different ailments. The phenolic and flavonoid contents of roots, stems, leaves, and flowers of both species were determined by using the Folin Ciocalteu and the aluminum chloride methods, respectively. Antioxidant activity of their aqueous and organic compounds was performed applying the robust and widely used method ABTS radical scavenging assay. *C. mixtus* had a higher phenolic content (115.25 mg GAE g<sup>-1</sup> DW) than *M. chamomilla* (82.99 mg GAE g<sup>-1</sup> DW). However, flavonoid content of both plants was very close (33.53 and 32.45 mg QE g<sup>-1</sup> DW in *M. chamomilla* and *C. mixtus*, respectively). The total phenolic and flavonoid content was high in flowers of both plants. Generally, for all the studied organs of the chosen plants, hydrophilic antioxidant activity was significantly higher than the lipophilic antioxidant activity. The correlation results show that these bioactive components in both plants are the main factor influencing the antioxidant activity.

**Key words:** *Cladanthus mixtus*, *Matricaria chamomilla*, Asteraceae, phenolic content, flavonoid content, antioxidant activity.

## INTRODUCTION

Medicinal plants are frequently used as remedies in developing countries. For example, in Africa, more than 80% of the inhabitants depend on traditional medicines

for their medication (Emmanuel and Didier, 2012; Komoreng et al., 2017; Sevindik et al., 2017). Also, the African continent is characterized for having a wide

\*Corresponding author. E-mail : [amina.el@um.es](mailto:amina.el@um.es). Tel: +34 603 22 07 55.

variety of plant species, used mainly for medicine, among 300 000 plant species identified, more than 200 000 species live in tropical Africa and are used for their healthy properties. Morocco is characterized for having a good geographical position (a country with Mediterranean and Atlantic coasts) which has allowed it to have an interesting plant diversity (Scherrer et al., 2005; Ghanmi et al., 2011). Effectively, amongst more than 7 800 species listed in North Africa, Morocco presents the greatest biodiversity (Dobignard and Chatelain, 2010-2013), Herbal medicine is an important cultural tradition in Morocco (Benkhniqne et al., 2010) and plays a central role in the daily life of many rural and urban Moroccans (Jouad et al., 2001; Eddouks et al., 2002; El-Hilaly et al., 2003; Merzouki et al., 2003; Tahraoui et al., 2007; Jamila and Mostafa, 2014; Eddouks et al., 2017).

Bioactive compounds of plants are natural products that show some Health-related and interesting activities such as anti-hyperglycemia, anti-inflammatory, anti-microbial, antioxidant, anti-cancer and, antiviral activities, and more generic actions such as pain reliever, nerve relaxant, tranquilizer, spasmodic, etc. (Winklhofer-Roob et al., 2003; Wang et al., 2014; Baser and Buchbauer, 2015; Akhlaghi and Foshati, 2017; Altemimi et al., 2017; Desmarchelier and Borel, 2017; Pehlivan et al., 2018; Ashktorab et al., 2019; Bohn, 2019; Mohammed et al., 2019; Agati et al., 2020; Mu et al., 2021). Considering their great efficacy and low toxicity, natural products have been extensively studied and introduced as a chemopreventive therapy for many diseases including cancer (Al-Hrout et al., 2018; Amin et al., 2021)

*Gladanthus mixtus* (L.) Oberpr. and Vogt. (synonymous with *Ormenis mixta* subsp. *Mixta*, *Anthemis mixta* L., Moroccan chamomile, simple leaved chamomile), is a traditional herbal medicine, widespread in Morocco, Algeria, Northern and Eastern part of the Mediterranean basin. The infusion of Moroccan chamomile leaves and flowers is used in Moroccan traditional medicine for treatment of different ailments, also as an anxiolytic and to rebalance the central nervous system. The essential oil showed significant antimicrobial, anti-inflammatory, antinociceptive, antioxidant and antibacterial effects (Satrani et al., 2007; Zrira et al., 2007; Merghoub et al., 2009; Hajjaj et al., 2016).

*Matricaria chamomilla* L. (synonymous with *Matricaria recutita*, German chamomile, chamomile and little chamomile) is commonly used as a medicinal plant due to its anti-inflammatory, analgesic, anti-microbial, anti-allergic, anti-hyperglycemia, anticancer, and anti-spasmodic effects in pregnant women, and for its sedative properties. Generally, essential oil of chamomile plant is used in pharmaceutical, cosmetic, and food industries (Wu et al., 2012; Dadashpour et al., 2018; Abai et al., 2019; Suroowan et al., 2019; Asadi et al., 2020; Hoferl et al., 2020; Ahanni Arazi and Danesh, 2021; Heidarianpour et al., 2021).

Many scientific studies have shown that free radicals play a major role in the development of different diseases

and dysfunctions, such as cancer, heart disease, aging, cataracts and immune system deterioration (García-Conesa and Larrosa, 2020; Sevindik, 2021). Free radicals can be eliminated by antioxidants, which inhibit the rate of lipid and protein oxidation and protect cells from oxidative damage (Asimi et al., 2013; Mohammed et al., 2020b). Antioxidant drugs are used for the prevention and treatment of oxidative stress related disease such as diabetes, Alzheimer's disease, atherosclerosis, heart stroke and cancer (Devasagayam et al., 2004; Khalipha et al., 2012; Mohammed et al., 2020a). Antioxidant activity in plant extracts is a widely used parameter to characterize the possible beneficial properties of plants (Shahidi and Ambigaipalan, 2015). The most widely used methods for measuring antioxidant activity are those that involve the generation of free radical species, and the presence of antioxidant compounds, which determine the disappearance of these free radicals (Arnao et al., 1999; Pehlivan et al., 2021). For the analysis of *in vitro* and *in vivo* antioxidant activity numerous methods have been developed, but only a few fast and reliable methods exist for a large number of plant samples. An extensive study on antioxidant activity measurement methods can be consulted in Apak et al. (2018).

We applied the end-point method using 2,2'-azino-bis-3-ethylbenzthiazoline-6-sulfonic acid (ABTS) as chromogen for the estimation of the total antioxidant activity (TAA) developed by our group (Arnao et al., 2001a). This method allows the calculation of the contribution of lipophilic (in organic media) and hydrophilic (in buffered media) compounds to total antioxidant activity (TAA), referred as hydrophilic and lipophilic antioxidant activities. This is a robust method, widely used and applied to diverse biological samples (Cano and Arnao, 2018). Although this method is originated for plant foodstuffs, it has also been applied to characterize other diverse biological materials such as plant extracts (Arnao, 2000; Arnao et al., 2001a, b; El Omari et al., 2003) and animal samples (Espinosa et al., 2012; Rubio et al., 2016).

In this work, we present a comparative study of bioactive components of German and Moroccan chamomile plants, studying the lipophilic and hydrophilic antioxidant activities in the different plant organs (root, stem, leaf, and flower). Also, total phenolic and flavonoid contents were determined in these two chamomile species. To the best of our knowledge, this is the first comparative study on these two plants from north-west Morocco (Tangier-Tetouan-Al Hoceïma region).

## MATERIALS AND METHODS

### Plant material and preparation of extracts

*C. mixtus* and *M. chamomilla* plants were collected at full maturity from Beni Hassane (Tangier-Tetouan-Al Hoceïma region, north-west of Morocco) in May 2018 and transported to the laboratory. Table 1 shows the plant's geographical localization and GPS

**Table 1.** Geographical location and GPS coordinates of studied plant species.

Plant species	GPS coordinates	Locality
<i>Matricaria chamomilla</i>	N 35° 21' 18.865" W 5° 22' 11.677"	Beni Hassane (Tangier-Tetouan-Al Hoceïma region)
<i>Cladanthus mixtus</i>	N 35° 21' 20.865" W 5° 22' 12.677"	

coordinates. Plant identification was done by Prof. Lamarti in Tetouan (Morocco). After separating the organs (roots, stems, leaves, and flowers) in fresh plants, the material was dried in an oven until stabilization of dry weight at 50°C in order to preserve the integrity of its natural composition as much as possible. After, it was ground in a Microtron-MB550 (KINEMATICA AG, Germany), at 8,000 rpm. The powder obtained was composed of particles with a diameter of around 0.2 mm, and stored in the dark at room temperature.

A mixture of 0.1 g of homogeneous powder, 5 ml of 50 mM Na phosphate buffer (pH 7.4) and 8 ml of ethyl acetate was crushed in a Euroturax T20 (IKA, Germany) for 2 min and transferred to a decantation funnel. The solid residue (totally colorless) was discarded. The aqueous phase was collected and analyzed as soon as possible for hydrophilic antioxidant activity, and aqueous phenolic and aqueous flavonoid contents. The organic phase (ethyl acetate) was collected and stored at -20°C until analysis to measure lipophilic antioxidant activity, and organic phenolic and organic flavonoid contents. In all cases, three replicate extraction were performed.

#### Total phenolic content

Folin-Ciocalteu's reagent was applied to determine the total content of phenolic compounds in the samples as described by Singleton and Rossi (1965) with some modifications. Fifty microliters of sample (hydrophilic or lipophilic phase) was placed in a glass test tube, then 950 µl of distilled water, 50 µl of 1 M sodium carbonate and 50 µl of Folin Ciocalteu reagent were added. The mixture was left to stand in a water bath for 15 min at 30°C, and the absorbance was measured at 715 nm. Gallic acid was used as standard. The results were expressed as mg gallic acid equivalents per gram of dry weight (mg GAE g<sup>-1</sup> DW). Photometric measurements were recorded on a Perkin-Elmer Lambda-2S UV-VIS spectrophotometer (Loughborough, UK). Experiments were conducted in triplicates.

#### Total flavonoids content

The aluminum chloride colorimetric method, modified by Woisky and Salatino (1998), was used to determine total flavonoid content. Quercetin, used as standard, was dissolved in 80% ethanol and then diluted to 25, 50, 100 and 150 µg·mL<sup>-1</sup>. The diluted standard solutions (0.5 ml) were separately mixed with 1.5 ml of 95% ethanol, 0.1 ml of 10% aluminum chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml of distilled water. After incubation at room temperature for 30 min, the absorbance of the reaction mixture was measured at 415 nm. A similar step was applied to the plant samples for the flavonoid content analysis. The results were expressed as mg of quercetin equivalents per gram of dry weight (mg QE·g<sup>-1</sup> DW). Experiments were conducted in triplicates.

#### Hydrophilic and lipophilic antioxidant activities using ABTS radical scavenging assay

Hydrophilic and lipophilic antioxidant activities were measured using our ABTS/H<sub>2</sub>O<sub>2</sub>/HRP decoloration methods (Arnao et al., 1999). These methods are based on the capacity of a sample to scavenge the ABTS radical cation (ABTS<sup>•+</sup>) compared to a standard antioxidant (Trolox) in a dose-response curve. The ABTS<sup>•+</sup> blue color changes to pale yellow. This method allows to measure separate antioxidant activity due to hydrophilic and lipophilic compounds, being able to operate mathematically with both values, obtaining the sum as the total antioxidant activity (TAA=HAA+LAA). The hydrophilic antioxidant activity (HAA) is measured in buffered medium and the lipophilic antioxidant activity (LAA) is measured in organic medium.

For the determination of the lipophilic antioxidant activity (LAA), the reaction mixture contained 1 mM ABTS, 75 µM hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), and 6 µM horseradish peroxidase (HRP) type VI in acidified ethanol (pure ethanol with phosphoric acid, 0.7%, w/v), in a total volume of 1 ml, at 25°C. In this case, 40 µl of the organic phase obtained during the plant extraction was added to the reaction medium and the decrease in absorbance at 730 nm was measured after 6 min. The total time needed to carry out each assay was approximately 6 min, including ABTS radical generation by peroxidase, the addition of antioxidant, and acquisition of the final absorbance value. The absorbance decrease was measured from the difference between the A<sub>730</sub> values before and 6 min after sample addition. Antioxidant activity was calculated as moles of ABTS<sup>•+</sup> quenched by 1 mole of Trolox. LAA was expressed as Trolox equivalents per g of dry weight (mg TE·g<sup>-1</sup> DW). Experiments were conducted in triplicates.

For hydrophilic antioxidant activity (HAA), the reaction mixture contained 2 mM ABTS, 60 µM H<sub>2</sub>O<sub>2</sub>, and 0.2 µM HRP in 50 mM Na-phosphate buffer (pH 7.5) in a total volume of 1 ml. The reaction was monitored at 730 nm until a stable absorbance was obtained. The assay temperature was 25°C. Then, 10 µl of the aqueous sample was added to the reaction medium and the decrease in absorbance, which is proportional to the ABTS<sup>•+</sup> quenched, was determined after 6 min, as Trolox equivalents per g of dry weight (mg Trolox·g<sup>-1</sup> DW). Experiments were conducted in triplicates. The concentrations of ABTS, H<sub>2</sub>O<sub>2</sub>, and Trolox were determined by measuring their absorbance using  $\epsilon_{340}=36$  mM<sup>-1</sup> cm<sup>-1</sup> for ABTS,  $\epsilon_{240}=43.6$  M<sup>-1</sup> cm<sup>-1</sup> for H<sub>2</sub>O<sub>2</sub> and  $\epsilon_{403}=100$  mM<sup>-1</sup> cm<sup>-1</sup> for HRP.

#### Statistical analysis

For statistical analysis of the results, the SPSS program (Chicago, USA) was used applying the One-way ANOVA to evaluate the statistical differences among the group and the Tukey multiple range test to establish significant differences between the evaluated parameters.

**Table 2.** Aqueous and organic phenolic contents in different organs of *Matricaria chamomilla* and *Cladanthus mixtus*.

Organ	<i>Matricaria chamomilla</i>		<i>Cladanthus mixtus</i>	
	Aqueous phase	Organic phase	Aqueous phase	Organic phase
Root	4.41 <sup>+</sup> ± 0.12 <sup>d</sup>	4.16 ± 0.43 <sup>b</sup>	11.31 ± 0.32 <sup>c</sup>	6.48 ± 0.28 <sup>c</sup>
Stem	13.45 ± 0.52 <sup>b</sup>	8.13 ± 0.48 <sup>a</sup>	18.09 ± 1.84 <sup>b</sup>	6.86 ± 0.23 <sup>bc</sup>
Leaf	10.56 ± 0.64 <sup>c</sup>	9.21 ± 1.05 <sup>a</sup>	25.15 ± 0.10 <sup>a</sup>	8.31 ± 0.67 <sup>b</sup>
Flower	21.78 ± 0.49 <sup>a</sup>	11.29 ± 0.82 <sup>a</sup>	22.60 ± 0.42 <sup>a</sup>	16.45 ± 0.04 <sup>a</sup>

\*Values are the mean ± SD of three replicates expressed as mg gallic acid equivalents (GAE)·g<sup>-1</sup> DW. Different letters indicates significant differences between the organs for the same column at p < 0.05.

## RESULTS AND DISCUSSION

### Determination of phenolic content of *M. chamomilla* and *C. mixtus*

Phenolic compounds, such as flavonols, flavones, and phenolic acids, are synthesized by plants in the so called secondary metabolism. *M. chamomilla* is known to have high phenolic content (Gee and Johnson, 2001; Guimarães et al., 2013) that may contribute to its antioxidant activity (Lim and Quah, 2007; Wojdyło et al., 2007).

Table 2 shows the phenolic content of different organs of *M. chamomilla* and *C. mixtus*. Aqueous phenolic content was higher in all organs of both plants compared to organic phenolic content. The calculations were made using a calibration curve that showed a good linearity for gallic acid with a correlation coefficient R<sup>2</sup> of 0.9988. Results showed, for the same phase, significant differences (p < 0.05) between the organs.

*M. chamomilla* showed values of aqueous phenolic content between 4.41 and 21.78 mg GAE·g<sup>-1</sup> DW, with the flowers showing the highest content, followed by the stems, leaves, and roots. Organic phenolic content was between 4.16 and 11.29 mg GAE·g<sup>-1</sup> DW, with the flowers showing the highest level followed by the leaves, stems, and roots. Consequently, total phenolic content, as the sum of the aqueous and organic phenolic contents, was highest in flowers (40%), followed by the stems (26%), leaves (24%) and roots (10%) (Figure 1).

The total phenolic content of flower extract of *M. chamomilla* found in this work (as the sum of the aqueous and organic phenolic contents = 33.07 mg GAE·g<sup>-1</sup> DW) is slightly higher than that found in the flower extract from a similar study with Italian chamomilla (2,689.2 mg GAE·100 g<sup>-1</sup> DW) (Formisano et al., 2015), and much higher than in a study with Egyptian chamomille (3.7 mg GAE·g<sup>-1</sup> DW) (Roby et al., 2013), and than in a commercial chamomile product from United Arab Emirates (21.4 mg GAE·g<sup>-1</sup> DW) (Al-Dabbagh et al., 2019). Thus, chamomile flower from this study still has the highest value of total phenolic content, possibly due to that the applied bi-phasic extraction method is more exhaustive.

The extracts of *C. mixtus* had the highest total phenolic content compared to *M. chamomilla* (Table 2). The aqueous phenolic content ranged between 11.31 and 25.15 mg GAE·g<sup>-1</sup> DW, with the leaves showing the highest content, followed by the flowers, stems, and roots. The organic phenolic content ranged between 6.48 and 16.45 mg GAE·g<sup>-1</sup> DW, following the order: flower > leaves > stems > roots. Therefore, the highest total phenolic content was observed in flowers (34%), followed by leaves (29%), stems (22%), and roots (15%) (Figure 1). In a previous report, extracts from aerial parts of *C. mixtus* (collected from Bouznika, Morocco) showed total phenolic content of 38.2 mg GAE·g<sup>-1</sup> DW in the aqueous extracts, and of 19.5 mg GAE·g<sup>-1</sup> DW in methanolic extracts (Elouaddari et al., 2019), showing lower content than the one found in this study (65.84 and 31.62 mg GAE·g<sup>-1</sup> DW, respectively).

### Determination of flavonoid content of *M. chamomilla* and *C. mixtus*

Table 3 shows the flavonoid content of different organs of *M. chamomilla* and *C. mixtus*. To quantify flavonoid content in plant extracts, we used quercetin as standard, obtaining a linear calibration curve and a correlation coefficient R<sup>2</sup> of 0.9994. Results showed that aqueous flavonoid content was significantly higher in all organs for both plants compared to organic flavonoid content. *M. chamomilla* showed values of aqueous flavonoid content between 2.11 and 7.97 mg QE·g<sup>-1</sup> DW, and of organic flavonoid content between 0.50 and 5.65 mg QE·g<sup>-1</sup> DW.

In *C. mixtus*, aqueous flavonoid content was between 3.70 and 10.20 mg QE·g<sup>-1</sup> DW, and the organic flavonoid content between 0.99 and 3.74 mg QE·g<sup>-1</sup> DW. Both plants showed the highest flavonoid content in the flowers, followed by leaves, stems, and roots. *M. chamomilla* and *C. mixtus* showed close proportions of total flavonoid content in the flowers (41 and 43%, respectively) and in stems (15 and 17%, respectively) (Figure 2).

In Italian chamomile flowers, a total flavonoid content of 710.7 mg QE·100 g<sup>-1</sup> DW was observed (Formisano et al., 2015), which is lower than in our flowers extract of *M.*




Figure 1. Percentage of phenolic compounds in different organs of *Matricaria chamomilla* and *Cladanthus mixtus*.

Table 3. Aqueous and organic flavonoid content in different organs of *Matricaria chamomilla* and *Cladanthus mixtus*.

Organ	<i>Matricaria chamomilla</i>		<i>Cladanthus mixtus</i>	
	Aqueous phase	Organic phase	Aqueous phase	Organic phase
Root	*2.11 ± 0.23 <sup>c</sup>	0.50 ± 0.04 <sup>b</sup>	3.70 ± 0.11 <sup>c</sup>	0.99 ± 0.02 <sup>c</sup>
Stem	3.98 ± 0.23 <sup>b</sup>	1.12 ± 0.41 <sup>b</sup>	4.34 ± 0.35 <sup>c</sup>	1.36 ± 0.08 <sup>c</sup>
Leaf	6.78 ± 0.24 <sup>a</sup>	5.42 ± 0.16 <sup>a</sup>	5.34 ± 0.09 <sup>b</sup>	2.78 ± 0.17 <sup>b</sup>
Flower	7.97 ± 0.36 <sup>a</sup>	5.65 ± 0.05 <sup>a</sup>	10.20 ± 0.07 <sup>a</sup>	3.74 ± 0.10 <sup>a</sup>

\*Values are the mean ± SD of three replicates expressed as mg quercetin equivalents (QE)·g<sup>-1</sup> DW. Different letters indicates significant differences at p < 0.05 differences between the organs for the same column at p < 0.05.


*chamomilla* (13.62 mg QE·g<sup>-1</sup> DW). In a previous study of aerial parts of Moroccan *C. mixtus*, the flavonoid content was 3.2 and 2.7 mg QE·g<sup>-1</sup> DW in methanolic and aqueous extracts, respectively (Elouaddari et al., 2019). In our case, *C. mixtus* contained a flavonoid content of 19.88 and 7.88 mg QE·g<sup>-1</sup> DW, in aqueous and organic extracts, respectively. Globally, a total flavonoid content in aerial parts of 27.76 mg QE·g<sup>-1</sup> DW was determined in our estimations, a much greater amount than in the mentioned study of Elouaddari et al. (2019).

**Determination of the antioxidant activity of *M. chamomilla* and *C. mixtus***

Preformed ABTS radical (ABTS<sup>•+</sup>) is considered to be an excellent chromogen for use in measurement of the

antioxidative properties of pure products and biological samples (Arnao et al., 1999, 2001b). Figure 3 shows hydrophilic antioxidant activity (HAA) and lipophilic antioxidant activity (LAA) of different organs of *M. chamomilla* and *C. mixtus*. Results of *M. chamomilla* showed that HAA ranged from 9.28 to 18.02 mg TE·g<sup>-1</sup> DW, while LAA ranged from 1.47 to 3.51 mg TE·g<sup>-1</sup> DW. Therefore, TAA (HAA+LAA) was formed mainly of HAA (87%) and the rest (12%) of LAA (Figure 4).

Differences of antioxidant activity were noted between the studied organs of *M. chamomilla*. Concerning HAA, the order was as followed: roots (18.02 mg TE·g<sup>-1</sup> DW) ≥ flowers (17.57 mg TE·g<sup>-1</sup> DW) > stems (12.27 mg TE·g<sup>-1</sup> DW) > leaves (9.28 mg TE·g<sup>-1</sup> DW). For LAA the order was as followed: flowers (3.51 mg TE·g<sup>-1</sup> DW) > stems (1.49 mg TE·g<sup>-1</sup> DW) ≅ roots (1.48 mg TE·g<sup>-1</sup> DW) ≅ leaves (1.47 mg TE·g<sup>-1</sup> DW). Significant difference was



**Figure 2.** Percentage of flavonoid compounds in different organs of *Matricaria chamomilla* and *Cladanthus mixtus*.

reported between LAA of flowers and other organs ( $p < 0.05$ ). Consequently, the flowers of *M. chamomilla* showed the highest TAA while leaves had the lowest TAA. *C. mixtus* also showed greater HAA than LAA, ranging from 12.89 to 23.40 mg TE·g<sup>-1</sup> DW for HAA, and from 1.34 to 3.99 mg TE·g<sup>-1</sup> DW for LAA. Thus, TAA was formed of 89% HAA and 11% LAA (Figure 4). Significant difference was noted between HAA of the studied organs ( $p < 0.05$ ), with stems showing the highest HAA (23.40 mg TE·g<sup>-1</sup> DW), followed by flowers (20.32 mg TE·g<sup>-1</sup> DW), roots (14.11 mg TE·g<sup>-1</sup> DW) and leaves (12.89 mg TE·g<sup>-1</sup> DW). As in *M. chamomilla*, the flowers of *C. mixtus* had the highest LAA. Significant difference was noted between LAA of flowers and other organs ( $p < 0.05$ ), with the order: flowers > roots > leaves > stems.


#### Relationship of the phenolic and flavonoid contents with the antioxidant activity of *M. chamomilla* and *C. mixtus*

Some natural products may act to regulate cell growth, differentiation, apoptosis, and oxidative stress (Hamza et al., 2018; Kamal et al., 2018, Mohammed et al., 2020b; Amin et al., 2021; Sevindik, 2021). Phenolic molecules are known for their antioxidant activity. For example, Gallic acid is reported to have antioxidant, antimicrobial, antitumor and anti-inflammatory properties activities (Lima et al., 2016). Since phenolics and flavonoids may contribute to the antioxidant activity of extracts, we have studied the correlation between these compound contents and the antioxidant activity (Tables 4 and 5).

The correlation coefficient is a statistical measure of linear association between two variables, ranging

between -1 and 1. A correlation value of 1 or -1 indicates that two variables are perfectly related in a positive or negative linear sense, and a value of 0 indicates that there is no linear relationship between the two variables (Asuero et al., 2006). Regarding phenolic content and antioxidant activities (Table 4), in stems, leaves, and flowers of *M. chamomilla*, an excellent significant positive correlation was obtained between HAA and aqueous phenolics ( $r^2 = 0.98$ ,  $p < 0.05$ ), and an acceptable significant positive correlation was obtained between TAA and total phenolic content ( $r^2=0.87$ ,  $p < 0.05$ ). In roots, stems, and flowers, a significant positive correlation was observed between LAA and organic phenolics ( $r^2=0.70$ ,  $p < 0.05$ ). In the case of *C. mixtus*, a good significant positive correlation was observed between aqueous phenolics and HAA ( $r^2= 0.85$ ,  $p < 0.05$ ) in stems, leaves, and flowers (Table 4). Between organic phenolics and LAA, an extremely significant positive correlation was found in all organs ( $r^2= 0.92$ ,  $p < 0.05$ ). Lower significant positive correlation ( $r^2=0.53$ ,  $p < 0.05$ ) was observed in roots, stems, and flowers of *C. mixtus*. Therefore, it can be concluded that stems and flowers always showed a good correlation between their phenolic compounds and antioxidant activities.

Regarding flavonoid content and antioxidant activities (Table 5), a significant positive correlation was also reported in roots, stems, and leaves of *M. chamomilla* between aqueous flavonoid content and HAA ( $r^2=0.91$ ,  $p < 0.05$ ), and between total flavonoids content and TAA ( $r^2=0.81$ ,  $p < 0.05$ ). Also, a significant positive correlation was observed between organic flavonoid content and LAA ( $r^2= 0.98$ ,  $p < 0.05$ ) in roots, stems, and flowers. In *C. mixtus*, a great significant positive correlation was found between HAA and aqueous flavonoid content ( $r^2 =$




**Figure 3.** Lipophilic and hydrophilic antioxidant activities of different organs extracts of *Matricaria chamomilla* and *Cladanthus mixtus*. Different letters indicate significant differences between the organs for one plant at  $p < 0.05$ . Values are means  $\pm$  S.D. for 3 replicates.

0.91;  $p < 0.05$ ) in roots, stems, and flowers. Also, a good significant positive correlation was observed between organic flavonoid content and LAA ( $r^2 = 0.84$ ,  $p < 0.05$ ), and, in a lesser extent, between total flavonoid content and TAA ( $r^2 = 0.75$ ,  $p < 0.05$ ) of roots, leaves, and flowers.

This analysis confirms that phenolic and flavonoid contents are the main contributors to the total antioxidant

activity in our study. Nonetheless, it is important to take into consideration that different phenolic compounds can contribute in different ways to TAA, resulting in different antioxidant activities. Accordingly, it is very important to obtain real TAA measurements by methods or tests that quantify antioxidant activities of aqueous and organic antioxidant compounds in their respective media (Cano and Arnao, 2018).



**Figure 4.** Percentage of antioxidant activity in different organs of *Matricaria chamomilla* and *Cladanthus mixtus*.

**Table 4.** Relationship between phenolic contents and antioxidant activities (HAA, LAA, and TAA) of different organs of *Matricaria chamomilla* and *Cladanthus mixtus*.

<i>Matricaria chamomilla</i>	Relationship between parameter		
	Organ	Equation	R <sup>2*</sup>
Aqueous phenolics vs HAA	Stem, leaf, flower	$y = 0.7165x + 2.1017$	0.98
Organic phenolics vs LAA	Root, stem, flower	$y = 0.0543x + 1.1914$	0.70
Total phenolics vs TAA	Stem, leaf, flower	$y = 0.6345x - 1.4994$	0.87
<b><i>Cladanthus mixtus</i></b>			
Aqueous phenolics vs HAA	Stem, leaf, flower	$y = -1.3937x + 49.462$	0.85
Organic phenolics vs LAA	Root, stem, leaf, flower	$y = 0.2528x - 0.2369$	0.92
Total phenolics vs TAA	Root, stem, flower	$y = 0.3338x + 12.572$	0.53

\* indicates significant correlation at  $p < 0.05$ .

**Table 5.** Relationship between flavonoid contents and antioxidant activities (HAA, LAA and TAA) of different organs of *Matricaria chamomilla* and *Cladanthus mixtus*.

<i>Matricaria chamomilla</i>	Relationship between parametres		
	Organ	Equation	R <sup>2*</sup>
Aqueous flavonoids vs HAA	Root, stem, leaf	$y = -1.8111x + 20.972$	0.91
Organic flavonoids vs LAA	Root, stem, flower	$y = 0.4135x + 1.1562$	0.98
Total flavonoids vs TAA	Root, stem, leaf	$Y = -0.8049x + 20.021$	0.81
<b><i>Cladanthus mixtus</i></b>			
Aqueous flavonoids vs HAA	Root, stem, flower	$y = 0.9002x + 0.5625$	0.91
Organic flavonoids vs LAA	Root, leaf, flower	$y = 1.0854x + 8.8089$	0.84
Total flavonoids vs TAA	Root, leaf, flower	$y = 0.9875x + 9.4099$	0.75

\*Indicates significant correlation at  $p < 0.05$ .

## Conclusion

This study investigated the bioactive components and the

total antioxidant activity levels measured in hydrophilic and lipophilic solutions of different organs of *C. mixtus* and *M. chamomilla*. The results indicate that hydrophilic



antioxidant activity, using ABTS radical scavenging assay, was significantly higher than the lipophilic antioxidant activity, in both plants. In addition, total phenolic and flavonoid contents of both plants are also highly relevant. Therefore, it can be concluded that *C. mixtus* and *M. chamomilla* are rich in phenolic and flavonoid contents and have a strong antioxidant activity, mainly in flowers. The correlation analysis results show that the phenolic and flavonoid contents in both plants are the main factors influencing the antioxidant activity. This study suggests that both plants present interesting health-related bioactivities, but more specific bioassays should be made to determine the therapeutic and food applications of interest.

## CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

## REFERENCES

- Abdi P, Mahdavi Ourtakand M, Honarmand Jahromy S (2019). The Effect of *Matricaria chamomilla* Alcoholic Extract on Phenotype Detection of Efflux Pumps of Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated from Skin Lesions. Iranian Journal of Medical Microbiology 13(3):220-231.
- Agati G, Brunetti C, Fini A, Gori A, Guidi L, Landi M, Sebastiani F, Tattini M (2020). Are flavonoids effective antioxidants in plants? Twenty years of our investigation. Antioxidants 9(11):1098.
- Ahani Azari A, Danesh A (2021). Antibacterial Effect of *Matricaria chamomilla* Alcoholic Extract against Drug-Resistant Isolates of *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Infection Epidemiology and Microbiology 7(1):29-35.
- Akhlaghi M, Foshati S (2017). Bioavailability and metabolism of flavonoids: A Review. International Journal of Nutrition Sciences 2(4):180-184.
- Al-Dabbagh B, Elhady IA, Elhaw M, Murali C, Al Mansoori A, Awad B, Amin A (2019). Antioxidant and anticancer activities of chamomile (*Matricaria recutita* L.). BMC Research Notes 12(1):1-8.
- Al-Hrout A, Chaiboonchoe A, Khraiweh B, Murali C, Baig B, El-Awady R, Tarazi H, Alzahrani A, Nelson DR, Greish YE, Ramadan W, Salehi-Ashtiani K, Amin A (2018). Safranal induces DNA double-strand breakage and ER-stress-mediated cell death in hepatocellular carcinoma cells. Scientific Reports 8(1):1-15.
- Altemimi A, Lakhssassi N, Baharlouei A, Watson GD, Lightfoot AD (2017). Phytochemicals: Extraction, isolation, and identification of bioactive compounds from plant extracts. Plants 6(4):42.
- Amin A, Farrukh A, Murali C, Soleimani A, Praz F, Graziani G, Brim H, Ashktorab H (2021). Saffron and Its Major Ingredients' Effect on Colon Cancer Cells with Mismatch Repair Deficiency and Microsatellite Instability. Molecules 26(13):3855.
- Apak R, Çapanoğlu E, Shahidi F (2018). Measurement of antioxidant activity and capacity: Recent Trends and Applications. Functional Foods Science and Technology Series. John Wiley and Sons Ltd. Published. Oxford, UK pp. 117-139.
- Arnao MB (2000). Some methodological problems in the determination of antioxidant activity using chromogen radicals: a practical case. Trends in Food Science and Technology 11(11):419-421.
- Arnao MB, Cano A, Acosta M (1999). Methods to measure the antioxidant activity in plant material: a comparative discussion. Free Radical Research 31(1):89-96.
- Arnao MB, Cano A, Acosta M (2001b). The hydrophilic and lipophilic contribution to total antioxidant activity. Food Chemistry 73(2):239-244.
- Arnao MB, Cano A, Alcolea JF, Acosta M (2001a). Estimation of free radical-quenching activity of leaf pigment extracts. Phytochemical Analysis 12(2):138-143.
- Asadi Z, Ghazanfari T, Hatami H (2020). Anti-inflammatory Effects of *Matricaria chamomilla* Extracts on BALB/c Mice Macrophages and Lymphocytes. Iranian Journal of Allergy, Asthma and Immunology. Available at: <https://publish.kne-publishing.com/index.php/IJAAI/article/view/2862/2108>
- Ashktorab H, Soleimani A, Singh G, Amin A, Tabtabaei S, Latella G, Stein U, Akhondzadeh S, Solanki N, Gondré-Lewis MC, Habtezion A, Brim H (2019). Saffron: the golden spice with therapeutic properties on digestive diseases. Nutrients 11(5):943.
- Asimi OA, Sahu NP, Pal AK (2013). Antioxidant activity and antimicrobial property of some Indian spices. International Journal Scientific Research Publications 3(3):1-8.
- Asuero AG, Sayago A, Gonzalez AG (2006). The correlation coefficient: An overview. Critical Reviews in Analytical Chemistry 3(1):41-59.
- Baser CHK, Buchbauer G (2015). Handbook of Essential Oils: Science, Technology, and Applications. CRC Press, BocaRaton, FL, USA. 2nd Edition.
- Benkhngue O, Zidane L, Fadli M, Elyacoubi H, Rochdi A, Douira A (2010). Etude ethnobotanique des plantes médicinales dans la région de Mechraâ Bel Ksiri (Région du Gharb du Maroc). Acta botánica barcinonensia pp.191-216.
- Bohn T (2019). Carotenoids and markers of oxidative stress in human observational studies and intervention trials: Implications for chronic diseases. Antioxidants 8(6):179.
- Cano A, Arnao MB (2018). ABTS/TEAC (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)/Trolox-Equivalent Antioxidant Capacity) radical scavenging mixed-mode assay. Measurement of antioxidant activity and capacity: Recent Trends and Applications. Functional Foods Science and Technology Series, chapter 7, pp. 117-139.
- Dadashpour M, Firouzi-Amandi A, Pourhassan-Moghaddam M, Maleki MJ, Soozangar N, Jeddi F, Nouric M, Zarghamie N, Pilehvar-Soltanahmadi Y (2018). Biomimetic synthesis of silver nanoparticles using *Matricaria chamomilla* extract and their potential anticancer activity against human lung cancer cells. Materials Science and Engineering C 92:902-912.
- Desmarchelier C, Borel P (2017). Overview of carotenoid bioavailability determinants: From dietary factors to host genetic variations. Trends in Food Science and Technology 69:270-280.
- Devasagayam TPA, Tilak JC, Boloor KK, Sane KS, Ghaskadbi SS, Lele RD (2004). Free radicals and antioxidants in human health: current status and future prospects. Journal of the Association of Physicians of India 52:794-804.
- Dobignard A, Chatelain C (2010-2013). Index synonymique de la flore d'Afrique du Nord. Conservatoire et Jardin botaniques.
- Eddouks M, Ajebli M, Hebi M (2017). Ethnopharmacological survey of medicinal plants used in Daraa-Tafilalet region (Province of Errachidia), Morocco. Journal of Ethnopharmacology 198:516-530.
- Eddouks M, Maghrani M, Lemhadri A, Ouahidi ML, Jouad H (2002). Ethnopharmacological survey of medicinal plants used for the treatment of diabetes mellitus, hypertension and cardiac diseases in the south-east region of Morocco (Tafilalet). Journal of Ethnopharmacology 82(2-3):97-103.
- El Omari B, Fleck I, Aranda X, Abadia A, Cano A, Arnao MB (2003). Total antioxidant activity in *Quercus ilex* sprouts after fire. Plant Physiology and Biochemistry 41(1):41-47.
- El-Hilaly J, Hmammouchi M, Lyoussi B (2003). Ethnobotanical studies and economic evaluation of medicinal plants in Taounate province (Northern Morocco). Journal of Ethnopharmacology 86(2-3):149-158.
- Elouaddari A, Elamrani A, Moutia M, Oubrim N, Habti N, JamalEddine J (2019). Chemical composition and evaluation of antioxidant, antimicrobial and cytotoxic activities of Moroccan *Cladanthus mixtus* essential oil and extracts. Journal of Essential Oil Bearing Plants 22(6):1450-1466.
- Emmanuel MM, Didier DS (2012). Traditional knowledge on medicinal plants use by ethnic communities in Douala, Cameroon. European Journal of Medicinal Plants 2:159-176.
- Espinosa C, López-Jiménez JÁ, Cabrera L, Larqué E, Almajano MP, Arnao MB, Zamora S, Pérez-Llamas F (2012). Protective effect of white tea extract against acute oxidative injury caused by adriamycin

- in different tissues. *Food Chemistry* 134(4):1780-1785.
- Formisano C, Delfino S, Oliviero F, Tenore GC, Rigano D, Senatore F (2015). Correlation among environmental factors, chemical composition and antioxidative properties of essential oil and extracts of chamomile (*Matricaria chamomilla* L.) collected in Molise (South-central Italy). *Industrial Crops and Products* 63:256-263.
- García-Conesa MT, Larrosa M (2020). Polyphenol-Rich Foods for Human Health and Disease. *Nutrients* 12:400.
- Gee JM, Johnson IT (2001). Polyphenolic compounds: interactions with the gut and implications for human health. *Current Medicinal Chemistry* 8(11):1245-1255.
- Ghanmi M, Satrani B, Aberchane M, Ismaili MR, Aafi A, El Abid A (2011). Plantes Aromatiques et Médicinales du Maroc, Les milles et une vertu, Centre de Recherche Forestière, Rabat, Maroc. pp. 130.
- Guimarães R, Barros L, Dueñas M, Calheta RC, Carvalho Maria A, Santos-Buelga C, João M, Ferreira ICFR (2013). Nutrients, phytochemicals and bioactivity of wild Roman chamomile: A comparison between the herb and its preparations. *Food Chemistry* 136(2):718-725.
- Hajjaj G, Bahlouli A, Tajani M, Cherrah Y, Zellou A (2016). *In vivo* antinociceptive and anti-inflammatory properties of *Ormenis mixta* L. from Morocco. *International Journal of Pharmacy and Pharmaceutical Sciences* 8:49-52.
- Hamza AA, Heeba GH, Elwy HM, Murali C, El-Awady R, Amin A (2018). Molecular characterization of the grape seeds extract's effect against chemically induced liver cancer: *In vivo* and *in vitro* analyses. *Scientific Reports* 8(1):1-16.
- Heidarianpour A, Mohammadi F, Keshvari M, Mirazi N (2021). Ameliorative effects of endurance training and *Matricaria chamomilla* flowers hydroethanolic extract on cognitive deficit in type 2 diabetes rats. *Biomedicine and Pharmacotherapy* 135:111230.
- Höferl M, Wanner J, Tabanca N, Ali A, Gochev V, Schmidt E, Kaul VK, Singh V, Jirovetz L (2020). Biological activity of *Matricaria chamomilla* essential oils of various chemotypes. *Planta Medica International Open* 7(03):e114-e121.
- Jamila F, Mostafa E (2014). Ethnobotanical survey of medicinal plants used by people in Oriental Morocco to manage various ailments. *Journal of Ethnopharmacology* 154(1):76-87.
- Joad H, Haloui M, Rhiouani H, El Hilaly J, Eddouks M (2001). Ethnobotanical survey of medicinal plants used for the treatment of diabetes, cardiac and renal diseases in the North centre region of Morocco (Fez-Boulemane). *Journal of Ethnopharmacology* 77(2-3):175-182.
- Kamal H, Jafar S, Mudgil P, Murali C, Amin A, Maqsood S (2018). Inhibitory properties of camel whey protein hydrolysates toward liver cancer cells, dipeptidyl peptidase-IV, and inflammation. *Journal of Dairy Science* 101(10):8711-8720.
- Khalifa ABR., Ahmed F, Rahman MM (2012). Antioxidant and antidiarrhoeal potentiality of *Diospyros blancoi*. *International Journal of Pharmacology* 8(5):403-409.
- Komoreng L, Thekiso O, Lehasa S, Tiwani T, Mzizi N, Mokoena N, Khambule N, Ndebele S, Mdletshe N (2017). An ethnobotanical survey of traditional medicinal plants used against lymphatic filariasis in South Africa. *South African Journal of Botany* 111:12-16.
- Lim YY, Quah EP (2007). Antioxidant properties of different cultivars of *Portulaca oleracea*. *Food Chemistry* 103(3):734-740.
- Lima VN, Oliveira-Tintino CD, Santos ES, Morais LP, Tintino SR, Freitas TR, Geraldo YS, Pereira RL, Cruz RP, Menezes IR, Coutinho HD (2016). Antimicrobial and enhancement of the antibiotic activity by phenolic compounds: Gallic acid, caffeic acid and pyrogallol. *Microbial Pathogenesis* 99:56-61.
- Merghoub N, Benbacer L, Amzazi S, Morjani H, El-Mzibri M (2009). Cytotoxic effect of some Moroccan medicinal plant extracts on human cervical cell lines. *Journal of Medicinal Plants Research* 3(12):1045-1050.
- Merzouki A, Ed-Derfoufi F, Molero Mesa J (2003). Contribution to the knowledge of rifian traditional medicine III: phytotherapy of diabetes in Chefchaouen province (North of Morocco). *Ars Pharmaceutica* 44:59-67.
- Mohammed FS, Günel S, Şabik AE, Akgül H, Sevindik M (2020a). Antioxidant and Antimicrobial activity of *Scorzonera papposa* collected from Iraq and Turkey. *Kahramanmaraş Sütçü İmam Üniversitesi Tarım ve Doğa Dergisi* 23(5):1114-1118.
- Mohammed FS, Pehlivan M, Sevindik M (2019). Antioxidant, antibacterial and antifungal activities of different extracts of *Silybum marianum* collected from Duhok (Iraq). *International Journal of Secondary Metabolite* 6(4):317-322.
- Mohammed FS, Şabik AE, Doğan M, Selamoğlu Z, Sevindik M (2020b). Antioxidant potential of *Hypericum spectabile* JAUB. ET SPACH. *Bulletin of Biotechnology* 1(2):43-45.
- Mu C, Sheng Y, Wang Q, Amin A, Li X, Xie Y (2021). Potential compound from herbal food of *Rhizoma Polygonati* for treatment of COVID-19 analyzed by network pharmacology: Viral and cancer signaling mechanisms. *Journal of Functional Foods* 77:10414.
- Pehlivan M, Mohammed FS, Şabik AE, Kına E, Dogan M, Yumrutaş Ö, Sevindik M (2021). Some Biological activities of ethanol extract of *Marrubium globosum*. *Turkish Journal of Agriculture-Food Science and Technology* 9(6):1129-1132.
- Pehlivan M, Mohammed FS, Sevindik M, Akgul H (2018). Antioxidant and oxidant potential of *Rosa canina*. *Eurasian Journal of Forest Science* 6(4):22-25.
- Roby MHH, Sarhan MA, Selim KAH, Khalel KI (2013). Antioxidant and antimicrobial activities of essential oil and extracts of fennel (*Foeniculum vulgare* L) and chamomile (*Matricaria chamomilla* L). *Industrial Crops and Products* 44:437-445.
- Rubio CP, Hernández-Ruiz J, Martínez-Subiela S, Tvarijonavičute A, Arnao MB, Ceron JJ (2016). Validation of three automated assays for total antioxidant capacity determination in canine serum samples. *Journal of Veterinary Diagnostic Investigation* 28(6):693-698.
- Satrani B, Ghanmi M, Farah A, Aafi A, Fougrach H, Bourkhis B, Bousta D, Talbi M (2007). Composition chimique et activité antimicrobienne de l'huile essentielle de *Cladanthus mixtus*. *Bulletin de la Société de Pharmacie de Bordeaux* 146:85-96.
- Scherrer AM, Motti R, Weckerle CS (2005). Traditional plant use in the areas of Monte Vesole and Ascea, Cilento National Park (Campania, Southern Italy). *Journal of Ethnopharmacology* 97(1):129-143.
- Sevindik M (2021). Anticancer, antimicrobial, antioxidant and DNA protective potential of mushroom *Leucopaxillus gentianeus* (Qué.) Kotl. *Indian Journal of Experimental Biology* (IJE) 59(05):310-315.
- Sevindik M, Akgul H, Pehlivan M, Selamoğlu Z (2017). Determination of the therapeutic potential of *Mentha longifolia* ssp. *longifolia*. *Fresenius Environmental Bulletin* 26(7):4757-4763.
- Shahidi F, Ambigaipalan P (2015). Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects. A review. *Journal of Functional Foods* 18:820-897.
- Singleton VL, Rossi JA (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American journal of Enology and Viticulture* 16(3):144-158.
- Suroowan S, Pynee KB, Mahomoodally MF (2019). A comprehensive review of ethnopharmacologically important medicinal plant species from Mauritius. *South African Journal of Botany* 122:189-213.
- Tahraoui A, El-Hilaly J, Israïli ZH, Lyoussi B (2007). Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in South-Eastern Morocco (Errachidia province). *Journal of Ethnopharmacology* 110(11):105-117.
- Wang Y, Chung SJ, McCullough ML, Song WO, Fernandez ML, Koo SI, Chun OK (2014). Dietary carotenoids are associated with cardiovascular disease risk biomarkers mediated by serum carotenoid concentrations. *The Journal of Nutrition* 144(7):1067-1074.
- Winkhofer-Roob BM, Rock E, Ribalta J, Shmerling DH, Roob JM (2003). Effects of vitamin E and carotenoid status on oxidative stress in health and disease. evidence obtained from human intervention studies. *Molecular Aspects of Medicine* 24(6):391-402.
- Woisky R, Salatino A (1998). Analysis of propolis: Some parameters and procedures for chemical quality control. *Journal of Apicultural Research* 37(2):99-105.
- Wojdyło A, Oszmiański J, Czemerys R (2007). Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chemistry* 105(3):940-949.
- Wu YN, Xu Y, Yao L (2012). Anti-inflammatory and anti-allergic effects of German chamomile (*Matricaria chamomilla* L.). *Journal of Essential Oil Bearing Plants* 15(1):75-83.

Zrira S, Menut C, Bessiere J, Benjlalii B (2007). Chemical composition of the essential oils of Moroccan *Ormenis mixta* (L.) Dumort. ssp. multicaulis. *Journal of Essential Oil Bearing Plants* 10(5):378-385.