

Full Length Research Paper

Effects of inorganic nitrogen source and $\text{NH}_4^+ : \text{NO}_3^-$ ratio on proliferation of dog rose (*Rosa canina*)

Mahdi Shirdel², Alireza Motallebi-Azar^{1*}, Sirous Masiha¹, Najmedin Mortazavi²,
Mansour Matloobi¹ and Yavar Sharafi³

¹Department of Horticultural Sciences, Faculty of Agriculture, Tabriz University, Tabriz, Iran.

²Department of Horticultural Sciences, Faculty of Agriculture, Zanjan University, Zanjan, Iran.

³Department of Horticultural Science, Islamic Azad University, Maragheh Branch, Maragheh, Iran.

Accepted 9 June, 2011

Dog rose (*Rosa canina*) is one of the most important ornamental and medicinal plants which are used as rootstock for ornamental roses as *Rosa hybrida* and *Rosa floribunda*. The effects of different concentrations of NH_4NO_3 (10-15 and 20 mM), KNO_3 (15-20 and 25 mM) and $\text{NH}_4^+:\text{NO}_3^-$ ratios (10:25, 10:30, 10:35, 15:30, 15:35, 15:40, 20:35 and 20:45) was investigated on proliferation stage of dog rose cultivated on MS media with total nitrogen levels (35; 40, 45, 50, 55, 60 and 65 mM). The relative amounts of NH_4^+ and NO_3^- ions influenced the tested parameters. The highest shoot length and most node numbers were obtained with 20:45 $\text{NH}_4^+:\text{NO}_3^-$ ratios. Chlorosis leaf number was less with 15:35 and 20:45 $\text{NH}_4^+:\text{NO}_3^-$ ratios. When NO_3^- was higher than NH_4^+ , shoot length and node numbers were produced more and chlorosis leaf number was decreased. This study also demonstrates that with increasing nitrogen levels, the growth of shoots and node numbers were more but leaf chlorosis was decreased, therefore proliferation was better. The percentage of axillary shoot was affected by NH_4NO_3 concentrations and it was highest in 20 mM concentration, but with increasing KNO_3 , the leaf necrotic and feeble callusing percentage at the cut ends of explants were more and lower in row.

Key words: Dog rose, *in vitro* culture, MS nitrogen source, $\text{NH}_4^+ : \text{NO}_3^-$ ratio, proliferation.

INTRODUCTION

Rosa canina is a medicinal plant that its fruits have many important medicinal properties but unknown for many people's especially in Iran. However, this crop can be produced commercially and its orchards can be established such as other fruit trees (Sharafi, 2010; Iuncna, 2005). Fruits of the rose species are rich in minerals, vitamins (A, B1, B2, B3, C and K), sugars, phenolic compounds, carotenoids, tokepherol, bioflavonoid, tannins, organic acids, amino acids, volatile oils, vanillin and other photochemical such as antioxidant and antimicrobials. Medicinal properties and benefits of Rose are: Nutrient, mild laxative, mild diuretic, mild astringent, carminative, ophthalmic, tonic and verminfuge (Sharafi, 2010). Active ingredients of *R. canina* can protect cold, flu, infectious diseases, vitamin C deficiency and fever (Iuncna, 2005). Commercial propagation of

roses is usually done by cutting, although they can also be propagated by budding and grafting, which is difficult, undesirable and tedious process (Horn, 1992). Micropropagation of *Rosa* species is done through the meristem tip culture, shoot tip culture and axillary buds (Khosh-Khui and Sink, 1982; Skirvin et al., 1990; Rout et al., 1999). In recent years, several reports have been published on *in vitro* proliferation of roses, and have shown that tissue culture provides an alternative method for rapid multiplication (Horn et al., 1988). This technique is a tool for propagation of many plant species and is important for mass propagation (George, 1993). Although most rose micropropagation studies have used a solid agar-gelled medium; Chu et al. (1993) and Nikbakht et al. (2005) have reported that shoot proliferation of roses is higher on liquid medium rather than solidified medium because; nutrient absorption may be facilitated in this kind of medium (George, 1993). Most plant tissue culture media, which are characterized by high levels of nitrogen in the forms of ammonium nitrate and potassium nitrate, are still based on the nutrient component of Murashige

*Corresponding author. E-mail: yavarsharafi@iau-maragheh.ac.ir. Tel: +989144200882. Fax: 984213254506.

Table 1. Types of different $\text{NH}_4^+:\text{NO}_3^-$ ratios and total nitrogen.

$\text{NH}_4^+:\text{NO}_3^-$ ratio (mM)	Treatments								
	10:25	10:30	10:35	15:30	15:35	15:40	20:35	20:40	20:45
Total nitrogen (mM)	35	40	45	45	50	55	55	60*	65

*Total nitrogen and $\text{NH}_4^+:\text{NO}_3^-$ ratio are according to standard MS medium.

Table 2. Analysis of variance table.

Sources	Df	Mean square					
		Shoot length (cm)	Node number	Axillary shoot (%)	Chlorosis leaf number	Necrotic leaf number	Feeble callusing (%)
NH ₄ NO ₃	2	100**	57.75**	0.88*	1.63 ^{ns}	1.3 ^{ns}	18 ^{ns}
KNO ₃	2	15.6 ^{ns}	12.20 ^{ns}	0.14 ^{ns}	4.13*	9.2**	72**
KNO ₃ ×NH ₄ NO ₃ ($\text{NH}_4^+:\text{NO}_3^-$ ratio)	4	53**	23.52*	0.37 ^{ns}	3.51*	0.95 ^{ns}	45 ^{ns}
Error	72	1.60	7.02	0.20	1.30	0.21	0.11

Levels of significance are: * $p < 0.05$, ** $p < 0.01$ and Ns: non significant by ANOVA.

and Skoog (MS) (1962). The form and amount of nitrogen in the *in vitro* medium have significant effects on the rate of cell growth, differentiation and totipotency (Kirby et al., 1987). In the media which nitrate is the only source of nitrogen, often more alkaline state is established with time. This situation slightly improved by adding ammonium salt. pH controlling not only is reason for using both form of nitrate and ammonium in the medium, but also free ammonium ions are toxic. Also optimum ratio of $\text{NH}_4^+:\text{NO}_3^-$ is effective on morphogenesis of many plants (Bonga and Von Aderkas, 1992). Sharafi (2010) was found the best *in vitro* medium for hawthorn and dog rose pollen germination and tube growth in different researches. In this research the effect of inorganic nitrogen sources (NH_4NO_3 and KNO_3) were studied on proliferation stage of dog rose.

MATERIALS AND METHODS

After disappearing chilling requirement of buds in February, 2010, axillary buds of dog rose (which are grown in botanical garden of University of Tabriz) were cut and then placed under running tap water (for 1 h) and washed for 15 min, with Tween 80 (0.1%) and then decontaminated with 70% Ethanol (for 5 min), 0.1% (w/v) solution of Mercury (II) Chloride (for 2 min), Sodium Hypochlorite (for 20 min). Finally; all the explants were washed three times by sterile distilled water. Cefotaxime 250 mg. l⁻¹ and Tetracycline 100 mg. l⁻¹ were used through direct addition on establishment medium for bacterial decontaminations. However; explants were transferred to the prepared medium after sterilization. MS (Murashige and Skoog's, 1962) basal medium was used for explants establishment stage which was contained 30 g. l⁻¹ sucrose, 0.8% Agar, 1 mg. l⁻¹ GA₃ and 1 mg. l⁻¹ BAP. In proliferation medium, MS medium was modified with combination of both concentrations of NH_4NO_3 (10, 15 and 20 mM) and KNO_3 (15, 20 and 25 mM) at the ratio of $\text{NH}_4^+:\text{NO}_3^-$ (Table 1) where supplemented with 20 g. l⁻¹ sucrose, 10 g. l⁻¹ sorbitol, 0.8% Agar, 5 mg. l⁻¹ AgNO₃, 1 mg. l⁻¹ GA₃ and 3 mg. l⁻¹

BAP. The pH of media was adjusted to 5.8 then culture medium was autoclaved at 121 °C at 105 kPa for 20 min. Explants were incubated in the culture room at temperature of $25 \pm 2^\circ\text{C}$ and photoperiod of 16 h light and 8 h dark cycle. In proliferation stage, after four weeks shoot length, node number, axillary shoot percentage, chlorosis leaf number, necrotic leaf number and feeble callusing percentage at the cut ends of explants were recorded. All experiments were carried out as in completely randomized design (CRD) with 8 replications. Data were analyzed with SPSS software (ver. 16) and comparison of means was performed by Duncan's New Multiple Range Test at $P < 0.05$.

RESULTS AND DISCUSSION

Analysis of variance showed that, shoot length significantly influenced by different concentrations of NH_4NO_3 and $\text{NH}_4\text{NO}_3 \times \text{KNO}_3$ interaction ($\text{NH}_4^+:\text{NO}_3^-$ ratio) ($p < 0.05$) (Table 2). The highest shoot length was observed in 20:45 ratio of $\text{NH}_4^+:\text{NO}_3^-$. In this ratio the amount of total nitrogen in culture medium was high (65 mM). The lowest shoot length was observed in 15:30 $\text{NH}_4^+:\text{NO}_3^-$ ratio (total N 45 mM) (Figure 1A). In comparison of 20:45 and 15:30 $\text{NH}_4^+:\text{NO}_3^-$ ratio; more concentration nitrate ion and high total nitrogen may be the main reason of increasing shoot length. The high levels of ammonium ion may be inhibited growth of shoot length due to decrease the activity of Nitrate Reductase and Glutamate Syntheses Enzymes in producing of amino acids (Gamborg and Shyluk, 1970). So nitrate is the most important form of nitrogen that used by the dog rose. Moreover; results showed that the shoot length was increased with rising of total nitrogen in different treatments. Nitrogen is a major element for amino acid, protein and nucleic acid biosynthesis and has important roles in many metabolism pathways, cell growth and




Figure 1. Effect of NH₄⁺:NO₃⁻ ratio (mM) in MS media on the shoot length (A), node number (B), chlorosis leaf number (C) of Dog rose after 4 weeks in culture. Bars with common letters are not significantly different at $p < 0.05$, according to LSD test.

differentiation. The deficiency of nitrogen quickly inhibits plant growth and development (Taiz and Zeiger, 2002).

The different concentrations of NH₄NO₃ and interaction of NH₄NO₃ × KNO₃ ($p < 0.05$) (Table 2) were influenced significantly node numbers. Node numbers in 20:45 NH₄⁺:NO₃⁻ ratio was maximum and it was minimum in 10:30 and 10:25 NH₄⁺:NO₃⁻ ratios (Figure 1B).

Nitrate is essential for axillary meristem and leaves formation. So; omission of nitrate from the medium, inhibit axillary meristem and leaves formation (Ramage and Williams, 2001). Moreover, in 10:25 and 10:30 NH₄⁺:NO₃⁻ ratios less node number was observed, because total nitrogen (35 and 40 mM respectively) was low in comparison with 20:45 NH₄⁺:NO₃⁻ ratio (total N 65 mM). Balancing of NH₄⁺ and NO₃⁻ ions in medium is necessary for better absorption of NO₃⁻ ion and inhibiting of NH₃ toxicity inside the tissues. Therefore; cell growth is

depending on optimum ratio of NH₄⁺:NO₃⁻ in medium (Lenee and Chupeau, 1989). This is in accordance with the results obtained by Ivanova and Van staden in *Aloe polyphylla* (2009).

Axillary shoot percentage was influenced significantly by different concentrations of NH₄NO₃ ($p < 0.05$) (Table 2) and interaction of NH₄NO₃ × KNO₃ was insignificantly influenced on axillary shoot percentage. Axillary shoot percentage increased with increasing of NH₄NO₃ from 10 to 20 mM. Minimum and maximum axillary shoot percentage was observed in 10 and 20 mM concentration respectively. There was insignificant difference between two concentrations of NH₄NO₃, 10 and 15 mM, for axillary shoot percentage (Figure 2). Chlorosis leaf number was influenced significantly by different concentrations of KNO₃ and interaction of NH₄NO₃ × KNO₃ ($p < 0.05$) (Table 2). Highest chlorosis leaf number was observed with




Figure 2. Effect of NH₄NO₃ concentrations in the culture media on the axillary shoot percentage. * Bars with common letters are not significantly different at p<0.05, according to LSD test.




Figure 4. Effect of NH₄NO₃ concentrations in the culture media on feeble callusing percentage at the cut ends of explants. * Bars with common letters are not significantly different at p<0.05, according to LSD test.




Figure 3. Effect of KNO₃ concentrations in the culture media on the necrotic leaf number. * Bars with common letters are not significantly different at p<0.05, according to LSD test.

increase NH₄⁺:NO₃⁻ ratio (Figure 1C). High ratio of NH₄⁺:NO₃⁻ in medium can be reason for leaves chlorosis, so that with cell inside ammonium ion ethylene produced increased and following this phenomenon; senescence of leaves was occurred (Barker et al, 1987). The amount of total nitrogen was reported less important than NH₄⁺:NO₃⁻ ratio for chlorosis leaf which may be due to the NH₄⁺ and NO₃⁻ ions role in changing of media pH and loss absorption of nutrient elements (Ramage and Williams, 2001). When total nitrogen on medium is high, some of the leaves are suffering from chlorosis due to total nitrogen deficiency in the medium. Necrotic leaf number was influenced significantly by different concentrations of KNO₃ (p<0.05) (Table 2). Interaction of NH₄NO₃× KNO₃ was none significant on necrotic leaf number. Maximum necrotic leaf number was observed in the 25 mM concentration of KNO₃ (Figure 3). High concentration of K⁺ in the medium can be lead to leaf necrotic, so that deficiency of calcium occurred with the more K⁺ inside tissues and deficiency of calcium can be lead to leaf necrotic (Barker et al, 1987). Different concentrations of KNO₃ were significantly influenced on feeble callusing percentage at the cut ends of explants (p<0.05) (Table 2) and interaction of NH₄NO₃× KNO₃ was none significantly influenced on feeble callusing percentage. Minimum bottom callus formation observed at 25 mM concentration of KNO₃ (Figure 4), KNO₃ in high concentration may be inhibited in the form of undifferentiated cells. This result is in accordance with the results obtained by Salehi shanjani (2003) in *Juniperus excels*.

Conclusion

In Dog rose optimum $\text{NH}_4^+:\text{NO}_3^-$ ratio affected shoot length, the node number and chlorosis of leaf. Ammonium ion was effective on produce axillary shoot and nitrate ion was effective on necrotic leaf number and feeble callusing percentage at the cut ends of explants. Various forms of N supplied overlooked importance of this nutrient in plant growth and morphogenesis *in vitro*.

ACKNOWLEDGEMENTS

The authors are thankful to Zeinab Khodajoo-Sefat and Gholamreza Gohari for their critical suggestions and helps.

REFERENCES

- Barker AV, Arker K, Corey A (1987). Ammonium-induced ethylene evolution by horticultural crops. *Hort. Sci.*, 22: 381.
- Bonga JM, Vonaderkas P (1992). *In vitro* Culture of Trees. Kluwer Academic Pub., Dordrecht, Boston, London, p. 255.
- Chu CY, Hu SL, Knightt MA, Smith L (1993). Effect of liquid culture on the growth and development of miniature rose (*Rosa chinensis*). *Plant Cell Tiss. Organ. Cult.*, 32: 34-329.
- Gamborg OL, Shyluk JP (1970). The culture of plant cells with ammonium salts as a sole nitrogen source. *Plant Physiol.*, 45: 598-600.
- George EF (1993). Plant propagation by tissue culture. Part 1: The technology. 2nded. Exegetics Ltd, England, pp. 654-670.
- Horn W (1992). Micropropagation of rose (*Rosa L.*). In: Y. P. S. Bajaj, editor. *Biotechnology in Agriculture and Forestry*. Vol. 20, High-tech and micropropagation IV. Germany: Springer, pp. 320-342.
- Horn W, Sohlegel G, Karin J (1988). Micropropagation of *Rosa (Rosa hybrida)*. *Acta Hortic.*, 228: 623-626.
- Iuncna A (2005). A guide to medicinal plants in North Africa. Center for mediterranean cooperation, Malaga (Spain), p. 256.
- Ivanova M, Vav SJ (2009). Nitrogen source, concentration, and $\text{NH}_4^+:\text{NO}_3^-$ ratio influence shoot regeneration and hyperhydricity in tissue cultured Aloe polyphylla. *Plant Cell Tiss. Organ. Cult.*, 99: 167-174.
- Khosh-Khui M, Sink ICA (1982). Micropropagation of new and old world rose species. *J. Hortic. Sci.*, 57(3): 9-315.
- Kirby EG, Leutek E, Lee MS (1987). Nitrogen nutrition. In: J. M. Bonga, D. J. Durzan. (eds.), *Cell and Tissue Culture in Forestry*. Martinus Nijhoff Publishers, Dordrecht, Boston, Lancaster, 1: 237.
- Lenee P, Chupeau Y (1989). Development of nitrogen assimilating enzymes during growth of cells derived from protoplasts of sunflower and tobacco. *Plant Sci.*, 59: 109-117.
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.*, 15: 97-473.
- Nikbakht A, Kafi M, Mirmasoudi M, Babalar M (2005). Micropropagation of Damask Rose (*Rosa damascena* Mill.) cvs Azaran and Ghamsar. *Int. J. Agric. Biol.*, 7: 535-538.
- Ramage CM, Williams RR (2001). Mineral nutrition and plant morphogenesis. *In vitro Cell. Dev. Biol. Plant*, 38: 116-124.
- Rout GR, Samantaray S, Mottley J, Das P (1999). Biotechnology of the rose: A review of recent progress. *Sci. Hortic.*, 81: 28-201.
- Salehi- Shanjani P (2003). Nitrogen effect on callus induction and plant regeneration of *Juniperus excels*. *Int. J. Agric. Biol.*, 4: 419-422.
- Skirvin RM, Chu MC, Hu H, Young J (1990). Rose. In: P. V. Ammirato, D. R. Evans, W. R. Sharp, Y. P. S. Bajaj editors. *Handbook of Plant Cell Culture*. Vol. 5. New York: McGraw Hill, New York: Springer, pp. 716-743.
- Sharafi Y (2010). Biological characteristics of pollens in some genotypes of *Rosa canina* as main factors affecting fruit set. *Afr. J. Med. Plants Res.*, 4(20): 2173-2175.
- Sharafi Y (2010). Suitable *In vitro* medium for studding pollen viability in some of the Iranian hawthorn genotypes. *J. Med. Plants Res.*, 4(19): 1967-1970.
- Taiz E, Zeiger L (2002). *Plant Physiology*. 3rded. publisher Sinauer Associates, p. 690.