

## Full Length Research Paper

## Biological and antioxidant activity of *Gunnera tinctoria* (Nalca)

Pedro Zamorano<sup>1</sup>, Benjamín Rojano<sup>2</sup>, Marcela Morales<sup>2</sup>, Haroldo Magariños<sup>3</sup>, Patricio Godoy<sup>4</sup> and Ociel Muñoz<sup>3\*</sup>

<sup>1</sup>Graduate School, Faculty of Agricultural Science, Universidad Austral de Chile, Valdivia, Chile.

<sup>2</sup>Laboratorio Ciencia de los Alimentos, Facultad de Ciencias, Universidad Nacional de Colombia, Medellín, Colombia.

<sup>3</sup>Food Science and Technology Institute (ICYTAL), Faculty of Agricultural Sciences, Universidad Austral de Chile, Campus Isla Teja s/n, Valdivia, Chile.

<sup>4</sup>Clinical Microbiology Institute, Medicine Faculty, Universidad Austral de Chile, Valdivia, Chile.

Received 23 March, 2017; Accepted 26 April, 2017

The endemic Chilean edible plant *Gunnera tinctoria* (Nalca) is highly appreciated in the south of Chile by the small farmers. Nevertheless, no background exists about his secondary metabolites. In the present study, in the leaf from *G. tinctoria* was investigated the content of bioactive compounds like coumaric acid, ascorbic acid and total phenols; the antioxidant capacity was evaluated by 3-ethylbenzothiazoline-6-sulfonic acid (ABTS), oxygen radical absorbance capacity (ORAC), ferric reducing antioxidant power), and 2,2-diphenyl-1-picrylhydrazyl (DPPH) methods reducing sugars were measured. Finally, the biological activity was evaluated against *Cladophialophora* and *Cryptococcus laurentii*. The results suggest that the most abundant constituent in the extract were catechin (1344.97 mg/100 g dry weight) and epicatechin (1429.28 mg/100g dry weight), and was confirmed and quantified by high performance liquid chromatography (HPLC-PDA); while the ORAC methodology showed a high antioxidant capacity (192000.0±5.91 umol Trolox Eq/100 g dry weight). On the other hand, the extract had a fungicide effect against both microorganism assayed, inhibiting the growth of *Cladophialophora*'s mold- and the yeast *Cryptococcus laurentii*. This is the first report of antioxidant capacity, bioactive compounds and biological activity of *G. tinctoria*, and these findings suggest that an extract prepared from the Nalca leaf may be a promising source of antioxidant and bioactive compounds and as a research object by being an antifungal and therapeutic alternative in development.

**Key words:** *Gunnera tinctoria*, Nalca, antioxidant capacity, biological activity, nutraceutical.

### INTRODUCTION

Numerous studies have established that a high consumption of fruits and vegetables leads to an important intake of bioactive compounds (Torres et al., 2011); which are strongly associated with the reduction in

\*Corresponding author. E-mail: [ocielmunoz@uach.cl](mailto:ocielmunoz@uach.cl).

the incidence of cardiovascular diseases (Pounis et al., 2013), various types of cancer (Zamora-Ros et al., 2013), type II diabetes (Anhê et al., 2013), immune disorders and neurodegenerative disorders (Marszalek et al., 2017), at the gastrointestinal level (Ozidal et al., 2016), and in general in all diseases in which oxidative stress has an important role (Quiles et al., 2006). Among importance bioactive compounds are p-coumaric acid, which has hepatoprotective (Calil Brondani et al., 2017) and UV-protection qualities (Aguilar-Hernández et al., 2017). Flavanols like catechin and epicatechin are acknowledged for their antibacterial capacity (Hamilton-Miller, 1995), anti-inflammatory activity and cancer preventives for humans (Fujiki et al., 2002), and ascorbic acid which can reduce the malignant potential of melanoma cells (Fischer and Miles, 2017). Also, these secondary metabolites can inhibit the growth of various mold, bacteria and yeast species (Gramza and Korczak, 2005; Almajano et al., 2008), acting as antimicrobials (Han et al., 2007).

Gunneracea is a monogeneric family (Genus *Gunnera* L.), of which 6 subgenera, with approximately 65 species on the planet have been described (Bergman et al., 1992). Its distribution extends to Central and Southern Africa, Madagascar, New Zealand, United Kingdom, Tasmania, Indonesia, Philippines, Hawaii, Mexico, Venezuela, Colombia, Chile and Argentina; having been reported in South America 23 species in Colombia, 12 in Chile, 7 in Ecuador, 6 in Peru, 5 in Argentina and Bolivia, 3 in Venezuela, 2 in Brazil and one in Uruguay, including *Gunnera tinctoria* (Mol.) Mirb., *Gunnera magellanica* Lam., *Gunnera brephogea* Linden and *Gunnera manicata* (Bergman et al., 1992; Wanntorp and Klackenberg, 2006).

*G. tinctoria* (Molina) Mirb. (*Gunnera chilensis*), commonly known as Nalca, Chilean Rhubarb or Pangue, is a non-timber forest product (NTFPs) (Valdebenito et al., 2003) of recognized importance in southern Chile, especially at smallholder agriculture. Which predominates subsistence exploitation, since it also provides resources and higher incomes to rural people and/or groups that are engaged in the collection and informal sale of this wild plant (Petzold et al., 2006). In South America, this species is distributed from Chile and Argentina, to Colombia and Venezuela; it has also been reported in Peru, Ecuador and Bolivia (EPPO, 2014). It is worth mentioning that in both New Zealand and the UK, *G. tinctoria* seeks to be controlled and eradicated, since it is classified as an invasive species of rapid propagation and difficult management (Hickey and Osborne, 1998, Williams et al., 2005).

In Chile, it is one of the few ornamental plants that has an edible petiole, so it is widely consumed during the spring and summer seasons. Its leaves can be consumed directly as an infusion, or they are used in the native kitchen of the island of Chiloé (Chile) being used to cover other foods during their cooking. It has been described that the maceration of its flowers and leaves constitutes a

refreshing infusion. On the other hand, in medicinal terms (at the popular and rural level) it is known as an antifebrile, and applied in hemorrhages, dysentery, diarrhea, mouth, throat, liver and stomach affections (Montes and Wilkomirsky, 1985; Valdebenito et al., 2003). It has also been described that, at the agronomic level, *G. tinctoria* is a very good nitrogen-fixing species, which could be useful for the recovery of soils depleted in this nutrient (Troncoso et al., 2013).

This study is the first work that seeks to determine the nutritional characteristics, antioxidants and effects against microorganisms of *G. tinctoria*. The objective of this research was to characterize this plant, both from the nutritional point of view and the bioactive compounds, determining the antioxidant capacity of the *G. tinctoria* leaf, as well as determining the content of reducing sugars, total polyphenols and related metabolites directly with antioxidant activity, specifically Catechin, Epicatechin and p-coumaric acid, in addition to evaluating its antifungal effect on *Cryptococcus laurentii* and *Cladophialophora*.

## MATERIALS AND METHODS

### Reagents and equipment

Diphenyl-1-2,2 picrylhydrazyl (DPPH); 2,2'-azobis (2-amidino-propane) (AAPH); fluorescein; 6-hydroxy-2,5,7,8-tetramethyl chromane-2-carboxylic acid (trolox); 2,4,6-tris-(2-pyridyl-s-triazine) (TPTZ); 3,4,5-trihydroxybenzoic acid (gallic acid) and (+)-catechin and (-)-epicatechin standards were obtained from Sigma-Aldrich® (St. Louis, MO, USA). Sodium carbonate and folin carbonate were purchased from Merck (Germany). Methanol and other solvents were purchased from Fisher Scientific Co. (Fair Lawn, NJ, USA). Ultraviolet-visible measurements were performed on a Multiskan Spectrum spectrophotometer (Thermo Scientific). The decrease in fluorescence intensity measured in the ORAC assay was performed on a Perkin-Elmer LS-55 (U.K.) Spectrofluorimeter. The chromatographic assays were carried out on a Shimadzu® series Prominence® UFLC liquid chromatograph.

### Vegetal material

The plant material was collected in the commune of Los Lagos -Los Ríos Region, Chile- (31 m.s.n.m and 12.3°C average temperature) (IPNI, 2012; The plant list, 2013). After being stored in perforated polypropylene bags were lyophilized and taken to the laboratory for the respective analyzes.

### Preparation of the sample

For the antioxidant and secondary metabolite assays, 2.0 g of samples dissolved and homogenized with 20 ml of acidified water (1% v/v HCl) were weighed into an Ultra-Turrax (IKA-WERK®). The extract was centrifuged at 5000 rpm for 12 min at room temperature, taking the supernatant and making the appropriate dilutions with type I water. For the biological activity tests 2.5 g of samples dissolved and homogenized were weighed with 25 ml of MeOH-Ac. Formic (99:1). The extract was centrifuged at 5000 rpm for 12 min at room temperature. The solutions were stored at 4°C

until analysis.

## Chemical and antioxidant capacity

### Proximal analysis

Total ash (Bateman, 1970; AOAC, 1996; method 942.05), moisture (Bateman, 1970; AOAC, 1996; method 930.15), crude protein (AOAC, 1996; method 990.03), ethereal extract (Bateman, 1970; AOAC, 1980; method 832p) and crude fiber (AOAC, 1996; method 978.10) were determined. Carbohydrates, was determined by difference. Each sample was analyzed in triplicate, and all the proximate values were reported in percentage.

### Determination of reducing sugars - DNS colorimetric method

The determination of reducing sugars was performed as described by Najmus and Whitney (2011). 500  $\mu$ L of a 1% solution of 3,5-dinitrosalicylic acid in NaOH is added to 500  $\mu$ L of the sample to be evaluated. The mixture was heated at 90°C for 5 min and its absorbance determined at 540 nm. The contents were determined by comparing with a standard curve using glucose as standard. The results were expressed as mg of equivalent glucose/100 g dry weight.

## Content of secondary metabolites with antioxidant properties

### Total phenols

The determination of phenols was done by the colorimetric method of Folin-Ciocalteu designed by Singleton and Rossi (1965). 50  $\mu$ L of sample were added to 125  $\mu$ L of Folin's reagent and 400  $\mu$ L of 7.1% (w/v) sodium carbonate, adjusting with distilled water to 1000  $\mu$ L, the spectrophotometric reading was performed at 760 nm and compared with Standard curve using gallic acid as standard. The results were expressed as mg gallic acid equivalent: GAE/100 g dry weight.

### Determination of ascorbic acid

The determination of ascorbic acid was performed through HPLC, following the method described by Kelebek et al. (2009). The aqueous extract was filtered with cellulose acrodiscs of pore size 0.45  $\mu$ m. The samples were analyzed on a Shimadzu® liquid chromatograph model LC-20AD, equipped with a SIL-20A/HT auto injector, a CBM-20A communication module and a PDA detector SPD-M20A; a Merk® (250 mm long and 4 mm internal diameter) LiChrospher® 100 C-18 (5  $\mu$ m, 250 mm  $\times$  4.6 mm) column. As a mobile phase 0.1% formic acid in purified water was used, by isocratic elution system, at a temperature of 35°C and flow of 0.8 mL/min. Detection was performed at 245 nm. The identification and quantification was done with calibration curves elaborated with different concentrations of ascorbic acid.

### Determination of (+)-catechin and (-)-epicatechin

Confirmation and quantification of (+)-catechin and (-)-epicatechin were analyzed by direct injection of the samples, previously filtered through a 0.45  $\mu$ m pore-size nylon filter, in a liquid chromatography (HPLC-DAD) using a Shimadzu LC-20AD/T HPLC equipped with a SPD-6AUV detector (Kyoto, Japan) and a Pinnacle (II) C-18 column (5  $\mu$ m) 250  $\times$  4.6 mm (Restek®, Bellefonte, USA) with an

autoinjector and a photodiode array detector (PDA). (+)-catechin and (-)-epicatechin were adopted as the standard of identification and quantification at 280 nm. The mobile phase was methanol (A) acidified water (0.1% formic acid) (B), with gradient elution of 0.01 min 60% A was used; 5-12 min 80% A; 13-14 min 60% A. Flow rate of mobile phase was 1.0 mL/min (Oliveiro et al., 2009).

### Phenolic acids

The p-coumaric phenolic acid was determined by direct injection of the samples, previously filtered through a 0.45  $\mu$ m pore-size nylon filter, in a HPLC-DAD using a Shimadzu LC-20AD/T HPLC equipped with a SPD-6AUV detector (Kyoto, Japan) and a Pinnacle (II) C-18 column (5  $\mu$ m) 250  $\times$  4.6 mm (Restek®, Bellefonte, USA) with an autoinjector and a photodiode array detector (PDA). P-coumaric acid was adopted as the standard for its identification at 320 nm. The mobile phase was a mixture of acetonitrile, acidified water (phosphoric acid at pH = 2.5) (40:60) v/v, supplied at a rate of 0.8 mL/min (Kelebek et al., 2009).

## Assessment of antioxidant capacity

### Ferric reducing/antioxidant power assay (FRAP)

This method evaluates the reducing potential of a sample based on its ability to reduce ferric iron ( $\text{Fe}^{+3}$ ) present in a complex with 2,4,6-tris (2-pyridyl-s-triazine) (TPTZ) to the ferrous form ( $\text{Fe}^{+2}$ ). The absorbance was determined at 593 nm. In brief, 50  $\mu$ L of sample were added to 900  $\mu$ L of pH 3.4 acetate buffer, TPTZ,  $\text{FeCl}_3$ , in a 10:1:1 ratio; after 30 min of reaction the absorbance was determined, this value was compared to the reference curve constructed with ascorbic acid as the standard, and the results were expressed as AEAC (Ascorbic Acid Equivalent Antioxidant Capacity: mg ascorbic acid per 100 g dry weight) (Benzie and Strain, 1996).

### DPPH assay

Radical scavenging activity against the stable radical DPPH was measured using the methods proposed by Brand-Williams et al. (1995), with certain modifications. The method is based on the reaction of 10  $\mu$ L of sample with 990  $\mu$ L of DPPH solution for 30 min at room temperature. The absorbance decrease, associated with a reduction in the DPPH concentration, was measured at 517 nm. The results were expressed in  $\mu$ mol trolox equivalents per 100 g dry weight.

### ABTS•+ assay

The method of Re et al. (1999) was used with some modifications. 10  $\mu$ L of sample was added to 990  $\mu$ L of a solution of ABTS•+. The ability of the samples to trap the cationic radical ABTS•+ was evaluated by means of the decrease in absorbance read after 30 min of reaction at a wavelength of 732 nm. The absorbance value was compared with the reference curve constructed with Trolox as the primary standard, and the results were expressed as TEAC values ( $\mu$ mol Trolox per 100 g dry weight).

### Oxygen radical absorbance capacity (ORAC) assay

This methodology is used to evaluate the ability of a sample to trap peroxy radicals ( $\text{ROO}\cdot$ ). The method described by Prior et al. (2005) and Romero et al. (2010) was used. 30  $\mu$ L of the sample was

**Table 1.** Physico-chemical composition of *Gunnera tinctoria*.

Proximate composition variable (%)	Nalca leaf
Humidity	90.42 ± 0.31
Ashes	7.23 ± 0.23
Protein	19.95 ± 0.48
Ethereal Extract	1.15 ± 0.1
Crude fiber	12.40 ± 0.2
Carbohydrates***	59.27

\*The results of each analysis are presented as the mean ± standard deviation (n=3). \*\* Results expressed on dry weight. \*\*\* Determination of Carbohydrate by difference.

added to 21 µl of  $1 \times 10^{-2}$  M fluorescein in PBS (75 mM); 2,899 µl PBS (75 mM); and 50 µl of 0.6 M AAPH in PBS (75 mM). The temperature was controlled at 37°C and the pH was maintained at 7.4. The readings were made at an excitation  $\lambda$  493 nm and excitation slit 10 nm; and an emission  $\lambda$  515 nm with an emission slit 15 nm, with 1% attenuator and no attenuator plate. The protective effect of the antioxidant is calculated using the differences in areas under the decay curve of fluorescein between the blank (reaction in the absence of the sample) and the sample. It was compared against the curve of the primary Trolox® standard. The results were expressed as TEAC, µmol equivalents of Trolox per 100 g dry weight according to Equation (1).

$$ORAC = \frac{AUC - AUC^\circ}{AUC_{Trolox} - AUC^\circ} f[Trolox] \quad (1)$$

Where AUC is the area under the curve of the sample,  $AUC^\circ$  area under the curve for the control,  $AUC_{Trolox}$  area under the curve for the Trolox,  $f$  is the dilution factor of the extracts.

## Biological activity

### Fungal strains

Fungal strains of *C. laurentii* and *Cladophialophora* were used for the determination of antifungal activity. The microorganisms used were isolated from a cheese ripening room, belonging to small producers in the region of Los Ríos. Sampling, culturing, isolation and identification were done at the Clinical Microbiology Institute of the Universidad Austral de Chile.

### Evaluation of antifungal activity

#### Suspension preparation

From each strain a suspension was prepared, which consisted of an inoculum of the microorganisms, resuspended in sterile physiological serum. It was homogenized to a turbidity equal to that presented in a 0.5 Mc Farland tube. This comparison was made visually.

#### Susceptibility testing

The susceptibilities of the isolated microorganisms were determined by the modified Kirby-Bauer disc diffusion method with Muller Hinton agar plates (Bauer et al., 1996). All the media used in the

present investigation were obtained from Oxoid, Hampshire, UK.

### Agar well diffusion method

Antifungal activity of *G. tinctoria* was determined by agar well diffusion method (Bauer et al., 1996). To apply this technique, 6 wells were made in the agar (6 mm diameter). Afterwards, the plate was inoculated, where the fungal suspension (0.5 Mc Farland) was absorbed with a sterile cotton swab and then smeared on the agar in three different directions at a 90° angle. In parallel, 5 dilutions were prepared from the pure extract, of which in each well were deposited 25 µl of extract. The final well contained 25 µl of solvent (70% alcohol), which corresponded to the negative control. Later, the plates were incubated at 25 and 32°C (mold and yeast respectively) for 20 days. The experiments were carried out in triplicates.

### Determination of fungal activity

Antifungal activity was assessed by observing and determining the presence and/or absence of inhibition halos formed around the well containing the pure and diluted extracts. The determination of the fungistatic and/or fungicidal effect was done visually considering the time of inhibition of growth in Mueller Hinton agar. These measurements were performed after seven and 20 days.

## RESULTS

The proximal physico-chemical composition of the *G. tinctoria* leaf is shown in Table 1, and Table 2 shows that the Nalca leaf has a significant amount of metabolites, which have been quantified as total gallic acid equivalent polyphenols. Also, reducing sugars and bioactive compounds such as ascorbic acid, epicatechin, catechin and p-coumaric acid were determined. The antioxidant and reducing activity of nalca leaf extracts was characterized by different assays (ABTS, FRAP, DPPH, ORAC). The results of these analyzes can be seen in Table 3.

Due to the high antioxidant potential and the abundant concentration of phenolic compounds, biological activity tests were carried out in which the pure extract exerted a fungistatic and fungicidal effect on yeast *C. laurentii*

**Table 2.** Bioactive compounds and reducing sugars quantified in *G. tinctoria* leaf.

Compound	Nalca leaf
Total phenols (mg gallic ac. eq/100 g DW)	10684.20±128.75*
p-coumaric acid (mg/100 g DW)	48.69
Catechin (mg/100 g DW)	1344.97
Epicatechin (mg/100 g DW)	1429.28
Ascorbic acid (mg/100 g DW)	46.79
DNS (mg glucose eq/100 g DW)	7353.39 ± 0.36*

\* The results of each analysis are presented as the mean ± standard deviation (n=3). \*\* Results expressed in 100 g DW.

**Table 3.** Antioxidant activity in *G. tinctoria* leaf.

Methodology	Nalca leaf
FRAP (mg ascorbic acid/100 g DW)	11194.50±303.42
ABTS (µmol Trolox Eq/100 g DW)	120094.22±5950.44
DPPH (µmol Trolox Eq/100 g DW)	83266.76±3944.77
ORAC-H (µmol Trolox Eq/100 g DW)	192006.07±5911.13

\* The results of each analysis are presented as the mean ± standard deviation (n=3). \*\* Results expressed in 100 g DW.

**Table 4.** Biological activity. Presence or absence of inhibition halos when applying *G. tinctoria* leaf extract against *Cryptococcus laurentii*.

Species/extract	Day 7			Day 20		
	S.E	F.T	F.G	S.E	F.T	F.G
<b><i>Cryptococcus laurentii</i></b>						
Well 1		√			√	√
Well 2		√			√	√
Well 3		-			-	-
Well 4		-			-	-
Well 5		-			-	-
Well 6 (N.C)		-			-	-

\*Well 1: Pure extract. Well 2: 1:10 dilution. Well 3: 1:100 dilution. Well 4: 1:1000 dilution. Well 5: 1:10000 dilution. Well 6: negative control. \*\* (√): Presence of halo (inhibitory effect obtained by the extracts); (-): Absence of halo; N.C: Negative control; SE: No inhibitory effect. FT: Fungistatic effect. FG: Fungicidal effect.

(Table 4 and Figure 1) and *Cladophialophora* mold (Table 5 and Figure 2).


## DISCUSSION

### Physico-chemical characterization

According to Bello (2000) in food or plant species the water content and its location within the matrix are factors that influence specific characteristics such as appearance, texture and color. The percentage of humidity for leaf of *G. tinctoria* is of 90.42% being a value of water content

similar to those reported for kiwi (83.98%) (Morillas-Ruiz and Delgado-Alarcón, 2012), orange (85.70%) Berlitz and Grosh (1999), mango (82.10%) (Moreiras et al., 2006), and slightly higher than those determined for avocado (72.07%) and cherry (75.84%) (Morilla-Ruiz and Delgado-Alarcón, 2012). On the other hand, the total ash, corresponding to the mineral elements of a plant structure, presented a percentage of 7.23%; which is slightly higher than that found in apricot (3,0%), peach (4.0%), cherry (4.7%) (Lazos, 1991), orange (5.02%) and lemon (5.17%) (Ibrahim et al., 2011).

Proteins are macronutrients whose main function in the body is to form and regenerate tissues. In the leaves of



**Figure 1.** *Cryptococcus laurentii*. (a) Microscopic examination (fresh examination, chinese ink), and (b) Effect exerted by leaf extract.

**Table 5.** Biological activity, presence or absence of inhibition halos when applying *G. tinctoria* leaf extract against *Cladophialophora*.

Species/extract	Day 7			Day 20		
	S.E	F.T	F.G	S.E	F.T	F.G
<b><i>Cladophialophora</i></b>						
Well 1		√			√	√
Well 2		-			-	-
Well 3		-			-	-
Well 4		-			-	-
Well 5		-			-	-
Well (N.C)		-			-	-

\*Well 1: Pure extract. Well 2: 1:10 dilution. Well 3: 1:100 dilution. Well 4: 1:1000 dilution. Well 5: 1:10000 dilution. Well 6: negative control. \*\* (√): Presence of halo (inhibitory effect obtained by the extracts); (-): Absence of halo; N.C: Negative control; SE: No inhibitory effect. FT: Fungistatic effect. FG: Fungicidal effect.

*G. tinctoria* a value of  $19.95 \pm 0.48\%$  was found which is much higher than that reported for fruit pulps such as cherry (4.23-5.91%) (Mahmood et al., 2013), orange (6.40%) and lemon (7.04%) (Ibrahim et al., 2011). The difference between the values obtained for leaf and pulp is based on the fact that the role of proteins in living organisms is mainly structural. The ethereal extract or lipid fraction recorded a percentage of 1.15% which is considerably lower than those of cherry (6.39%), kiwi (4.36), plum (3.49%), the order of orange (0.33%) and papaya (0.16%) (Morillas-Ruiz and Delgado-Alarcón, 2012).


For crude fiber, a percentage of 12.40% was determined, which is similar to those reported for cherry (9.50%) (Lazos, 1991), and avocado (9.11%) (Morillas-Ruiz and Delgado-Alarcón, 2012); and considerably higher than orange (3.30%), kiwi (5.71%) and plum (3.0%) (Morillas-Ruiz and Delgado-Alarcón, 2012). Since raw fiber is the non-digestible portion of food and consists of cellulose, hemicellulose and lignin, this finding gives

the *G. tinctoria* leaf a high dietary value.

Although the leaves of one plant species are not comparable with the pulp of other matrices (because they are different structural zones); in the case of *G. tinctoria* it is important to make such comparisons (in addition to highlighting its high percentage of raw fiber, high protein content and low ethereal fraction) since in the peasant culture of southern Chile the leaf is consumed directly as a salad and/or as infusions or beverages.

### Secondary metabolites

The phenolic compounds constitute a group of secondary metabolites that present antioxidant capacity with multiple biological benefits for the human being. In the present work the concentration determined for polyphenols was  $10684.20 \pm 128.75$  mg gallic acid/100 g DW; Similar to that reported for Curuba (10584.7 mg gallic acid/100 g DW), and higher than blackberry (1864.4 mg gallic



**Figure 2.** *Cladophialophora*. (a) Microscopic examination (fresh examination, Chinese ink), and (b) effect exerted by leaf extract.

acid/100 g DW), strawberry (1638.4 mg gallic acid/100 g DW), mortiño (1237.2 mg acid (Gallon acid/100 g DW), red apple (426.7 mg gallic acid/100 g DW), grape (290.6 mg gallic acid/100 g DW), cauliflower (440.3 mg gallic acid/100 g DW), kiwi (199.3 mg gallic acid/100 g DW) and carrot (160.0 mg gallic acid/100 g DW) (Zapata et al., 2014).

Compared to leaves of *Swietenia macrophylla*, *Swietenia humilis* and *Swietenia mahagoni*, the total polyphenol content of Nalca is vastly higher (10684.20 mg gallic acid/100 g DW versus 290.5; 250.91, and 146.4 mg gallic acid/100 g DW respectively) (Preciado et al., 2016). It also exceeds the concentrations reported in *Vaccinium meridionale Swart* (Green tea: 85.6 mg gallic acid/100 g DW and Black tea: 189.8 mg gallic acid/100 g DW) (Zapata-Vahos et al., 2015); and, to those of *Ilex paraguariensis* (commonly known as yerba mate) and *Ilex laurina*, which present a minor concentration (10.78 mg gallic acid/100 g DW and 23.70 mg gallic acid/100 g DW, respectively) (Perez et al., 2014).

Usually the leaf is one of the structural parts of plant species with higher content of bioactive compounds, since the synthesis of them is recognized as a necessity by supporting filter functions and protection against ultra violet light (Morales et al., 2017), support mechanical and intervention in the ecological interactions between the plant, its environment and predators, among others (Zobel et al., 1999, Chaves and Escudero, 1999, Zoratti et al., 2014, Ramakrishna and Ravishankar, 2011).


p-coumaric acid (pCA) is present in a wide variety of edible plants, fruits and vegetables (Clifford, 2000). It has been reported to have antioxidant, antimicrobial, anti-inflammatory properties, and it has also been suggested that its participation reduces the risk of stomach cancer (Ferguson et al., 2005) through the reduction of carcinogenic nitrosamines (Kikugawa et al., 1983). It was determined by HPLC-UV, obtaining a value of 38.41 mg/L. Similar to extracts of yerba mate (*Ilex laurina*):

47.32 mg/L; and that of *Ilex paraguariensis* 24.44 mg/L (Pérez et al., 2014). In terms of dry basis (48.69 mg p-coumaric acid/100 g DW) the results of *G. tinctoria* are much higher than those reported for leaves and ginger root: *Boesenbergia pulchella var attenuata*, *Boesenbergia armeniaca* (each with 4.0 mg/100 g DW) (Jing et al., 2010); and greater than Turmeric (*Curcuma longa*) (5.96 mg /100 g DW) (Wojdylo et al., 2007).

The metabolites catechin and epicatechin due to their antioxidant activity are reported as inhibitors and/or retarders of various types of cancer-skin, breasts, prostate and lung (Yang et al., 2002); and for having shown the capacity of arrest the cell cycle and induce apoptosis in cancerous cells (Gupta et al., 2010; Chena et al., 2010). The calculated values for these bioactive compounds were 1060.98 and 1127.49 mg epicatechin/L, respectively (Figure 3). In terms of dry basis, concentrations of 1344.97 mg catechin/100 g DW and 1429.28 mg epicatechin/100 g was determined. These results are higher than those reported for black tea from Mortiño leaves (*V. meridionale Swartz*): 437-972 mg of catechin/100 g DW and 470-1014 mg epicatechin/100 g DW (Zapata-Vahos et al., 2015). On the other hand, compared to the quantified for *Camellia sinensis* (green tea), the leaf extract of *G. tinctoria* showed higher values of both catechin (1060.98 mg/L v/s 270 mg/L) and epicatechin (1127.49 mg/L v/s 738mg/L) (Moore et al., 2009).

### Antioxidant capacity

One of the most reliable methodologies for measuring antioxidant capacity is DPPH; since it quantifies only antioxidant compounds, and also presents a high selectivity due to its high molecular impediment, so that the antioxidants that neutralize this radical must be very efficient (Prior et al., 2005; Osman et al., 2006).



**Figure 3.** HPLC-PDA chromatograms at 280 nm of (1) catechin, (2) epicatechin from Nalca leaf collected in the commune of Los Lagos, XIV region of Chile.

In this investigation the antioxidant capacity of *G. tinctoria* was 83266.76  $\mu\text{mol Trolox}/100\text{ g DW}$ . On the other hand, the antioxidant activities described for *S. macrophylla*, *S. mahagoni* and *S. humilis* (Preciado et al., 2016) are largely overcome by the leaves of Nalca (83266.76  $\mu\text{mol Trolox}/100\text{ g DW}$  versus 16190; 10710 and 8900  $\mu\text{mol Trolox}/100\text{ g DW}$  respectively). As well as it is larger and in the order of the extracts made from green tea leaf and black tea of mortiño (49158 and 111815  $\mu\text{mol Trolox}/100\text{ g DW}$ , respectively) (Zapata-Vahos et al., 2015).

Considering the characteristics and nature of the bioactive compounds it is necessary to combine and execute other techniques that measure the antioxidant capacity of these compounds. In this way, the ABTS methodology has been used extensively to determine it. This test is based on the quantification of the discoloration of the ABTS $\cdot+$  radical, due to the interaction with hydrogen or electron donor species (Moon and Shibamoto, 2009). According to Lu et al. (2010) the ABTS methodology is a fast and stable technique, which is applied mainly to antioxidants of an aqueous nature. Results obtained from ABTS for Nalca leaf (*G. tinctoria*) indicate a concentration of 120094.22  $\mu\text{mol Trolox}/100\text{ g DW}$ . Values higher than those obtained by Kopjar et al. (2015), who in Croatia reported for infusions of yellow,

green and black tea leaves concentrations of 40733, 32815 and 27032  $\mu\text{mol Trolox}/100\text{ g DW}$  respectively; at 1810  $\mu\text{mol Trolox}/100\text{ g DW}$  reported for leaf of *Punica granatum L.* (Pomegranate) (Elfalleh et al., 2012).

In the comparison of the results of the antioxidant activity obtained by both methods, it can be observed that the values for DPPH are lower than those obtained in ABTS. According to Heim et al. (2002), the OH groups at the 3'-, 4'-, 5'- positions of the B ring of flavonoids increase the antioxidant activity of the compound (unlike phenolic compounds with a single group Hydroxyl). And, from the point of view of free radicals, these cause different reactivity with phenolic compounds depending on their structure (Singleton and Rossi, 1965). According to this the ABTS radical has low selectivity in reactions with hydrogen donor atoms (it reacts with any hydroxylated aromatic compound independent of its real antioxidant potential) (Campos and Lissi, 1997). Unlike DPPH, it does not react with flavonoids lacking OH groups in a B ring, nor does it with aromatic acids containing only one OH group (Roginsky and Lissi, 2005). In addition, the results of the study clearly suggest that the antioxidant compounds present in *G. tinctoria* leaves are highly hydrophilic and therefore more sensitive to the ABTS technique.

Through the FRAP assay, the presence of metabolites

with the capacity to stabilize free radicals through the electron transfer mechanism (SET) is expressed. The leaf extract showed an antioxidant capacity of 11194.50 mg ascorbic acid/100 g DW; higher by 42.06% than that reported for Mortiño (*V. meridionale*) green tea leaf (7880 mg Ascorbic acid/100 g DW) (Zapata-Vahos et al., 2015). And, of the order of black tea leaf of Mortiño (15780 mg ascorbic acid/100 g DW) (Zapata-Vahos et al., 2015).

The antioxidant efficacy determined by this assay depends on the redox potentials of the compounds present in the extract, so that, from the study results, it is feasible to indicate that the high reducing capacity of polyphenols is related to the degree of hydroxylation and conjugation of the polyphenol compounds (Pulido et al., 2000).

According to Prior et al. (2005), both the FRAP and Folin-Ciocalteu methods to quantify total phenols are capable of detecting any reducing substance, presenting interference with, among others, free reducing sugars and organic acids such as ascorbic acid. In order to compare this information, the concentration of reducing sugars was quantified by means of the colorimetric technique with DNS (3,5 dinitroxylic acid), which was 7353.39 mg equivalent glucose/100 g DW; and by HPLC technique a Vitamin C value of 46.79 mg ascorbic acid/100 g DW.

Since the total phenol test showed results much higher than those reported by other studies, and that on the contrary FRAP was of the order of the mentioned vegetal species; it is possible that the compounds involved were more sensitive to basic pH. Therefore, taking into account that both methodologies may be responding positively to reducing sugars or other components that are not involved in the antioxidant activity, the yields were verified by techniques such as ABST, DPPH and ORAC, which quantify only antioxidant metabolites.

The ORAC methodology measures the ability of the polyphenols to trap the peroxy radicals generated *in situ* through a hydrogen atom transfer (HAT) proton transfer mechanism; and is the most commonly used technique to measure the total antioxidant potential of foods and nutritional supplements (Wu et al., 2004). It provides a controllable source of peroxy radicals and can be adapted to detect both hydrophilic and hydrophobic antioxidants by handling the source of the radical and the solvent (Prior et al., 2003; Ou et al., 2005). The antioxidant activity measured by the ORAC method for the Nalca leaf yielded a value of 192006.07  $\mu\text{mol Trolox}/100\text{ g DW}$ . These values suggest that the *G. tinctoria* leaf has a high content of hydrophilic antioxidants with the ability to neutralize radicals  $\text{ROO}\cdot$  peroxides. These are superior to those reported for *S. mahagoni* (125520  $\mu\text{mol Trolox}/100\text{ g DW}$ ) (Preciado et al., 2016); green tea and black of Mortiño (*V. meridionale*) (77880 and 149682  $\mu\text{mol Trolox} / 100\text{ g DW}$ ) (Zapata-Vahos et al., 2015); yerba mate (*Ilex laurina*) (58880  $\mu\text{mol Trolox} / 100\text{ g DW}$ ), *Ilex paraguariensis* (35990  $\mu\text{mol}$

$\text{Trolox}/100\text{ g DW}$ ) (Pérez et al., 2014); *Camellia euphlebia*, *Camellia microcarpa* and *Camellia nitidissima* (85640, 59630 and 52490  $\mu\text{mol Trolox}/100\text{ g DW}$  respectively) (Song et al., 2011); freeze-dried leaves of Boldo (*Peumus boldus Molina*) (46160  $\mu\text{mol Trolox}/100\text{ g DW}$ ) (Soto et al., 2014), acai (43795.7  $\mu\text{mol Trolox}/100\text{ g DW}$ ), guayava (11988.2  $\mu\text{mol Trolox}/100\text{ g DW}$ ), cauliflower (10617.7  $\mu\text{mol Trolox}/100\text{ g DW}$ ), and kiwi (4651.1  $\mu\text{mol Trolox}/100\text{ g DW}$ ) (Zapata et al., 2014).

The metabolite content and the antioxidant activity obtained through the ORAC method (together with those quantified by the other methodologies described) are relevant evidence for the study; since they are above those described for other species of recognized pharmaceutical capacity. The antioxidant capacity of *G. tinctoria* leaf measured through ORAC is 147 and 28% higher than the leaf of green tea and black mortiño, a plant species which has shown the ability to reduce the cell viability of SW480 (colon adenocarcinoma cell line) (Zapata-Vahos et al., 2015). A 300% superior to the infusions of *Paemus boldus*, endemic tree of Chile recognized for behaving as an antioxidant, chemopreventive, digestive stimulant, diuretic, relaxing and in the treatment of disorders of liver and gallbladder (Soto et al., 2014; Simirgiotis and Schmeda-Hirschmann, 2010, Fernandez et al., 2009), and, 226 and 433% higher than extracts prepared from leaves of species of the *Ilex family* (which are widely consumed as tea-like beverage in Argentina, Uruguay, southern Brazil, Bolivia, Chilean Patagonia), and which have been reported to be capable of inhibiting the growth of metastatic-derived cells (SW620) and of primary human colon adenocarcinoma cells SW480, HT-29 and CaCo-2 (Perez et al., 2014; González de Mejia et al., 2010).

### Antifungal activity

According to Gramza and Korczak (2005) polyphenols can inhibit the growth of species of the genus *Clostridium* and *Helicobacter pylori*. It has also been shown that certain extracts of antioxidant tea inhibitors inhibit some food-borne pathogenic microorganisms such as *Bacillus cereus* (Almajano et al., 2008; Si et al., 2006), *Staphylococcus aureus* (Kim et al., 2004; Si et al., 2006; Taguri et al., 2004), *Vibrio cholerae* (Taguri et al., 2004), *Listeria monocytogenes* (Kim et al., 2004; Kim and Fung, 2004; Si et al., 2006), *Escherichia coli* (Cho et al., 2007; Kim and Fung, 2004; Si et al., 2006), *Salmonella typhimurium* (Si et al., 2006), *Salmonella enteritidis* (Kim and Fung, 2004), *Micrococcus luteus* y *Pseudomona aeruginosa* (Almajano et al., 2008); and that in particular green tea catechins may decrease the resistance of *S. aureus* to antibiotics such as oxacillin (Stapleton et al., 2004).

It is possible to attribute this antagonistic effect to the high content of metabolites reported in the Nalca leaf and its synergistic effect, which is directly related to its

antioxidant and biological activity. It is precisely the complex mixture of compounds (usually of low molecular weight) that are part of the leaves of the plants (glucosides, saponins, tannins, alkaloids, polyphenols, flavonols, flavonoids, catechins, epigallocatechin, organic acids and volatile components, terpenes, among others) (Tajkarimi et al., 2010) that contribute to the antimicrobial activity.

The high levels of catechin and epicatechin obtained from leaf of *C. sinensis* have shown the ability to inhibit the *in vitro* growth of *Candida albicans* (Anand and Rai, 2017; Ning et al., 2015), the most common human fungal pathogen (Pfaller and Diekema, 2007), associated with superficial infections of the skin, hair, oral or vaginal tract and with life-threatening systemic infections (Luo et al., 2013; Paramythiotou et al., 2014). Also, *in vitro* antibacterial activity was reported by Riso et al. (2002) against *E. coli*, *Pseudomonas aeruginosa* and *S. aureus*.

According to Avello et al. (2012), the possible antifungal mechanisms may be due to the volatility and low molecular weight of most of the compounds mentioned, since they can cross resistance structures of the fungal species. Thus, both cytoplasmic membrane damage, interference in folate metabolism and reduction in ergosterol content, promote the formation of pores through which ions and molecules (mainly  $K^+$  and  $H^+$ ) diffuse, which ultimately compromise viability of the membrane and that according to the concentrations of the extract applied would affect the production of ATP. Also, due to the mixing of a large number of components present, they are considered low risk in the development of resistance by the pathogens, since each of the components has its own biological activity, and it is difficult to correlate the action of a single compound or classes of compounds, because the fungitoxic potential is the result of the synergistic action of the constituents (Burt, 2004; Yahyzadeh et al., 2008).

The *in vitro* antifungal activity of *G. tinctoria* leaf extract was evaluated qualitatively by the presence of inhibition halos. Figure 2 shows the inhibition halos against two fungi, *C. laurentii* (yeast) and *Cladophialophora* (mold).

In this research and supported by the studies conducted by Neri et al. (2006) and Amiri et al. (2008) the fungistatic effect was considered when a reactivation of mycelial growth was observed after seven days of incubation, and effect fungicide when there was no mycelial growth of the pathogen after the period of incubation and observation that corresponded to the period of 20 days. According to Gajanan et al. (2011), the fungicidal effect in the laboratory refers to the agent that decreases the colonies by 99.9% from a pure subculture inoculum. In this way, the pure extract of leaf, produced in laboratory conditions have inhibitory effects (fungicidal effect) on both microorganisms; since the effect remained unchanged until the observation made on day 20. In parallel, there were negative controls (Well 6), which after their incubation showed an abundant growth.

These results are promising because in cheeses conserved (even in refrigeration) species of the genus *Cryptococcus* predominate; which interferes in the maturation of this product by metabolizing the lactic acid, raising the pH, favoring the growth of proteolytic bacteria and generating fructifications, undesirable leuconiform flavors and an unpleasant texture for the appearance of the product (Pereira-Dias et al., 2000; Orberá, 2004). On the other hand, in the food industry, *Cladophialophora* has been isolated in non-integral maize flour, tomatoes, carrots, apples, strawberries and pears (Badali et al., 2008; Romão-Dumaresq et al., 2016). In addition, taking into account the content of metabolites and the antioxidant capacity presented by *G. tinctoria*, it is possible to project its biological activity on various microorganisms and to present itself as an alternative to the antagonistic action exerted by extracts from native and endemic Chilean trees such as Laurel (*Laurelia sempervirens*), Boldo (*Peumus boldus*) and Tepa (*Laureliopsis philippiana*) against pathogenic strains of plant species such as *Rhizoctonia solani*, *Pythium irregulare*, *Ceratocystis pilifera*, *Phragmidium violaceum* and *Fusarium oxysporum* (Bittner et al., 2009).

## Conclusions

In this research, the antioxidant and antifungal potential of *G. tinctoria* was reported, highlighting its added value compared with other plants, fruits and vegetables that are of mass consumption. The results of this research suggest that the *G. tinctoria* leaf is a rich source of polyphenols and secondary metabolites, which intervene in the high antioxidant capacity quantified by this study. According to this it is possible to consider it as (1) a functional food and rich source of antioxidants and bioactive compounds, and (2) as an antifungal and/or therapeutic alternative.

## CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

## ACKNOWLEDGEMENTS

The authors express their gratitude to the entities in charge of the Platform for Student and Academic Mobility of the Pacific Alliance (VIII Call) -ICETEX, APC (Presidential Agency of International Cooperation of Colombia), Colombian Ministry of Foreign Affairs- and to the Food Science Laboratory from the National University of Colombia, headquarters in Medellín.

## REFERENCES

Aguilar-Hernández I, Kristian N, López-Luke T, Contreras-Torres FF, Petter J, Ornelas-Soto N (2017). Vibrational Spectroscopy Surface

- enhanced Raman spectroscopy of phenolic antioxidants: A systematic evaluation of ferulic acid, p-coumaric acid, caffeic acid and sinapic acid. *Vib. Spectrosc.* 89:113-122.
- Anand J, Rai N (2017). Anticandidal synergistic activity of green tea catechins, antimycotics and copper sulphate as a mean of combinational drug therapy against candidiasis. *J. Med. Mycol.* 27(1):33-45.
- Anhê FF, Desjardins Y, Pilon G, Dudonné S, Genovese MI, Lajolo FM, Marette A (2013). Polyphenols and type 2 diabetes: A prospective review. *Pharm. Nutr.* 1(4):105-114.
- Almajano MP, Carbo R, Jimenez AP, Gordon MH (2008). Antioxidant and antimicrobial activities of tea infusions. *Food Chem.* 108:55-63.
- Amiri A, Dugas R, Pichot AL, Bompeix G (2008). In vitro and in vivo activity of eugenol oil (*Eugenia caryophyllata*) against four important postharvest apple pathogens. *Int. J. Food Microbiol.* 15:126(1-2):13-9.
- Association of Official Analytical Chemists (AOAC) (1980). Official Methods of Analysis of AOAC International, 13th Ed., AOAC International, Gaithersburg, MD, USA.
- Association of Official Analytical Chemists (AOAC) (1996). Official Methods of Analysis of AOAC International, 16th Ed., AOAC International, Gaithersburg, MD, USA.
- Avello M, López C, Gatica C, Bustos E, Brieva A, Pastene E (2012). Efectos antimicrobianos de extractos de plantas chilenas de las familias Lauraceae y Atherospermataceae. *Rev. Cubana Plant Med.* 17(1):73-83.
- Badali H, Gueidan C, Najafzadeh MJ, Bonifaz A, Van den Ende AHGG, de Hoog GS (2008). Biodiversity of the genus *Cladophialophora*. *Stud. Mycol.* 61:175-191.
- Bateman JV (1970). Nutrición Animal: Manual de Métodos Analíticos. Centro Regional de Ayuda Técnica, México D.F. 468 p.
- Bauer AW, Kirby WM, Sherris JC, Turc M (1996). Antibiotic susceptibility testing by standardized single method. *Am. J. Clin. Pathol.* 45:493-496.
- Bello J (2000). Ciencia Bromatológica. Principios generales de los alimentos. Ediciones Díaz de Santos, S.A. Madrid. España.
- Benzie IF, Strain JJ (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Anal. Biochem.* 239(1):70-76.
- Bergman B, Johansson C, Soderback E (1992). The Nostoc-Gunnera symbiosis. *New Phytol.* 122:379-400.
- Berlitz HD, Grosch W (1999). Tablas de Composición de Alimentos. Editorial Acribia. Zaragoza. España.
- Bittner M, Aguilera MA, Hernández V, Arbert C, Becerra J, Casanueva ME (2009). Fungistatic Activity Of Essential Oils Extracted from *Peumus boldus* Mol., *Laureliopsis philippiana* (Looser) Schodde and *Laurelia sempervirens* (Ruiz & Pav.) Tul. (*Chilean Monimiaceae*). *Chilean J. Agric. Res.* 69(1):30-37.
- Brand-Williams W, Cuvelier ME, Berset C (1995). Use of a free radical method to evaluate antioxidant activity. *Food Sci. Technol.* 28:25-30.
- Burt S (2004). Essential oils: their antibacterial properties and potential applications in foods – a review. *Int. J. Food Microbiol.* 94:223-253.
- Calil Brondani J, Reginato FZ, da Silva Brum E, de Souza Vencato M, Lima Lhamas C, Viana C, Manfron MP (2017). Evaluation of acute and subacute toxicity of hydroethanolic extract of *Dolichandra unguis-cati* L. leaves in rats. *J. Ethnopharmacol.* 202(3):147-153.
- Campos AM, Lissi EA (1997). Kinetics of the reaction between 2,2-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) derived radical cations and phenols. *Int. J. Chem. Kinet.* 29:219-224.
- Chaves N, Escudero J (1999). Variation of flavonoid synthesis induced by ecological factors. In: Inderjit, KMM Dakhini, CL Foy (eds): Principles and Practices in Plant Ecology Allelochemical Interactions. CRC Press, Boca Raton, Florida, USA.
- Chena G, Wang F, Trachoothama D, Huang P (2010). Preferential killing of cancer cells with mitochondrial dysfunction by natural compounds. *Mitochondrion* 10:614-25.
- Cho YS, Schiller NL, Kahng HY, Oh KH (2007). Cellular responses and proteomic analysis of *Escherichia coli* exposed to green tea polyphenols. *Curr. Microbiol.* 55:501-506.
- Clifford MN (2000). Chlorogenic acids and other cinnamates, occurrence, dietary burden, absorption and metabolism. *J. Sci. Food Agric.* 80:1033-1043.
- Elfalleh W, Hannachi H, Tlili N, Yahia Y, Nasri N, Ferchichi A (2012). Total phenolic contents and antioxidant activities of pomegranate peel, seed, leaf and flower. *J. Med. Plants Res.* 6(32):4724-4730.
- EPPO (2014). PQR-EPPO database. Paris, France: European and Mediterranean Plant Protection Organization. Available at: <http://www.eppo.int/DATABASES/pqr/pqr.html>
- Ferguson LR, Zhu S, Harris PJ (2005). Antioxidant and antigenotoxic effects of plant cell wall hydroxycinnamic acids in cultured HT-29 cells. *Mol. Nutr. Food Res.* 49:585-593.
- Fernandez J, Lagos P, Rivera P, Zamorano-Ponce E (2009). Effect of boldo (*Peumus boldus* Molina) infusion on lipoperoxidation induced by cisplatin in mice liver. *Phytother. Res.* 23(7):1024-1047.
- Fischer AP, Miles SL (2017). Ascorbic acid, but not dehydroascorbic acid increases intracellular vitamin C content to decrease Hypoxia Inducible Factor -1 alpha activity and reduce malignant potential in human melanoma. *Biomed. Pharmacother.* 86:502-513.
- Fujiki H, Suganuma M, Imai K, Nakachi K (2002). Green tea: cancer preventive beverage and/or drug. *Cancer Lett.* 188:9-13.
- Gajanan B, Zore D, Sitaram S, Mohan K (2011). Terpenoids inhibit *Candida albicans* growth by affecting membrane integrity and arrest of cell cycle. Sponsored School of Life Sciences, SRTM University, Nanded 431606, MS, India. *Phytomedicine* 18:1181-1190.
- González de Mejía E, Song YS, Heck CI, Ramírez-Mares MV (2010). Yerba mate tea (*Ilex paraguariensis*): Phenolics, antioxidant capacity and in vitro inhibition of colon cancer cell proliferation. *J. Funct Foods* 2:23-34.
- Gramza A, Korczak J (2005). Tea constituents (*Camellia sinensis* L.) as antioxidants in lipid systems. *Trends Food Sci. Technol.* 16(8):351-358.
- Gupta S, Ahmad N, Nieminem AL, Mukhtar H (2000). Growth inhibition, cell cycle dysregulation and induction of apoptosis by green tea constituents (-) epigallocatechin-3-gallate in androgen sensitive and androgen in sensitive human prostate carcinoma cells. *Toxicol Pharmacol.* 164:82-90.
- Hamilton-Miller JM (1995). Antimicrobial properties of tea (*Camellia sinensis* L.). *Antimicrob. Agents Chemother.* 39:2375-2377.
- Han X, Shen T, Lou H (2007). Dietary polyphenols and their biological significance. *Int. J. Mol. Sci.* 8:950-988.
- Heim KE, Tagliaferro AR, Bobilya DJ (2002). Flavonoid antioxidants: chemistry, metabolism and structure-activity relationship. *J. Nutr. Biochem.* 13:572-584.
- Hickey B, Osborne B (1998). Effect of *Gunnera tinctoria* (Molina) Mirbel on semi-natural grassland habitats in the west of Ireland. In: Starfinger U, Edwards K, Kowarik I, Williamson M (eds.), Plant Invasions: Ecological Mechanisms and Human Responses. Backhuys Publishers, Leiden, the Netherlands. pp. 195-208.
- Ibrahim MR, El-Banna HM, Omara II, Suliman MA (2011). Evaluation of nutritive value of some citrus pulp as feedstuffs in rabbits diets. *Pak. J. Nutr.* 10(7):667-674.
- International Plant Names Index (IPNI) (2012). The International Plant Names Index. Published on the Internet <http://www.ipni.org>
- Jing LJ, Mohamed M, Rahmat A, Bakar MFA (2010). Phytochemicals, antioxidant properties and anticancer investigations of the different parts of several ginger species (*Boesenbergia rotunda*, *Boesenbergia pulchella* var *attenuata* and *Boesenbergia armeniaca*). *J. Med. Plants Res.* 4(1):27-32.
- Kelebek H, Selli S, Canbas A, Cabaroglu T (2009). HPLC determination of organic acids, sugars, phenolic compositions and antioxidant capacity of orange juice and orange wine made from a Turkish cv. Kosan. *Microchem. J.* 91(2):187-92.
- Kikugawa K, Hakamada T, Hasunuma M, Kurechi T (1983). Reaction of p-hydroxycinnamic acid derivatives with nitrite and its relevance to nitrosamine formation. *J. Agric. Food Chem.* 1(4):780-785.
- Kim S, Fung DYC (2004). Antibacterial effect of crude water-soluble arrowroot (*Puerariae radix*) tea extracts on food-borne pathogens in liquid medium. *Lett. Appl. Microbiol.* 39(4):319-325.
- Kim S, Ruengwilysup C, Fung D (2004). Antibacterial effect of water-soluble tea extracts on foodborne pathogens in laboratory medium and in a food model. *J. Food Protect.* 67:2608-2612.
- Kopjar M, Tadic M, Pilizota V (2015). Phenol content and antioxidant activity of green, yellow and black tea leaves. *Chem. Biol. Technol. Agric.* 2(1):1-6.

- Lazos E (1991). Composition and oil characteristics of apricot, peach and cherry kernel. *Grasas Aceites* 42(2):127-131.
- Lu JM, Lin PH, Yao Q, Chen C (2010). Chemical and molecular mechanisms of antioxidants: Experimental approaches and model systems. *J. Cell Mol. Med.* 14(4):840-860.
- Luo S, Skerka C, Kurzai O, Zipfel PF (2013). Complement and innate immune evasion strategies of the human pathogenic fungus *Candida albicans*. *Mol. Immunol.* 56(3):161-169.
- Mahmood T, Anwar F, Bhatti IA, Iqbal T (2013). Effect of maturity on proximate composition, phenolics and antioxidant attributes of cherry fruit. *Pak. J. Bot.* 45(3):909-914.
- Marszałek K, Woźniak Ł, Kruszewski B, Skąpska S (2017). The Effect of High Pressure Techniques on the Stability of Anthocyanins in Fruit and Vegetables. *Int. J. Mol. Sci.* 18(2):277.
- Montes M, Wilkomirsky T (1985). *Medicina Tradicional Chilena, 1a Edición*, Editorial Universidad de Concepción, Concepción, Chile. P 98.
- Moon J, Shibamoto T (2009). Antioxidant assays for plant and food components. *J. Agric. Food Chem.* 57(5):1655-1666.
- Moore RJ, Jackson KG, Minihane AM (2009). Green tea (*Camellia sinensis*) catechins and vascular function. *Br. J. Nutr.* 102:1790-1802.
- Morales M, Zapata S, Jaime T, Rosales S, Alzate F, Maldonado M, Zamorano P, Rojano B (2017). Mangiferin content, carotenoids, tannins and oxygen radical absorbance capacity (ORAC) values of six mango (*Mangifera indica*) cultivars from the Colombian Caribbean. *J. Med. Plants Res.* 11(7):144-152.
- Moreiras O, Carbajal A, Cabrera L, Cuadrado C (2006). *Tablas de Composición de Alimentos*. Ediciones Piramide. Madrid. España.
- Morillas-Ruiz JM, Delgado-Alarcón JM (2012). Nutritional analysis of vegetable food with different origins: Evaluation of antioxidant capacity and phenolic total compounds. *Nutr. Clin. Diet. Hosp.* 32(2):8-20.
- Najmus AA, Whitney PJ (2011). Differential behaviour of the dinitrosalicylic acid (DNS) reagent towards mono and di-saccharide sugars. *Biomass Bioenergy* 35(11):4748-4750.
- Neri F, Mari M, Brigati S (2006). Control of *Penicillium expansum* by plant volatile compounds. *Plant Pathol.* 55(1):100-105.
- Ning Y, Ling J, Wu CD (2015). Synergistic effects of tea catechin epigallocatechin gallate and antimicrobials against oral *Candida* species. *Arch. Oral Biol.* 60(10):1565-1570.
- Oliveiro T, Capuano E, Cammerer B, Fogliano V (2009). Influence of roasting on the antioxidant activity and HMF formation of a cocoa bean model systems. *J. Agric. Food Chem.* 57(1):147-52.
- Orberá T (2004). Métodos moleculares de identificación de levaduras de interés biotecnológico. *Rev. Iberoamericana Micol.* 21:15-19.
- Osman AM, Wong KKY, Fernyhough A (2006). ABTS radical-driven oxidation of polyphenols: Isolation and structural elucidation of covalent adducts. *Biochem. Biophys. Res. Commun.* 346(1):321-329.
- Ou B, Prior RL, Huang D (2005). The chemistry behind dietary antioxidant capacity assays. *J. Agric. Food Chem.* 53:1841-1856.
- Ozdal T, Sela DA, Xiao J, Boyacioglu D, Chen F, Capanoglu E (2016). The reciprocal interactions between polyphenols and gut microbiota and effects on bioaccessibility. *Nutrients* 8(2):1-36.
- Paramythiotou E, Frantzeskaki F, Flevari A, Armaganidis A, Dimopoulos G (2014). Invasive fungal infections in the ICU: how to approach, how to treat. *Molecules* 19(1):1085-1119.
- Pereira-Dias S, Potes ME, Marinho A, Malfeito-Ferreira M, Loureiro V (2000). Characterisation of yeast flora isolated from an artisanal Portuguese ewes' cheese. *Int. J. Food Microbiol.* 60(1):55-63.
- Pérez JM, Maldonado ME, Rojano BA, Alzate F, Sáez J, Cardona W (2014). Comparative Antioxidant, Antiproliferative and Apoptotic Effects of *Ilex laurina* and *Ilex paraguariensis* on Colon Cancer Cells. *Trop. J. Pharm. Res.* 13(8):1279-1286.
- Petzold G, Catril G, Duarte Carlos (2006). Caracterización fisicoquímica de peciolos de pague (Gunnera tinctoria). *Rev. Chilena Nutr.* 33(3):539-543.
- Pfaller MA, Diekema DJ (2007). Epidemiology of invasive candidiasis: a persistent problem. *Clin. Microbiol. Rev.* 20:133-63.
- Pounis G, Costanzo S, Di Giuseppe R, De Lucia F, Santimone I, Sciarretta A, Barisciano P, Persichillo M, De Curtis A, Zito F, Di Castelnuovo AF, Sieri S, Benedetta Donati M, De Gaetano G, Iacoviello L (2013). Consumption of healthy foods at different content of antioxidant vitamins and phytochemicals and metabolic risk factors for cardiovascular disease in men and women of the Moli-sani study. *Euro. J. Clin. Nutr.* 67(2):207-13.
- Pulido R, Bravo L, Saura-Calixto F (2000). Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/ antioxidant power assay. *J. Agric. Food Chem.* 48:3396-3402.
- Preciado LM, Lobo-Echeverri T, Pereañez JA, Zapata K, Rojano BA (2016). Antioxidants from three *Swietenia* Species (Meliaceae). *J. Med. Plants Res.* 10(2):8-17.
- Prior RL, Hoang H, Gu L, Wu X, Bacchiocca M, Howard L, Hampsch-Woodill M, Huang D, Ou B, Jacob R (2003). Assays for hydrophilic and lipophilic antioxidant capacity (oxygen radical absorbance capacity (ORACFL) of plasma and other biological and food samples. *J. Agric. Food Chem.* 51:3273-3279.
- Prior RL, Wu X, Schaich K (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J. Agric. Food Chem.* 53(10):4290-4302.
- Quiles J, Ramirez M, Yaqoob P (2006). Olive oil and health. CABI Head Office International. Oxfordshire, UK.
- Ramakrishna N, Ravishankar GA (2011). Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signal. Behav.* 6(11):1720-1731.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic. Biol. Med.* 26(9-10):1231-1237.
- Riso P, Erba D, Criscuoli F, Testolin G (2002). Effect of green tea extract on DNA repair and oxidative damage due to H<sub>2</sub>O<sub>2</sub> in jurkat T cells. *Nutr. Res.* 22:1143-1150.
- Roginsky V, Lissi EA (2005). Review of methods to determine chain-breaking antioxidant activity in food. *Food Chem.* 92:235-254
- Romão-Dumaresq AS, Dourado MN, Fávoro LC, Mendes R, Ferreira A, Araújo WL (2016). Diversity of Cultivated Fungi Associated with Conventional and Transgenic Sugarcane and the Interaction between Endophytic *Trichoderma virens* and the Host Plant. *PLoS ONE* 11(7):e0158974.
- Si W, Gong J, Tsao R, Kalab M, Yang R, Yin Y (2006). Bioassay-guided purification and identification of antimicrobial components in chinese green tea extract. *J. Chromatogr. A.* 1125:204-210.
- Simirgiotis MJ, Schmeda-Hirschmann G (2010). Direct identification of phenolic constituents in Boldo folium (*Peumus boldus Mol.*) infusions by high-performance liquid chromatography with diode array detection and electrospray ionization tandem mass spectrometry. *J. Chromatography A.* 1217:443-449.
- Singleton VL, Rossi JA (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Viticult.* 16:144-158.
- Song L, Wang X, Zheng X, Huang D (2011). Polyphenolic antioxidant profiles of yellow camellia. *Food Chem.* 129:351-357.
- Soto C, Caballero E, Pérez E, Zúñiga ME (2014). Effect of extraction conditions on total phenolic content and antioxidant capacity of pretreated wild *Peumus boldus* leaves from Chile. *Food Bioprod. Process* 92(3):328-333.
- Stapleton P, Shah S, Hamilton-Miller J, Hara Y, Nagaoka Y, Kumagai A, Uesato S, Taylor P (2004). Anti-*Staphylococcus aureus* activity and oxacillin resistance modulating capacity of 3-O-acyl-catechins. *Int. J. Antimicrob. Agents* 24:374-380.
- Taguri T, Tanaka T, Kouno I (2004). Antimicrobial activity of 10 different plant polyphenols against bacteria causing food-borne disease. *Biol. Pharm. Bull.* 27(12):1965-1969.
- Tajkarimi M, Ibrahim S, Cliver D (2010). Antimicrobial herb and spice compounds in food. *Food Control.* 21(19):1199-1218.
- The Plant List (2010). Version 1. Published on the Internet; <http://www.theplantlist.org>
- Torres B, Tiwari BK, Partas A, Cullen PJ, Brunton N, O'Donnel CP (2011). Stability of anthocyanins and ascorbic acid of high pressure processes blood orange juice during storage. *Innov. Food Sci. Emerg.* 12:93-97.
- Troncoso P, Pérez C, Larrain J, Ardiles V (2013). Desarrollo de la fijación simbiótica de nitrógeno en una cronosecuencia primaria en la Isla Santa Inés, Región de Magallanes, Chile. *Rev. Chilena Historia Nat.* 86(3):345-356.
- Valdebenito G, Campos J, Larrain O, Aguilera M, Kahler C, Ferrando M,

- García E, Sotomayor A (2003). Boletín Divulgativo N° 12 *Gunnera chilensis* Lam. Innovación Tecnológica y Comercial de Productos Forestales No Madereros (PFNM) en Chile. Proyecto Fondef-Infor-Fundación Chile.
- Wanntorp L, Klackenborg J (2006). *Gunnera morae* (Gunneraceae). A new species from Colombia. *Caldasia* 28(2):221-225.
- Williams PA, Ogle CC, Timmins SM, La Cock GD, Clarkson J (2005). Chilean rhubarb (*Gunnera tinctoria*): biology, ecology and conservation impacts in New Zealand. DOC Research & Development Series 210. Department of Conservation, Wellington. P 27.
- Wojdylo A, Osmianski J, Czermerys R (2007). Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chem.* 105(3):940-949.
- Wu X, Beecher GR, Holden JM, Haytowitz DB, Gebhardt SE, Prior RL (2004). Lipophilic and hydrophilic antioxidant capacities of common foods in the United States. *J. Agric. Food Chem.* 52(12):4026-4037.
- Yahyzadeh M, Omidbaigi R, Zare R, Taheri H (2008). Effect of some essential oils on mycelial growth of *Penicillium digitatum*. *Sacc. World J. Microbiol. Biotechnol.* 24:1445-2145.
- Yang CS, Maliakal P, Meng X (2002). Inhibition of carcinogenesis by tea. *Annu. Rev. Pharmacol. Toxicol.* 42:25-54.
- Zamora-Ros R, Fedirko V, Trichopoulou A, González CA, Bamia C, Trepo E, Nöthlings U, Duarte-Salles T, Serafini M, Bredsdorff L, Overvad K, Tjønneland A, Halkjaer J, Fagherazzi G, Perquier F, Boutron Ruault MC, Katzke V, Lukanova A, Floegel A, Boeing H, Lagiou P, Trichopoulos D, Saieva C, Agnoli C, Mattiello A, Tumino R, Sacerdote C, Bueno-de-Mesquita HB, Peeters PH, Weiderpass E, Engeset D, Skeie G, Arguëlles MV, Molina-Montes E, Dorronsoro M, Tormo MJ, Ardanaz E, Ericson U, Sonestedt E, Sund M, Landberg R, Khaw KT, Wareham NJ, Crowe FL, Riboli E, Jenab M (2013). Dietary flavonoid, lignan and antioxidant capacity and risk of hepatocellular carcinoma in the European prospective investigation into cancer and nutrition study. *Int. J. Cancer* 133:2429-2443.
- Zapata S, Piedrahita AM, Rojano B (2014). Capacidad atrapadora de radicales oxígenos (ORAC) y fenoles totales de frutas y hortalizas de Colombia. *Perspect. Nutr. Hum.* 16:25-36.
- Zapata-Vahos IC, Villacorta V, Maldonado ME, Castro-Restrepo D, Rojano B (2015). Antioxidant and cytotoxic activity of black and green tea from *Vaccinium meridionale* Swartz leaves. *J. Med. Plants Res.* 9(13):445-453.
- Zobel AM, Clarke PA, Lynch JM (1999). Production of phenolics in response to UV irradiation and heavy metals in seedlings of *Acer* species. Narwal, S.S., ed. *Recent advances in allelopathy*, Boca Raton, CRC Press. pp. 1-12.
- Zoratti L, Karppinen K, Luengo Escobar A, Häggman H, Jaakola L (2014). Light-controlled flavonoid biosynthesis in fruits. *Frontiers Plant Sci.* 5:534.