

## Full Length Research Paper

# Oxidative stabilization of palm olein with isoespintanol (2-isopropyl-3,6-dimethoxy-5-methylphenol) isolated from *Oxandra cf xylopioides*

Karol Zapata<sup>1</sup>, Juan P. Arias<sup>1</sup>, Farid B. Cortés<sup>2</sup>, Camilo L. Alarcon<sup>3</sup>, Diego L. Durango Restrepo<sup>1</sup> and Benjamín A. Rojano<sup>1\*</sup>

<sup>1</sup>Research Group Chemistry of Natural Products and Food, Universidad Nacional de Colombia, Medellin, Colombia.

<sup>2</sup>Research Group Surface Phenomena-Michael Polanyi- Universidad Nacional de Colombia, Medellin, Colombia.

<sup>3</sup>Department of Pharmacy, Faculty of Chemistry – Pontificia Universidad Católica de Chile, Santiago de Chile, Chile.

Received 5 October, 2016; Accepted 3 March, 2017

**Antioxidants are used to retard oxidative processes and improve the sensory and nutraceutical quality of cooking oils. Isoespintanol, a monophenol isolated from *Oxandra cf. xylopioides*, has shown a greater free radical scavenger ability than thymol (biosynthetic analog). Isoespintanol antioxidant capacity was evaluated and compared to butylated hydroxytoluene (BHT). Cytotoxicity, thermal stability and effect on olein stability studies were performed. The results revealed that Olein oxidation time was reduced 17.2 and 4.2% in the presence of Isoespintanol and BHT, respectively. Thermogravimetric curves indicated higher thermal resistance for Isoespintanol than for BHT. Finally, cytotoxicity tests for Isoespintanol against murine macrophages revealed no effect on cell viability, indicating their possible use as a safe food additive.**

**Key words:** Oxidative stabilization, isoespintanol, bht, antioxidant capacity, cytotoxicity, safe food additive.

## INTRODUCTION

Frying is a wet method of cooking where food is subjected to a quick immersion in a bath of fat or oil at temperatures between 150 and 180°C. The resulting food remains dry, crispy, browned and has the best culinary characteristics. However, frying oils are subjected to elevated temperatures for extended periods of time that leads to oxidative reactions. The reaction of dissolved oxygen and water produces compounds that confer

unpleasant flavors to oils and food. Common compounds include volatile carbonyls, hydroxyl acids, keto acids and epoxy acids, providing unpleasant flavors to oils and food (Alvis and Velez, 2008; Wu et al., 2013, Paz et al., 2015).

Oil used for frying should possess chemical characteristics that contribute to its useful life. For instance, saturated and unsaturated fatty acid content ratio affects the stability of the product during the heating

\*Corresponding author. E-mail: brojano@unal.edu.co. Tel: +57 4 4309381.

process. Different studies suggest that oils with high content of monounsaturated fatty acids improve stability of fried food. Oleic acid from palm, olives and sunflower is an example of such oils (Casal et al., 2010; Aladedunye and Przybylski, 2013).

According to the Department of Agriculture of United States of America, palm oil is the second type of oil with higher production volume in the world (USDA, 2004). Palm olein was introduced to the market in the late twentieth century and it has remained in use because of its technical and economic advantages compared to other oils and fats. It is estimated that millions of tons of palm olein and oil in their natural forms or mixtures with other oils, are used annually for domestic and industrial frying. In addition, numerous studies demonstrate the benefits of the use of palm oil in frying because of its greater oxidative stability compared to other oils (Ismail, 2005; Kalogianni et al., 2011; Mozzon et al., 2013). However, palm olein products are susceptible to lipid peroxidation due to the presence of unsaturated fatty acids. For this reason, antioxidants are used to retard oxidative processes and improve the sensory and nutraceutical quality of the products (Sanchez, 2011, Rojano et al., 2008). Moreover, addition of natural metabolites provides extra protection against oxidative stress enhancing the antioxidant system of the consumer. Although antioxidants in food determine the lifetime and reduce the nutritional losses, they never regenerate the quality of a highly-oxidized food.

Butylhydroxyanisole (BHA) and butylatedhydroxytoluene (BHT) are synthetic antioxidants widely used in the food industry; however, they can cause adverse health effects, such as increased cholesterol, liver enlargement and induction of liver cancer. Therefore, the search for natural alternatives with high antioxidant activity and without cytotoxic and genotoxic effects have become a necessity for the food industry (Niaiforovia et al., 2010; Krishnaiah et al., 2011). Recent years have witnessed a growing interest for natural antioxidants. For instance, a study done by Piedrahita et al. (2015) showed that rosemary extracts have shown higher thermal resistance and antioxidant capacity than synthetic antioxidants (BHA, BHT and TBHQ) when working with potato frying processes in olein palm; additionally, some studies have described the stabilizing capacity of rosemary extracts on oily matrices (Upadhyay et al., 2006); Upadhyay and Mishra (2015a) show that the oxidative stability in sunflower oil given by rosemary oleoresin (1200 and 1500 mg/kg) and tertiary butyl hydroquinone (200 mg/kg) are comparable.

Isoespintanol is a isopropylphenol extracted with low polarity solvents from leaves of *Oxandra cf. xylopioides* (*Annonaceae*) a native plant from Colombia and it has opened new alternatives in the world of fats and oils, due to its chemical structure that makes it soluble in liposoluble media; some studies have shown the antioxidant capacity of Isoespintanol in oily matrices

(Rojano et al., 2008a). Likewise, Isoespintanol (2-isopropyl-3,6-dimethoxy-5-methylphenol) have shown a greater free radical scavenging ability than its biosynthetic analog (thymol) (Rojano et al., 2008b).

The industry of frying processes requires antioxidants with high thermal stability that preserve the sensory quality and oxidative stability of oils subjected to high temperatures. In the present study, the authors assess the antioxidant capacity of Isoespintanol to trap reactive oxygen species compared to BHT. We compare the effect of adding Isoespintanol or BHT on the oxidative stability of palm olein under accelerated oxidative conditions. Finally, we calculate the *in vitro* thermal stability of Isoespintanol and BHT and estimate the cytotoxicity of Isoespintanol in biological systems.

## MATERIALS AND METHODS

### Reagents

Fluorescein, dichlorofluoresceindiacetate, pyrogallol red, AAPH and BHT were bought from Sigma Aldrich through Colombia G & G Successors SAS. Palm olein (antioxidant free) was provided by a local supplier (TECNAS).

### Plant material

The *Oxandra cf. xylopioides* species were collected in the municipality of Montería (Córdoba, Colombia) in coordinates 08°48'17" north latitude and 75°42'07" west longitude; in October at warm weather under 24°C. The material was identified by the botanist Francisco Javier Roldan Palacios. A sample under file number 037852 was deposited in the Botanical Garden Joaquín Antonio Uribe, Medellín (Colombia). The plant material was cleaned and stored at 4°C until further analysis.

### Extraction and isolation

The powdered leaves (1.0 kg) of *Oxandra cf. xylopioides* were extracted with petroleum ether (15 L) by percolation at room temperature and concentrated in vacuum to obtain a crude extract (150 g). The petroleum ether extract (30 g) was subjected to passage over a silica gel column (5 x 80 cm) eluting with a step gradient of hexane: dichloromethane (95:5, 80:20, 70:30, 0:100 each 1.0 L), to obtain four fractions. Fraction II was dissolved in dichloromethane and compound 4 recrystallized in hexane (3.0 g) (1.5%). The compound was characterized by NMR (1H and 13C COSY bi-dimensional and, NOESY, HMQC, HMBC), MS, UV and IR, and their physical properties; these results were previously reported (Rojano et al., 2007).

### Fatty acid composition of palm olein

The fatty acid profile of the oil was performed on an Agilent 6890N GC coupled to an Agilent 5973N MS selective detector equipped with a split/split-less injector. The injector temperature was 300°C and the sample previously derivatized with KOH-methanol. The sample was automatically injected into the split-less mode. A column HP-5ms (5% phenyl methyl siloxane) of 30 m, 0.25 mm with a film thickness of 0.25 µm and at a maximum temperature of

325°C was used. The temperature program started at 50°C and went up to 200°C in 5 min, then it reached a final temperature of 300°C (14 min) at a rate of 10°C/min. Helium was used as carrier at a constant flow of 1 mL/min gas. The detector temperature was 300°C. The software used to calculate all parameters was MSD ChemStation D 02.00.275 Copyright © Agilent Technologies 1989-2005. For determination of saturated and unsaturated fatty acids as methyl esters the database NIST 2005 was used (Zapata et al., 2013).

### Oxidative stability of palm olein

The active oxygen method (AOM) was used to monitor the oxidative condition of oil. 30 mL of sample were placed in a heater at a constant temperature of  $99 \pm 0.1^\circ\text{C}$  and aerated at 1.150 mL air/min until the time in which a peroxide value (PV) of 100 meq. O<sub>2</sub>/kg. oil was reached (Negative Control). Other samples were assessed under the same conditions, one in presence of Isoespiptanol and the other in presence of BHT. Both antioxidants were at a concentration of 200 mg/L, and were named as IE200 and BHT200, respectively. The concentrations were defined as the maximum concentration permitted for this application in different countries.

### Peroxide value (PV)

The peroxide value (meq.O<sub>2</sub>/kg.oil) was determined by spectrophotometry. This method is based on the oxidation of Fe<sup>+2</sup> to Fe<sup>+3</sup> by hydroperoxides to form a complex Fe<sup>+3</sup>-thiocyanate. 0.01 g of sample were added to 3 mL of a solution chloroform: methanol (7:3), the mixture was stirred for 10 s and 1 mL of this solution was mixed with 50 µL of FeSO<sub>4</sub>-BaCl<sub>2</sub> (0.144 M - 0.4 M) and 50 µL of NH<sub>4</sub>SCN (0.44 M). The mixture was incubated for 20 min in the darkness; the absorbance was determined at 510 nm in a spectrophotometer Jenway® 6405 UV/Vis (Zapata et al., 2015).

### Thermal stability of antioxidants

The thermal stability of the antioxidants was estimated using a thermogravimetric analyzer (Q50, TA Instruments Inc., New Castle, DE). Analyses were carried out by heating the samples in air from 30 to 300°C at a rate of 20°C.min<sup>-1</sup>. Airflow rate was constant throughout the experiment (100 cm<sup>3</sup>/min). Initially, the instrument was calibrated for mass changes and temperature, using nickel as a reference material. For the export of graphical and numerical data, Q Series software was used (Q50-1459 TGA Q-50 @ Mfg-TGA).

### Antioxidant capacity - tests ORAC (Oxygen Radical Absorbance Capacity)

The following techniques were used to determine the antioxidant activity of Isoespiptanol compared with BHT: ORAC Fluorescein Hydrophilic (H-ORAC<sub>FL</sub>), ORAC Fluorescein Lipophilic (L-ORAC<sub>FL</sub>), ORAC dichlorofluoresceindiacetate (ORAC<sub>DCFH</sub>) and ORAC pyrogallol red (ORAC<sub>RPG</sub>); the ORAC methodologies differ in the target molecule.

#### H-ORAC<sub>FL</sub>

The method was described in the beginning by Prior et al. (2005) and is currently carried out with modifications described as follows: A volume of 30 µL of the sample was added to 2,899 µL of PBS (75 mM), 21 µL fluorescein  $1 \times 10^{-2}$  M in PBS (75 mM) and 50 µL of

AAPH 0.6 M in PBS (75 mM). Temperature and pH were controlled at 37°C and 7.4 respectively. Readings were performed at  $\lambda$  excitation of 493 nm and 10 slit;  $\lambda$  emission of 515 nm and emission slit 15, runs with attenuator 1%. The protective effect of the antioxidant is calculated using the differences in areas under the curve of decay of fluorescein in the absence and presence of sample, and compared against the primary standard Trolox. Results were expressed as µmol Trolox equivalents/100 grams of sample according to the following equation.

$$\text{ORAC} = \frac{(\text{AUC} - \text{AUC}^0)}{(\text{AUC}_{\text{Trolox}} - \text{AUC}^0)} f[\text{Trolox}] \quad (1)$$

Where AUC is the area under the curve, in the presence of the sample AUC<sup>0</sup> is the area under the curve for the control (no sample), AUC Trolox is the area under the curve using Trolox, f is the dilution factor of the sample assessed (Prior et al., 2005).

#### L-ORAC<sub>F</sub>

Assay is carried out as explained in the previous section. However, the sample preparation is performed according to the protocol reported by Rautenbach and Venter (2010) described as follows: 0.5 g of sample were dissolved in 4 mL of hexane (lipophilic solution), this solution is mixed with 10 mL of a solution of water: acetone (1:1, v / v) with 7%  $\beta$ -cyclodextrin (micellar agent).

A modified ORAC<sub>DCFH</sub> method described by Martin-Romero et al. (2008), was used and described as follows. The nonfluorescent probe 2,7 dichlorodihydrofluorescein (DCFH) reacts with reactive oxygen species (ERO's) generated by the azo compound APPH (2,2'-Azobis(2-amidinopropane) dihydrochloride) in an aqueous medium to form 2,7 dichlorofluorescein (DCF), a fluorescent molecule. The compound tested captures free radicals and reduces the fluorescence emitted by decreasing DCF formed. To carry out the reaction; 50 µL of a solution of AAPH 0.3 M were mixed with 50 µL of 2,7-dichlorofluorescein diacetate 2.4 mM, 2850 µL of phosphate buffer - 75 mM pH 7.4 and 50 µL of the sample at 37°C. The intensity of the fluorescence emitted during the first 10 minutes was read ( $\lambda$  emission 490 nm and  $\lambda$  excitation 530 with 10 nm slit emission/excitation). The protective effect of the antioxidant is calculated using the increase in fluorescence in the presence and absence of the compound assessed, and compared with the calibration curve of the primary standard Trolox. Results were expressed as µmol Trolox equivalents/100 g of sample.

The ORAC<sub>RPG</sub> method described by Lopez-Alarcon and Lissi (2006) was used with the following modifications: Pyrogallol Red (PGR) reacts with reactive oxygen species (ERO's) generated by the azo compound APPH (2,2'-Azobis(2-amidinopropane) dihydrochloride) in an aqueous medium and forms a colorless oxidized product. The compound tested traps free radicals and prevents the reduction of the absorbance due to decreased oxidized RPG formed. To perform the reaction, 10 µL of an aqueous solution of AAPH 0.3 M were mixed with 50 µL of an aqueous solution of Pyrogallol Red (PGR) ( $5 \times 10^{-5}$  M), 30 µL of sample and 210 µL buffer phosphate (75 mM, pH 7.4), the mixture was incubated for 4000 s and the absorbance was monitored during this time at 540 nm. For the spectrophotometric readings, a Thermo Scientific Multiskan Spectrophotometer UV-VIS Spectrum was used. The protective effect of the antioxidant is calculated using the percentage of inhibition of decay in absorbance when the reaction contains the sample; this value is compared against a primary standard (Trolox). Results were expressed as µmol Trolox equivalents/100 g of sample.

#### Isoespiptanol cytotoxicity

MTT method described by Mosmann (Mosmann, 1983) with murine

**Table 1.** Fatty acid composition of palm olein

Time (s)	Fatty acid	Carbons: Unsaturation	% Fatty acid
19.4	Palmitic Acid	16: 0	5.91
21.1	Methyl pentadecanoate	15: 0	5.86
22.5	Oleic Acid	18: 1 <sup>Δ9</sup>	31.94
24.9	Stearic Acid	18: 0	10.01
25.3	vaccenic acid	18: 1 <sup>Δ11</sup>	20.00
26.2	Linoleic acid (Omega 6)	18: 2 <sup>Δ9,12</sup>	0.82
27.1	11 - eicosenoic	20: 1 <sup>Δ11</sup>	5.50
29.9	eicosanoic	20: 0	1.04
30.3	docosanoic	22: 0	8.43
31.3	15-tetracosenoic	24: 1 <sup>Δ9</sup>	7.22
35.9	tetradecanoic	24: 0	0.25
36.7	hexacosanoic	26: 0	0.52

macrophages RAW 264.7 was used to measure the cytotoxic effect of Isoespiritanol. This assay is based on reduction of 3-(4,5-dimethylthiazol-2-yl) -2,5-diphenyl tetrazolium bromide (MTT) to Formazan in a reaction catalyzed by mitochondrial succinyl dehydrogenase.

The cells stimulated with LPS (1 µg/mL) were mixed with Isoespiritanol (100 µM) and incubated in microplate 96-well at 37°C for 24 h in atmosphere with 5% CO<sub>2</sub> in DMEM medium containing 2 mM L-glutamine (Gibco), 100 U/ml penicillin (Gibco), 100 µg/ml streptomycin without fetal bovine serum. The control contained 0.5% DMSO. After incubation, the culture medium was removed and 100 µL of MTT (0.5 mg/ml) were added. When deposits appeared blue, all liquid was removed and 100 µL of DMSO was added to dissolve Formazan crystals. Absorbance at 490 nm was measured (Spectrophotometer Multiskan Spectrum UV-VIS (Thermo Scientific)). The number of living cells is proportional to the amount of Formazan produced. The increase in absorbance indicates cell viability and this was determined using the following equation:

$$\% \text{ Viability} = \frac{A_p}{A_c} \times 100 \quad (2)$$

$A_p$  = Arithmetic mean of the absorbance of the test sample;  $A_c$  = Arithmetic mean absorbance of control.

### Statistical analysis

All the antioxidants experiments were performed four times. Data are presented as the mean ± the standard error (SE) and were processed by one-way analysis of variance (ANOVA). Multiple comparisons of the means were performed using Duncan's New Multiple Range Test ( $p < 0.05$ ). All of the statistical analysis was performed using the Statgraphics Centurion V statistical package for Windows. For testing cytotoxic effect, data was generated with an experimental design of randomized complete block, taking sample from each donor as blocking factor. The analysis of variance and the corresponding mean tests (Duncan) were performed using the statistical software Statgraphics Plus, version 5.0.

## RESULTS AND DISCUSSION

Degree of unsaturation is a factor that affects directly the


kinetics of degradation of oils (Guillén and Cabo, 2002; Choe and Min, 2006; García-Moreno et al., 2013). Oils with a high content of unsaturated fatty acids, especially polyunsaturated, are more susceptible to oxidation. Palm olein has 12 main fatty acids, with a ratio of unsaturated/saturated (I/S) of 0.71 and with chains of 15-26 carbons. Table 1 summarizes the percentage of fatty acids. The oleic acid (ω9) (31.9%), vaccenic (20%), 15-tetracosenoic (7.22%), stearic (10.01%) and docosanoic (8.43%) percentages are highlighted; also, a low amount of polyunsaturated fatty acids was found (less than 1%).

Palm olein has a high content of monounsaturated (64.66%) and saturated (32.02%) fatty acids and a low content of polyunsaturated fatty acids (0.82%). This gives high stability against oxidative processes as frying, as it is recommended for use in domestic and industrial level (Mozzon et al., 2013).

The ratio of monounsaturated/saturated fatty acids is an indicator of oxidative stability in oils; however, this does not completely define their behavior in the frying process where high temperatures are used for longer periods. In addition, the presence of dissolved oxygen in the oil and water from the food causes a series of reactions where different oxidized compounds are produced. Hence, the addition of antioxidants becomes an important need in the food industry.

Figure 1 shows the formation of hydroperoxides in the palm olein supplemented with Isoespiritanol (IE200) and BHT (BHT200) and without antioxidants. The concentration of hydroperoxides increases with time after 60 h of heating and is less in the presence of antioxidants. However, Isoespiritanol delays hydroperoxides production in greater proportion than BHT.

The Active Oxygen Method (AOM) (Method Cd 12-57) was used to monitor the oxidative state of palm olein, which determines the time that the oil takes to reach a peroxide value (PV) of 100 meq O<sub>2</sub>/kg oil (induction time; IT). Induction times were 89.68, 93.95 and 105.09 h for the control, BHT200 and IE 200, respectively. This shows



**Figure 1.** Formation of peroxides in the Palm Olein supplemented with (▲) Isoespintanol 200 mg / L, (○) BHT 200 mg / L and (◆) Control without antioxidant.

that in the presence of Isoespintanol, the induction time is reduced in 17.2% compared to the control, while the use of BHT only implies a reduction of 4.2%. These findings are attributed to the higher capacity of Isoespintanol to trap peroxy radicals ( $\text{ROO}\cdot$ ) generated in the oxidative process compared to BHT. Previous studies showed that the effectiveness of rosemary extracts in the oxidative stabilization of lipids is greater than commercial mixtures of synthetic antioxidants such as Butyl Hydroxytoluene (BHT), Butyl Hydroxyanisole (BHA) and Propyl Gallate (PG) (Upadhyay et al., 2006; Piedrahita et al., 2015).


Antioxidants are used to improve the oxidative stability of fats and oils that are used at temperatures above  $110^\circ\text{C}$  and under aeration for extended periods (Santos et al., 2012; Upadhyay and Mishra 2015b). However, some natural and synthetic antioxidants decompose or volatilize below  $110^\circ\text{C}$ , which leads to a decrease in their protective ability. We used thermogravimetry to assess the thermal stability of the studied antioxidants. Dynamic thermogravimetric curves indicated that mass loss is initiated at temperatures below  $100^\circ\text{C}$  for both, Isoespintanol and BHT (Figure 2). However, Isoespintanol exhibits a higher overall thermal resistance compared to BHT. The complete evaporation of Isoespintanol occurs at  $158.6^\circ\text{C}$  unlike BHT that is volatilized completely at  $144.8^\circ\text{C}$ .

The volatility and instability of synthetic antioxidants at high temperatures are known issues in the industry which have prompted the search for natural alternatives. The results suggest that BHT provides lower levels of protection compared to Isoespintanol due to its volatility. Similar results were obtained by Santos et al. (2012), who found that natural antioxidants display higher thermal resistance than synthetic antioxidants, a phenomenon


attributed to the difference on chemical bonds type of the natural antioxidants. Although BHT and Isoespintanol have similar molecular masses (220.35 and 210 g/mol, respectively) (Figure 3), the atomic bonding pattern is different. For instance, the type of substitution in the phenolic nucleus makes them energetically different.

H-ORACFL, L-ORACFL, ORACDCFH and ORACRPG values for Isoespintanol and BHT are presented in Figure 4. The results show that the Isoespintanol has substantially greater antioxidant capacity than BHT under all experimental scenarios, which could explain its ability to oxidatively stabilize palm olein under accelerated conditions of oxidation.

ORAC assays using fluorescein (FL) have limitations because they are influenced more by stoichiometry than by reactive factors (Lopez-Alarcon and Lissi, 2006). An additional methodology has been proposed to determine the ORAC value based on pyrogallol red discoloration (RPG), which provides values that are exclusively dependent on the reactivity of the tested compounds rather than the amount; thereof, this concept is important considering the function of an antioxidant. This difference is due to the high reactivity of RPG and the compound concentrations used in the experiments compared to low levels of ORAC probes with fluorescein. One way to check the quantity and the quality of the test compound is to determine  $\text{ORAC}_{\text{FL}}$  and  $\text{ORAC}_{\text{RPG}}$ . The results show that Isoespintanol exhibits the perfect setting, between quantity and quality, revealing greater results in both methods compared to BHT. Besides, these results are positive considering that the National Institutes of Health considers that BHT is a promoter of carcinogenesis in humans, so its use as a food additive is not recommended. BHT was toxic for mouse lung cells



**Figure 2.** Curves (a) TG and (b) DTG Isoespintanol and BHT.



**Figure 3.** Chemical structure of antioxidants (a) Isoespintanol (IE) and (b) Butylated hydroxytoluene (BHT).

(necrosis type I pneumocytes and endothelial cells), and there is also evidence of hepatotoxicity. BHT also has adverse effects on reproduction and increases the effect of certain carcinogens such as urethane and N-fluorenilacetamida.

This study revealed that Isoespintanol at 100  $\mu$ M does not have significant cytotoxic effects against murine macrophages (RAW 264.7) (Figure 5). The results are positive considering the possible use of Isoespintanol as

a food additive.

Although there is not a clear mechanism that describes the way that the mono- and phenols act on cell protection, we can suggest that due to the high capacity of Isoespintanol to trap oxygen radicals, as evidenced by their ORAC values, the harmful effects of reactive species present in the cell are counteracted (Duthie et al., 2003). Previous studies found that thymol and carvacrol, structural analogues of Isoespintanol, showed protective




Figure 4. Oxygen Radical Antioxidant Capacity (ORAC) for Isoespitanol and BHA.




Figure 5. Evaluation by MMT of Isoespitanol Cytotoxicity in murine macrophages (RAW 264.7)

effects on DNA from human lymphocytes at low concentrations (0.5 to 50 mM). However, pro-oxidant effects were observed at concentrations greater than 100 µM (Aydin et al., 2005). Numerous studies have shown that some polyphenolic antioxidants such as quercetin, rutin or epicatechingallate have genotoxic effects at high concentrations (Perez, 2003; Johnson and Loo, 2000) contrary to the findings of this study in respect of Isoespitanol, which represents a great advantage.

## Conclusions

Palm olein has a high content of monounsaturated and

saturated fatty acids (96.68%) and a negligible content of polyunsaturated fatty acids (0.82%), reflecting stability against oxidative phenomena. However, processes such as frying are adverse and accelerate deterioration reactions. For this reason, the addition of antioxidants becomes an important step in the food industry. The results revealed that hydroperoxides production under oxidation accelerated conditions in palm olein is 39% slower in the presence of Isoespitanol (200 mg/L) and only 10% slower in the presence of BHT. These results are attributed to the superior antioxidant capacity observed for Isoespitanol and demonstrate the ability of this molecule to extend the conservation process in oils subjected to high temperatures. The dynamic thermo-

gravimetric curves for Isoespintanol and BHT indicated that the mass loss for both antioxidants is initiated at temperatures below 100°C, however, Isoespintanol exhibited higher overall thermal resistance than BHT, since complete decomposition took place at 157 and 144, respectively. Finally, cytotoxicity tests with 100 µM Isoespintanol against murine RAW 264.7 macrophages showed no effect on cell viability, indicating their possible use as a safe food additive.

## CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

## REFERENCES

- Aladedunye F, Przybylski R (2013). Frying stability of high oleic sunflower oils as affected by composition of tocopherol isomers and linoleic acid content. *Food Chem.* 141(3):2373-2378.
- Alvis A, Vélez C (2008). Modeling of the process of frying yams (*Dioscoreaalata*) through reological measurements using the methodology of response surface. *Información Tecnol.* 19(5):11-18.
- Aydin S, Başaran A, Başaran N (2005). The effects of thyme volatiles on the induction of DNA damage by the heterocyclic amine IQ and mitomycin C. *Mutat. Res.* 581:43-53.
- Casal S, Malheiro R, Sendas A, Oliveira B (2010). Pereira, J.A. Olive oil stability under deep-frying conditions. *Food Chem. Toxicol.* 48(10):2972-2979.
- Choe E, Min D (2006). Chemistry and Reactions of Reactive Oxygen Species in Foods. *Crit. Rev. Food Sci. Nutr.* 46:1-22.
- Duthie S, Collins AR, Duthie GG, Dobson V (2003). Quercetin and myrcetin protect against hydrogen peroxide-induced DNA damage (strand breaks and oxidized pyrimidines) in human lymphocytes. *Mutat. Res.* 393:223-231.
- García-Moreno PJ, Pérez-Gálvez R, Guadix A, Guadix EM (2013). Influence of the Parameters of the Rancimat Test on the Determination of the Oxidative Stability Index of Cod Liver Oil. *Food Sci. Technol.* 51:303-308.
- Guillén MD, Cabo N (2002). Fourier Transform Infrared Spectra Data versus Peroxide and Anisidine Values to Determine Oxidative Stability of Edible Oils. *Food Chem.* 77:503-510.
- Ismail R (2005). Palm oil and palm olein frying applications. *Asia Pac. J. Clin. Nutr.* 14(4):414-419.
- Johnson MK, Loo G (2000). Effects of epigallocatechingallate and quercetin on oxidative damage to cellular DNA. *Mutat. Res.* 459:211-218.
- Kalogianni EP, Karapantsios TD, Miller R (2011). Effect of repeated frying on the viscosity, density and dynamic interfacial tension of palm and olive oil. *J. Food Eng.* 105(1):169-179.
- Krishnaiah D, Sarbaty R, Nithyanandam RA (2011). Review of the antioxidant potential of medicinal plant species. *Food Bioprod. Processing* 89:217-233.
- Lopez-Alarcón C, Lissi EA (2006). Novel and simple ORAC methodology based on the interaction of Pyrogallol Red with peroxyl radicals. *Free Rad. Res.* 40(9):979-985.
- Martin-Romero FJ, Miguel-Lasobras EM, Domínguez-Arroyo JA, González-Carrera E, Alvarez IS (2008). Contribution of culture media to oxidative stress and its effect on human oocytes. *Reprod. BioMed.* Online 17(5):652-661.
- Mosmann T (1983). Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *J. Immunol. Methods* 65(1-2):55-63.
- Mozzon M, Pacetti D, Lucci P, Balzano M, Frega NG (2013). Crude palm oil from interspecific hybrid *Elaeioleifer a* × *Elaeisguineensis*: Fatty acid regiodistribution and molecular species of glycerides. *Food Chem.* 141(1):245-252.
- Niaiforovia N, Mihailovia V, Maskovia P, Solujia S, Stojkovic A, Pavlovic D, Muratspahia D (2010). Antioxidant activity of selected plant species; potential new sources of natural antioxidants. *Food Chem. Toxicol.* 48:3125-3130.
- Perez G (2003). Flavonoids: Antioxidants or prooxidants. *Rev. Cubana InvestBiomed.* 22(1):48-57.
- Piedrahita AP, Peñaloza J, Cogollo A, Rojano BA (2015). Kinetic Study of the Oxidative Degradation of Choibá Oil (*Dipteryxoleifer a*Benth.) with Addition of Rosemary Extract (*Rosmarinus officinalis* L.). *Food Nutr. Sci.* 6:466-479.
- Prior RL, Wu X, Schaich K (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J. Agric. Food Chem.* 43:4290-4302.
- Rautenbach F, Venter I (2010). Hydrophilic and lipophilic antioxidant capacity of commonly consumed South African fruits, vegetables, grains, legumes, fats/oils and beverages. *J. Food Compos. Anal.* 23(7):753-761.
- Rojano B, Gaviria CA, Gil MA, Saez JA, Schinella G, Tournier H (2008b). Antioxidant activity of the isoespintanol in different media. *Vitae* 15(1):173-181.
- Rojano B, Saez J, Schinella G, Quijano J, Velez E, Gil A (2008a). Experimental and theoretical determination of the antioxidant properties of isoespintanol (2-Isopropyl-3,6-dimethoxy-5-methylphenol). *J. MolStruct.* 877:1-6.
- Rojano B, Figadere B, Martin MT, Recio MC, Giner R, Schinella G (2007). Constituents of Oxandra cf. xylopioides with anti-inflammatory activity. *J. Nat. Prod.* 70(5):835-838.
- Sanchez A (2011). Vegetable oils and fats as functional ingredient in meat products. *Nacameh* 5:108-118.
- Santos NA, Cordeiro AM, Damasceno SS, Aguiar RT, Rosenhaim R, Carvalho-Filho JR, Santos IM, Maia AS, Souza A (2012). Commercial antioxidants and thermal stability evaluations. *Fuel* 97:638-643.
- Shantha NC, Decker EA (1994). Rapid, Sensitive, Iron-Based Spectrophotometric Methods for Determination of Peroxide values of food lipids. *J. AOAC Int.* 77:421-424.
- United States Department of Agriculture (USDA) 2004). United States Department of Agriculture, Agricultural Statistics 2004. Table 3-51.
- Upadhyay R, Mishra HN (2015a). Classification of Sunflower Oil Blends Stabilized by Oleoresin Rosemary (*Rosmarinus officinalis* L.) Using Multivariate Kinetic Approach. *J. Food Sci.* 80(8):1746-1754.
- Upadhyay R, Mishra HN (2015b). Multivariate Analysis for Kinetic Modeling of Oxidative Stability and Shelf Life Estimation of Sunflower Oil Blended with Sage (*Salvia officinalis*) Extract Under Rancimat Conditions. *Food Bioprocess Technol.* 8(4):801-810.
- Upadhyay R, Sehwag S, Mishra HN (2017). Chemometric approach to develop frying stable sunflower oil blends stabilized with oleoresin rosemary and ascorbyl palmitate. *Food Chem.* 1(218):496-504.
- Wu H, Karayiannis TG, Tassou SA (2013). Two-dimensional frying model for the investigation and optimisation of continuous industrial frying systems. *Appl. Ther. Eng.* 51(1-2):926-936.
- Zapata A, Cogollo PA, Rojano BA (2013). Nutraceuical potential of almond oil of *choibá* or mountain almond (*Dipteryxoleifer a*Benth.). *Rev. Cub. Plantas Med.* 18:368-380.
- Zapata K, Piedrahita AM, Alzatea AF, Cortés FB, Rojano BA (2015). Oxidative Stabilization of Sacha Inchi (*Plukenetia Volubilis* Linneo) Oil with Mortiño (*Vaccinium Meridionale* SW) suspensions addition. *Rev. Cien. Desarrollo* 6(2):141-153.