

Full Length Research Paper

Antimicrobial activity of native and naturalized plants of Minnesota and Wisconsin

Joy R. Borchardt¹, Donald L. Wyse¹, Craig C. Sheaffer¹, Kendra L. Kauppi², R. Gary Fulcher³
Nancy J. Ehlke¹, David D. Biesboer⁴ and Russell F. Bey²

¹Department of Agronomy and Plant Genetics, University of Minnesota, 411 Borlaug Hall, St. Paul MN 55108-6026

²Department of Veterinary and Biomedical Sciences, University of Minnesota, 205 Veterinary Science, St. Paul, MN 55108-6187.

³Department of Food Science and Nutrition, University of Manitoba, Winnipeg, MB, R3T 2N2.

⁴Department of Plant Biology, University of Minnesota, 250 Biological Science Center, St. Paul, MN 55108-6022

Accepted 8, February, 2008

The antimicrobial activity of aqueous ethanol extracts of stems, leaves, flowers and roots from 336 native and naturalized species (597 extracts) collected in Minnesota and Wisconsin was tested against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Candida albicans*. Twenty-four percent, or 142 extracts, exhibited antimicrobial activity. Extracts from *Betula papyrifera* Marshall (Betulaceae), *Centaurea maculosa* Lam. (Asteraceae), *Epilobium angustifolium* L. (Onagraceae), *Hypericum perforatum* L. (Clusiaceae), *Lythrum salicaria* L. (Lythraceae), and *Rhus glabra* L. (Anacardiaceae) inhibited all four microorganisms. Extracts from two species inhibited three microorganisms, 11 extracts (10 species) inhibited two, and 119 extracts (98 species) inhibited one microorganism with four species having inhibition zones greater than 15 mm. This is the first report describing the antimicrobial activity of *Clintonia* sp. (Liliaceae), *Comptonia peregrina* (L.) J.M. Coult. (Myricaceae), *Desmodium illinoense* A. Gray (Fabaceae), *Geum virginianum* L. (Rosaceae), leaves of *Scirpus americanus* Pers. (Cyperaceae), flower clusters of *Eupatorium maculatum* L. (Asteraceae), berries of *Smilacina racemosa* (L.) Desf (false Solomon's seal) and frozen *Hypericum perforatum* L. (Clusiaceae).

Key words: Antimicrobial, medicinal, native plants, antibacterial.

INTRODUCTION

New antimicrobial agents are needed to treat diseases in humans and animals caused by drug resistant microorganisms. In addition, there is a continuing consumer demand for "natural" and/or "preservative-free" microbiologically safe foods and cosmetic products (Wijesekera, 1991; Zink, 1997). As public demand for these products increases, an opportunity exists to satisfy consumer demands while providing wholesome and safe products from plants.

Antimicrobial compounds of plant origin may occur in stems, roots, leaves, bark, flowers and fruits of plants. Plant derived phytoalexin (Beuchat et al., 1994) isothiocyanates (Delaquis and Mazza, 1995) alllicins, anthocya-

nins (Somaatmadja et al., 1964) and essential oils (Lis-Balchin and Deans, 1997) tannins and polyphenols and terpenoids (Cutter, 2000; Hao et al., 1998; Puupponen-Pimia et al., 2001) have demonstrated antibacterial and/or antifungal activities. These compounds are bactericidal and/or bacteriostatic influencing lag time, growth rate and maximum growth of microorganisms.

Herbal medicine expertise of North American Native Indian cultures has been documented for the states of Minnesota and Wisconsin in many publications in the popular literature. Indigenous herbal medicine includes knowledge regarding the appropriate plant parts, extraction, and manner of preparation as infusions, decoctions, or poultices. Often, different plant parts have had specific ethnomedical applications. For instance, the flowers, seeds and roots of *Rhus* and *Epilobium* spp. have had antiemetic, antidiarrheal, oral and respiratory aid, anti-hemorrhagic, dermatological aid, and analgesic applications

*Corresponding author. E-mail: wyses001@umn.edu. Tel: 612-625-7064; Fax: 612-625-1268.

(Moerman, 2004). Although a specific plant part might have a reported use, other parts of the plant and additional applications may remain uninvestigated or a plant may not have a recorded ethnomedical use.

Research has identified some North American species as potential medicinal crops (Small and Catling, 1999). An example is *Taxus brevifolia* Nutt., the Pacific Yew. This species has a taxane diterpenoid in the bark that is useful in treating some forms of cancer (Wani et al., 1971). In addition, *Achillea millefolium* L., *Acorus calamus* L., *Caulophyllus* species, *Echinacea pallida* (Nutt.) Nutt. var. *angustifolia* (DC.) Cronq., *Epilobium angustifolium* L., *Oenothera biennis* L., *Podophyllum peltatum* L., and *Taraxacum* species have been considered as medicinal crops (Small and Catling, 1999). Another example is *P. peltatum* L. that is being considered as an alternative crop in the southern United States (Cushman et al., 2005).

Another source of antimicrobial compounds is naturalized non-native plant introductions into North America, yet few of these species have been studied for this purpose. An example is *Lythrum salicaria* L. (purple loosestrife) that has invaded North American wetlands. *L. salicaria* has styptic and antibacterial action that can assist wound healing (Thompson et al., 1987).

A significant opportunity exists to identify new, natural plant derived antimicrobial agents for treatment of diseases or as food or cosmetic preservatives. Our objective was to evaluate the activity of aerial parts of native and naturalized species in the Upper Mississippi River Basin against organisms that cause disease and spoilage, that is, *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Candida albicans*.

MATERIALS AND METHODS

Plant collection

In 2003 and 2004 aerial parts (leaves, stems and flowers) and an occasional root of native and naturalized plants were collected in Minnesota and Wisconsin. The availability of flowers on some species depended on the time of year. In 2003, 301 plant samples were collected and in 2004, 296 plant samples were collected for a total of 336 species. In 2003, all the plant samples were frozen at -10°C immediately after collection and stored from one to four months before extraction and antimicrobial activity determination. In 2004, plant samples were frozen (-10°C) immediately after collection, and within 48 h were extracted and tested.

Preparation of extracts

Ten grams of plant material were cut into 2.5 cm pieces and combined with 10 ml of 80:20 aqueous ethanolic solvent. The mixture was pulverized with a stomacher for 2 min. The plant material varied in absorbency; if 2 ml of extract could not be pipetted, an additional 2 ml of solvent was added. The macerated mixture was left at room temperature (20°C) for 24 h. The stomacher bags were kneaded and approximately 2 ml of supernatant was transferred into a sterile microfuge tube and centrifuged at 15,000 xg for 15 min to remove remaining plant solids. Extracts

of 50 µl were placed on 6 mm sterile paper discs then the discs and solvent were placed in a biological safety cabinet where the solvent was allowed to evaporate for one hour.

Antimicrobial screening

The extracts were screened for antimicrobial activity using the disk diffusion technique (Bauer et al., 1966). Test microorganisms included *S. aureus* (ATCC 12600), *E. coli* (ATCC 8677), *P. aeruginosa* (ATCC 9721) and *C. albicans* (ATCC 10231). *E. coli* and *P. aeruginosa* were maintained on MacConkey's agar (Becton Dickinson and Company, Microbiology Systems, Sparks, MD), *S. aureus* on blood agar and *C. albicans* on Sabouraud dextrose agar (Becton Dickinson and Company, Microbiology Systems, Sparks, MD). After 18 to 20 h of culture, each microorganism was diluted in sterile saline to an optical density of approximately 0.5 using a MacFarland standard (Becton Dickinson and Company, Microbiology Systems, Sparks, MD). A Mueller-Hinton agar plate was swabbed on three axes with a sterile cotton tipped swab, which was dipped in the freshly prepared, diluted culture. Discs containing the dried extract were placed on the freshly swabbed plates along with controls. Discs with evaporated solvent were used as a negative control and an antibiotic disc (ticarcillin, 75 mcg or chloramphenicol 30 mcg) was used as a positive control. The plates were incubated at 37°C for 18 h and zones of inhibition were measured in millimeters on three axes and the mean value reported.

RESULTS AND DISCUSSION

Of the 597 samples, 142 samples (24%), representing 109 different species and 53 plant families exhibited antimicrobial activity against at least one microorganism (Table 1). Extracts from six plants (*B. papyrifera* Marshall, *Centaurea maculosa* Lam, *E. angustifolium* L., *Hypericum perforatum* L., *L. salicaria* L. and *Rhus glabra* L.) inhibited growth of the four test microorganisms. Extracts from 455 samples exhibited no antimicrobial activity (Table 2).

Plant extracts inhibiting four microorganisms

Betula papyrifera

Extracts of *B. papyrifera* Marshall (paper birch) leaves were most effective against *S. aureus* and had slightly inhibited the other three microorganisms. Previous research showed that leaf extracts (aqueous acetone) of *Betula pubescens* Ehrh, the old-world counterpart to *B. papyrifera* also had antimicrobial activity against *S. aureus*, with slight activity against *E. coli*, and no activity against *C. albicans* or *Aspergillus niger* (Rauha et al., 2000). Ethanol extracts of *B. papyrifera* bark and wood exhibited antimicrobial activity against our four bacterial species (two Gram-positive and two Gram-negative) but no antifungal activity (Omar et al., 2000). Methanolic extracts of air-dried *B. papyrifera* branches had activity against 6 of 11 bacteria (four Gram-positive, two Gram-negative) and three fungal species (McCutcheon et al., 1992; 1994). The antimicrobial activity appears to be eva-

Table 1. Antimicrobial activity of aqueous/ethanol plant extracts from aerial parts of plants collected in Minnesota and Wisconsin, U.S.A. in 2003 and 2004.

| Botanical name | Common name | Plant part tested | Microorganisms ^a | | | |
|---|------------------------|--------------------------|-----------------------------|----|----|----|
| | | | Inhibition zones in mm | | | |
| Inhibition against four microorganisms | | | Sa | Ec | Pa | Ca |
| <i>Betula papyrifera</i> Marshall ¹⁰ | paper birch | leaves | 9 | sl | sl | sl |
| <i>Centaurea maculosa</i> Lam. ⁸ | spotted knapweed | leaves | 11 | 10 | 7 | 7 |
| <i>Epilobium angustifolium</i> L. ³⁷ | fireweed | leaves | 11 | 7 | 7 | 15 |
| <i>Epilobium angustifolium</i> L. ³⁷ | fireweed | flowering aerial organs | 17 | 6 | 7 | 20 |
| <i>Hypericum perforatum</i> L. ¹⁷ | St. John's Wort | flowering aerial organs | 15 | 6 | 6 | 7 |
| <i>Lythrum salicaria</i> L. ³² | purple loosestrife | flowering spikes, leaves | 15 | 10 | 7 | 7 |
| <i>Rhus glabra</i> L. ³ | sumac, smooth | green flower clusters | 16 | 10 | 12 | 13 |
| <i>Rhus glabra</i> L. ³ | sumac, smooth | leaves | 13 | 7 | 10 | 15 |
| Inhibition against three microorganisms | | | | | | |
| <i>Desmodium illinoense</i> A. Gray ²³ | prairie tick trefoil | flowering aerial organs | sl | | sl | sl |
| <i>Scirpus americanus</i> Pers. ¹⁹ | bulrush | leaves | 10 | sl | sl | |
| Inhibition against two microorganisms | | | | | | |
| <i>Clintonia</i> sp. ³¹ | Clintonia | leaves, roots | 11 | | | 11 |
| <i>Comptonia peregrina</i> (L.) J.M. Coult. ³⁵ | sweet fern | aerial organs | 9 | | | 15 |
| <i>Comptonia peregrina</i> (L.) J.M. Coult. ³⁵ | sweet fern | leaves | 14 | | | 12 |
| <i>Cotinus coggygria</i> Scop. ³ | smoke tree | leaves | 13 | | 10 | |
| <i>Desmanthus illinoensis</i> (Michx.) MacMill. ³³ | Illinois bundle flower | leaves, stems, flowers | 15 | | | 17 |
| <i>Epilobium angustifolium</i> L. ³⁷ | fireweed | flowers | 14 | | | 17 |
| <i>Epilobium ciliatum</i> Raf. ³⁷ | Amer. willow herb | leaves | 11 | | | 15 |
| <i>Eupatorium maculatum</i> L. ⁸ | Joe pye weed | flower clusters | 9 | | | sl |
| <i>Geum virginianum</i> L. ⁴⁴ | rough avens | immature floral organs | 10 | | | 10 |
| <i>Juglans nigra</i> L. ²⁹ | black walnut | leaves | 15 | | | 12 |
| <i>Polygonum coccineum</i> L. ⁴¹ | swamp smartweed | flowers | 11 | | | 7 |
| Inhibition against a single microorganism | | | | | | |
| <i>Adiantum pedatum</i> L. ¹ | maidenhair fern | leaves | 7 | | | |
| <i>Aesculus glabra</i> Willd. ²⁸ | Ohio buckeye | leaves | 8 | | | |
| <i>Allium ramosum</i> L. ² | Chinese chives | leaves | 7 | | | |
| <i>Ambrosia artemesifolia</i> L. ⁸ | ragweed | flowering aerial organs | 10 | | | |
| <i>Amorpha canescens</i> L. ²³ | lead plant | leaves | 7 | | | |
| <i>Amorpha canescens</i> L. ²³ | lead plant | flowering aerial organs | 8 | | | |
| <i>Anaphalis margaritacea</i> (L.) Benth. and Hook ⁸ | pearly everlasting | leaves | 18 | | | |
| <i>Anaphalis margaritacea</i> (L.) Benth. and Hook ⁸ | pearly everlasting | flowering aerial organs | 15 | | | |
| <i>Anaphalis margaritacea</i> (L.) Benth. and Hook ⁸ | pearly everlasting | flower | 10 | | | |
| <i>Anemone quinquefolia</i> L. ⁴³ | snowdrop | leaves | | | | 12 |
| <i>Apocynum andromaeifolium</i> L. ⁷ | spreading dogbane | leaves stem, stalk | 10 | | | |
| <i>Apocynum androsaemifolium</i> L. ⁷ | spreading dogbane | leaves | 12 | | | |
| <i>Apocynum cannabinum</i> L. ⁷ | prairie dogbane | leaves | 7 | | | |
| <i>Asarum canadense</i> L. ⁶ | wild ginger | leaves w/ stems | 15 | | | |
| <i>Asclepias tuberosa</i> L. ⁷ | butterfly weed | flowering aerial organs | 7 | | | |

Table 1. contd.

| | | | | | | |
|---|----------------------|----------------------------------|----|--|--|----|
| <i>Baptisia australis</i> (L.) R.Br. ²³ | false indigo | leaves w/ few flowers | sl | | | |
| <i>Cannabis sativa</i> L. ¹⁴ | hemp | leaves, stems | 25 | | | |
| <i>Chamaecrista fasciculata</i> (L.) Moench ²⁴ | partridge pea | leaves | 11 | | | |
| <i>Chrysanthemum leucanthemum</i> L. ⁸ | oxeye daisy | flowers | 7 | | | |
| <i>Cicuta maculata</i> L. ⁴ | water hemlock | flowering aerial organs | sl | | | |
| <i>Cicuta maculata</i> L. ⁴ | water hemlock | flower clusters | sl | | | |
| <i>Claytonia virginica</i> L. ⁴² | spring beauty | leaves | sl | | | |
| <i>Coreopsis palmata</i> Nutt. ⁸ | prairie coreopsis | leaves | | | | 8 |
| <i>Cornus amomum</i> L. ¹⁸ | silky dogwood | leaves | 10 | | | |
| <i>Cornus stolonifera</i> Mill. ¹⁸ | red-osier dogwood | flower clusters, leaves | 10 | | | |
| <i>Corylus</i> sp. ¹⁰ | hazelnut | seed clusters, leaves | 10 | | | |
| <i>Cotinas coggygria</i> Scop. ³ | smoketree | flowers, leaves, stem | 11 | | | |
| <i>Desmodium canadense</i> (L.) DC ²³ | showy tick trefoil | leaves | 12 | | | |
| <i>Dicentra eximia</i> (Ker Gawl.) Torr. ²⁵ | bleeding heart | leaves, stem | 10 | | | |
| <i>Diervilla lonicera</i> Mill. ¹⁵ | bush honeysuckle | leaves, stem, flowers | 9 | | | |
| <i>Echinacea purpurea</i> (L.) Moench ⁸ | purple coneflower | leaves | 10 | | | |
| <i>Equisetum sylvaticum</i> L. ²⁰ | woodland horsetail | leaves | 10 | | | |
| <i>Eucalyptus</i> sp. ³⁶ | eucalyptus | leaves | 15 | | | |
| <i>Eupatorium maculatum</i> L. ⁸ | Joe pye weed | leaves, stems | 8 | | | |
| <i>Eupatorium perfoliatum</i> L. ⁸ | boneset | leaves | 11 | | | |
| <i>Eupatorium perfoliatum</i> L. ⁸ | boneset | flower clusters | 7 | | | |
| <i>Euphorbia corollata</i> L. ²² | flowering spurge | leaves | 10 | | | |
| <i>Euphorbia corollata</i> L. ²² | flowering spurge | aerial parts | 7 | | | |
| <i>Euphorbia esula</i> L. ²² | leafy spurge | flowers | | | | 9 |
| <i>Euphorbia maculata</i> L. ²² | milk purslane | leaves | | | | 14 |
| <i>Filipendula rubra</i> (Hill) B.L.Rob. ⁴⁴ | Queen of the prairie | leaves | 12 | | | |
| <i>Geranium maculatum</i> L. ²⁶ | wild geranium | flowers | 7 | | | |
| <i>Geranium maculatum</i> L. ²⁶ | wild geranium | leaves, stems | 9 | | | |
| <i>Geum triflorum</i> Pursh ⁴⁴ | prairie smoke | leaves, stems | 7 | | | |
| <i>Glycyrrhiza lepidota</i> Pursh ²³ | wild licorice | leaves | 11 | | | |
| <i>Helenium autumnale</i> L. ⁸ | sneezeweed | leaves | 7 | | | |
| <i>Helianthus giganteus</i> L. ⁸ | giant sunflowers | flowers | sl | | | |
| <i>Heracleum lanatum</i> L. ⁴ | cow parsnip | seed heads, stems | 10 | | | |
| <i>Heracleum lanatum</i> L. ⁴ | cow parsnip | leaves | 7 | | | |
| <i>Hesperis matronalis</i> L. ¹² | dame's rocket | leaves | 14 | | | |
| <i>Hesperis matronalis</i> L. ¹² | dame's rocket | flower | 12 | | | |
| <i>Heuchera richardsonii</i> R.Br. ⁴⁷ | prairie alum root | leaves | 13 | | | |
| <i>Hypericum perforatum</i> L. ¹⁷ | St. John's Wort | aerial parts, immature fruits | 11 | | | |
| <i>Ilex verticillata</i> L. ⁵ | winterberry | branch w/ leaves, berries | 9 | | | |
| <i>Impatiens capensis</i> Meerb ⁹ | jewelweed | leaves | sl | | | |
| <i>Juglans nigra</i> L. ²⁹ | black walnut | leaves | 11 | | | |
| <i>Larix laricina</i> (Du Roi) K. Koch ³⁹ | tamarack | leaves | 7 | | | |
| <i>Ledum groenlandicum</i> Oeder ²¹ | Labrador tea | leaves | 9 | | | |
| <i>Liatris pycnostachya</i> Michx. ⁸ | prairie blazing star | leaves | 10 | | | |
| <i>Lythrum salicaria</i> L. ³² | purple loosestrife | leaves | 9 | | | |
| <i>Miscanthus giganteus</i> (hybrid) ⁴⁰ | Chinese silver grass | leaves, stalk | 12 | | | |

Table 1. contd.

| | | | | | |
|--|----------------------|---|----|--|----|
| <i>Monarda fistulosa</i> L. ³⁰ | wild bergamot | leaves | 6 | | |
| <i>Monarda fistulosa</i> L. ³⁰ | bergamot | flowering aerial organs | 7 | | |
| <i>Morus rubra</i> L. ³⁴ | red mulberry | leaves | 11 | | |
| <i>Oenothera biennis</i> L. ³⁷ | evening primrose | leaves | sl | | |
| <i>Picea mariana</i> (Mill.) Britton, Sterns and Poggenb. ³⁹ | black spruce | leaves | 9 | | |
| <i>Pinus strobes</i> L. ³⁹ | white pine | flowers | 10 | | |
| <i>Polygonum coccineum</i> L. ⁴¹ | swamp smartweed | leaves | 14 | | |
| <i>Polygonum coccineum</i> L. ⁴¹ | swamp smartweed | flowering aerial organs | 11 | | |
| <i>Polygonum cuspidatum</i> L. ⁴¹ | Japanese knotweed | leaves | 9 | | |
| <i>Polygonum persicaria</i> L. ⁴¹ | lady's thumb | leaves, stems | 7 | | |
| <i>Potentilla arguta</i> Pursh ⁴⁴ | tall potentilla | leaves | | | 16 |
| <i>Potentilla simplex</i> Michx. ⁴⁴ | common cinquefoil | flowering aerial organs | 11 | | |
| <i>Prunella vulgaris</i> L. ³⁰ | heal-all | flowering aerial organs | 7 | | |
| <i>Prunus americana</i> Marshall ⁴⁴ | plum | leaves | 7 | | |
| <i>Pycnanthemum virginianum</i> (L.) Durand B.D. Jacks. ³⁰ | mountain mint | leaves | 9 | | |
| <i>Quercus rubra</i> L. ²⁴ | red oak | leaves | | | 9 |
| <i>Rhus typhina</i> L. ³ | staghorn sumac | berries | sl | | |
| <i>Ribes</i> sp. ²⁷ | gooseberry | leaves, twigs | 10 | | |
| <i>Rosa palustris</i> Marshall ⁴⁴ | swamp rose | aerial parts w/ rose hips | 13 | | |
| <i>Rubus</i> sp. ⁴⁴ | wild raspberry | leaves | 9 | | |
| <i>Rumex acetosella</i> L. ⁴¹ | sheep sorrel | immature flowers | 11 | | |
| <i>Rumex crispus</i> L. ⁴¹ | curly dock | immature flowers | 12 | | |
| <i>Rumex crispus</i> L. ⁴¹ | curly dock | stems of immature flowers | 11 | | |
| <i>Ruta graveolens</i> L. ⁴⁵ | rue | leaves | sl | | |
| <i>Salix petolaris</i> Sm. ⁴⁶ | meadow willow | leaves | 9 | | |
| <i>Sanguinaria Canadensis</i> L. ³⁸ | bloodroot | leaves w/ stems | sl | | |
| <i>Scirpus americanus</i> Pers. ¹⁹ | bulrush | flower clusters | 8 | | |
| <i>Scirpus validus</i> Vahl ¹⁹ | great bulrush | leaves | 7 | | |
| <i>Senna marilandica</i> (L.) Link ¹³ | wild senna | leaves | 10 | | |
| <i>Silene latifolia</i> Poir ¹⁶ | white campion | stems, leaves | 9 | | |
| <i>Smilacina racemosa</i> (L.) Desf. ³¹ | false Solomon's seal | leaves stem, seeds | | | 14 |
| <i>Smilacina racemosa</i> (L.) Desf. ³¹ | false Solomon's seal | berries | | | 17 |
| <i>Solanum dulcamara</i> L. ⁴⁹ | nightshade | berries, mixed maturity flower, leaves, stems, | | | 11 |
| <i>Solanum dulcamara</i> L. ⁴⁹ | nightshade | fruits | | | 11 |
| <i>Solidago rigida</i> L. ⁸ | stiff goldenrod | leaves | sl | | |
| <i>Solidago</i> sp. ⁸ | golden rod | leaves | 7 | | |
| <i>Spiraea alba</i> Du Roi ⁴⁴ | meadowsweet | flowers, leaves, stems | 15 | | |
| <i>Spiraea tomentosa</i> L. ⁴⁴ | steeplebush | leaves | 15 | | |
| <i>Spiraea tomentosa</i> L. ⁴⁴ | steeplebush | flowers, leaves, stems | 10 | | |
| <i>Spiraea tomentosa</i> L. ⁴⁴ | steeplebush | flowering aerial organs | 13 | | |
| <i>Symphytum officinale</i> L. ¹¹ | comfrey | leaves | 9 | | |
| <i>Tanacetum vulgare</i> L. ⁸ | tansy | leaves, stems | sl | | |
| <i>Tanacetum vulgare</i> L. ⁸ | tansy | flower clusters | 7 | | |
| <i>Taxus</i> sp. ⁵⁰ | yew | new spring growth | sl | | |
| <i>Tephrosia virginiana</i> (L.) Pers. ²³ | goats rue | leaves | 11 | | |

Table 1. contd.

| | | | | | | |
|--|--------------------|---------------------------|----|----|----|----|
| <i>Thalictrum dasycarpum</i> Fisch. and Ave'-Lall. ⁴³ | purple meadow rue | leaves, stem | 7 | | | |
| <i>Thalictrum dioicum</i> L. ⁴³ | early meadow rue | leaves, stem | 7 | | | |
| <i>Tilia americana</i> L. ⁵¹ | basswood | developing seeds | 10 | | | |
| <i>Tilia americana</i> L. ⁵¹ | basswood | flowers clusters, bracts | 7 | | | |
| <i>Tragopogon pratensis</i> L. ⁸ | oyster plant | seed heads | 9 | | | |
| <i>Trillium grandiflorum</i> (Michx.) Salisb ³¹ | big white trillium | leaves | | | | sl |
| <i>Vaccinium angustifolium</i> Aiton ²¹ | blueberry | leaves, stems | 7 | | | |
| <i>Verbascum thapsus</i> L. ⁴⁸ | mullein | flowering spike | sl | | | |
| <i>Vernonia fasciculata</i> Michx. ⁸ | common ironweed | leaves | sl | | | |
| <i>Veronicastrum virginicum</i> (L.) Farw. ⁴⁸ | Culver's root | leaves | 7 | | | |
| <i>Vitis aestivalis</i> Michx. ⁵² | wild grape | leaves | 7 | | | |
| <i>Zizia aurea</i> L. ⁴ | golden alexanders | seed heads, stems, leaves | 15 | | | |
| Controls | | | | | | |
| Aqueous ethanol | | | | | | |
| Ticarcillin | | | | 27 | 20 | |
| Chloramphenicol | | | 27 | | | |

Sa=Staphylococcus aureus; Ec= Escherichia coli; Pa= Sa=Staphylococcus aureus; Ec= Escherichia coli; Pa=Pseudomonas aeruginosa; Ca=Candida albicans; sl=slight Numerical superscripts refer to familial names for each species:

¹Adiantaceae, ²Alliaceae, ³Anacardiaceae, ⁴Apiaceae, ⁵Aquifoliaceae, ⁶Aristolochiaceae, ⁷Asclepiadaceae, ⁸Asteraceae, ⁹Balsaminaceae, ¹⁰Betulaceae, ¹¹Boraginaceae, ¹²Brassicaceae, ¹³Caesalpiniaceae, ¹⁴Cannabaceae, ¹⁵Caprifoliaceae, ¹⁶Caryophyllaceae, ¹⁷Clusiaceae, ¹⁸Cornaceae, ¹⁹Cyperaceae, ²⁰Equisetaceae, ²¹Ericaceae, ²²Euphorbiaceae, ²³Fabaceae, ²⁴Fagaceae, ²⁵Fumariaceae, ²⁶Geraniaceae, ²⁷Grossulariaceae, ²⁸Hippocastanaceae, ²⁹Juglandaceae, ³⁰Lamiaceae, ³¹Liliaceae, ³²Lythraceae, ³³Mimosaceae, ³⁴Moraceae, ³⁵Myricaceae, ³⁶Myrtaceae, ³⁷Onagraceae, ³⁸Papaveraceae, ³⁹Pinaceae, ⁴⁰Poaceae, ⁴¹Polygonaceae, ⁴²Portulacaceae, ⁴³Ranunculaceae, ⁴⁴Rosaceae, ⁴⁵Rutaceae, ⁴⁶Salicaceae, ⁴⁷Saxifragaceae, ⁴⁸Scrophulariaceae, ⁴⁹Solanaceae, ⁵⁰Taxaceae, ⁵¹Tiliaceae, ⁵²Vitaceae

associated with butenolols that were found in the buds (Demirci et al., 2000).

Centaurea maculosa

This study showed *C. maculosa* leaf extracts had inhibition zones of 7 to 10 mm (Table 1). A (+)-catechin enantiomer synthesized by this plant has been shown to possess antibacterial and antifungal activities while a (-)-catechin was phytotoxic (Veluri et al., 2004). Antimicrobial activity has been reported from methanol extracts of *Centaurea aintensis* and *Centaurea erengoides* flowers (Barbour et al., 2004), the essential oils of *Centaurea sessilis* and *Centaurea armena* (Yayli et al., 2005) and the dry heads of *Centaurea diffusa* (Skliar et al., 2005). Secondary metabolites including sesquiterpene lactones were identified in extracts of the aerial parts of *Centaurea deusta* and had antibacterial and antifungal activity (Karioti et al., 2002).

Epilobium angustifolium

Two separate extracts of *E. angustifolium* L. (fireweed) showed antimicrobial activity against all four microorganisms with inhibitions zones from 6 to 18 mm. Inhibition zones produced by the leaf extract ranged from 7 to 15

mm and extract from the flowering aerial parts were 7 to 20 mm (Table 2). These results agree with those from a previous study in which the antimicrobial activity of air-dried *E. angustifolium* was reported (Rauha et al., 2000). Extracts from fresh aerial parts of five *Epilobium* species including *E. angustifolium* had antimicrobial activity (Battinelli et al., 2001). *E. angustifolium* also showed good inhibitory action against *Klebsiella pneumoniae*, *P. aeruginosa* with an especially high level of activity against *Microsporum canis*; however, no inhibitory effect was recorded for *E. coli* (Battinelli et al., 2001). Methanolic extracts of *Epilobium minutum* inhibited three microorganisms including Gram-negative *K. pneumoniae*, *P. aeruginosa*, and methicillin resistant *S. aureus* but had no activity against nine fungal species (McCutcheon et al., 1992; 1994).

Hypericum perforatum

A number of studies have reported the antimicrobial activity of *Hypericum* species throughout the world, including *H. perforatum* (McCutcheon et al., 1992, 1994; Sakar and Tamer, 1990; Rabanal et al., 2002; Avato et al., 2004; Dall'Agnol et al., 2003, 2005; Barnes et al., 2001). These latter studies prepared extracts from dried aerial plant parts. Our study is the first to test an extract prepared from frozen plant material and also confirms the

Table 2. Alphabetical list of species that did not show antimicrobial activity. Generally, aerial portions of the plants were sampled including leaves, flowers, stems and fruits.

| | |
|---|---|
| <i>Acer negundo</i> L. | <i>Artemisia verlotiorum</i> LaMotte |
| <i>Achillea millefolium</i> L. | <i>Asclepias incarnata</i> L. |
| <i>Aconitum carmichealii</i> DeBeaux | <i>Asclepias syriaca</i> L. |
| <i>Agastache foeniculum</i> (Pursh) Kuntze | <i>Asclepias variegata</i> L. |
| <i>Alliaria petiolata</i> (M.Bieb.) Cavara and Grande | <i>Asclepias verticillata</i> L. |
| <i>Allium stellatum</i> Ker Gawl. | <i>Aster ericoides</i> L. |
| <i>Allium tuberosum</i> Rottl. Ex Spreng | <i>Aster umbellatus</i> Mill |
| <i>Amaranthus retroflexus</i> L. | <i>Astragalus canadensis</i> L. |
| <i>Ambrosia trifida</i> L. | <i>Astragalus mollissimus</i> Torr. |
| <i>Amorpha fruticosa</i> L. | <i>Baptisia lactea</i> (Raf.) Thieret |
| <i>Andromeda glaucophylla</i> Link | <i>Barbarea vulgaris</i> R. Br. |
| <i>Andropogon gerardii</i> Vitman | <i>Belamcanda chinensis</i> (L.) DC. |
| <i>Andropogon gerardii</i> Vitman | <i>Berteroa incana</i> (L.) DC. |
| <i>Anemone canadensis</i> L. | <i>Bidens vulgata</i> Greene |
| <i>Anemone patens</i> L. | <i>Bouteloua curtipendula</i> (Michx.) Torr. |
| <i>Anemone virginiana</i> L. | <i>Brassica</i> sp. |
| <i>Angelica</i> sp. | <i>Calendula officinalis</i> L. |
| <i>Antennaria parvifolia</i> Nutt. | <i>Calla palustris</i> L. |
| <i>Apios americana</i> Medik. | <i>Caltha palustris</i> L. |
| <i>Aquilegia canadensis</i> L. | <i>Campanula americana</i> L. |
| <i>Arctium lappa</i> L. | <i>Campanula rapunculoides</i> L. |
| <i>Artemisia absinthium</i> L. | <i>Campanula rotundifolia</i> L. |
| <i>Artemisia ludoviciana</i> Nutt. | <i>Carex</i> sp. |
| <i>Castilleja coccinea</i> (L.) Spreng | <i>Elymus trachycaulus</i> (Link) Gould ex Shinners |
| <i>Catalpa speciosa</i> Warder | <i>Equisetum arvense</i> L. |
| <i>Celtis occidentalis</i> L. | <i>Equisetum hyemale</i> L. |
| <i>Centaurea maculosa</i> Lam. | <i>Erigeron canadensis</i> L. |
| <i>Chelone glabra</i> L. | <i>Erigeron philadelphicus</i> L. |
| <i>Chenopodium album</i> L. | <i>Erigeron strigosus</i> Muhl. |
| <i>Chrysanthemum leucanthemum</i> L. | <i>Eryngium yuccifolium</i> Michx. |
| <i>Cichorium intybus</i> L. | <i>Eupatorium aromaticum</i> L. |
| <i>Cimicifuga racemosa</i> (L.) Nutt. | <i>Eupatorium purpureum</i> L. |
| <i>Cirsium arvense</i> (L.) Scop. | <i>Eupatorium rugosum</i> Houttuyn. |
| <i>Commelina communis</i> L. | <i>Euphorbia maculata</i> L. |
| <i>Convallaria majalis</i> L. | <i>Euphorbia marginata</i> Pursh |
| <i>Cornus alternifolia</i> L.f. | <i>Euthamia graminifolia</i> (L.) Nutt. |
| <i>Cornus canadensis</i> L. | <i>Euthamia tenuifolia</i> (Pursh) Nutt. |
| <i>Coronilla varia</i> L. | <i>Galeopsis tetrahit</i> L. |
| <i>Dalea candida</i> Michx. | <i>Galinsoga quadriradiata</i> Ruiz and Pav. |
| <i>Dalea purpurea</i> Vent | <i>Galium aparine</i> L. |
| <i>Daucus carota</i> L. | <i>Galium boreale</i> L. |
| <i>Dryopteris filix-mas</i> L. | <i>Gaura biennis</i> L. |
| <i>Echinacea pallida</i> Nutt. | <i>Gentiana puberulenta</i> J.S. Pringle |
| <i>Echinacea purpurea</i> (L.) Moench | <i>Glechoma hederaceae</i> L. |
| <i>Echinocystis lobata</i> (Michx.) Torr. and A. Gray | <i>Glycyrrhiza lepidota</i> Pursh |
| <i>Elymus canadensis</i> L. | <i>Grindelia squarrosa</i> (Pursh) Dunal |
| <i>Helianthus laetiflorus</i> Pers. | <i>Lithospermum canescens</i> (Michx.) Lehm. |
| <i>Helianthus maximiliani</i> Schrad. | <i>Lobelia siphilitica</i> L. |
| <i>Helianthus pauciflorus</i> Nutt. | <i>Lonicera japonica</i> Thunb. |
| <i>Helianthus tuberosus</i> L. | <i>Lonicera oblongifolia</i> (Goldie) Hook |

Table 2. contd.

| | |
|---|--|
| <p> <i>Heliopsis helianthoides</i> (L.) Sweet <i>Heracleum lanatum</i> Michx. <i>Hieracium aurantiacum</i> L. <i>Hydrangea</i> sp. <i>Hydrophyllum virginianum</i> L. <i>Hylotelephium spectabile</i> (Boreau) H. Ohba <i>Impatiens pallida</i> Nutt. <i>Inula helenium</i> L. <i>Iris</i> sp. <i>Iris virginica</i> L. <i>Juniperus communis</i> L. <i>Kuhnia eupatorioides</i> L. <i>Lactuca</i> sp. <i>Lathyrus ochroleucus</i> Hook. <i>Leonurus cardiaca</i> L. <i>Leonurus</i> sp. <i>Lepidium virginicum</i> L. <i>Liatris aspera</i> Michx. <i>Lilium michiganense</i> Farw <i>Linum rigidum</i> Pursh. <i>Osmunda cinnamomea</i> L. <i>Osmunda claytoniana</i> L. <i>Oxalis stricta</i> L. <i>Panicum</i> sp. <i>Panicum virgatum</i> L. <i>Parthenium integrifolium</i> L. <i>Pastinaca sativa</i> L. <i>Pedicularis lanceolata</i> Michx. <i>Pediomelum argophyllum</i> (Pursh) J.W. Grimes <i>Penstemon grandiflorus</i> Nutt. <i>Phlox pilosa</i> L. <i>Phragmites australis</i> (Cav.) Trin. Ex Steud. <i>Physostegia virginiana</i> (L.) Benth. <i>Pilea punula</i> Lindl., nom. Conserv. <i>Plantago major</i> L. <i>Podophyllum peltatum</i> L. <i>Polygonatum biflorum</i> (Walter) Elliott <i>Polygonum amphibium</i> L. <i>Polygonum aviculare</i> L. <i>Polygonum cilinode</i> Michx. <i>Polygonum sagittatum</i> L. <i>Prenanthes alba</i> L. <i>Prenanthes racemosa</i> Michx. <i>Solidago</i> <i>astate</i> <i>is</i> L. <i>Solidago ptarmicoides</i> (Nees) B. Boivin <i>Sonchus asper</i> (L.) Hill <i>Sorghastrum nutans</i> (L.) Nash <i>Spartina pectinata</i> Link <i>Stachys palustris</i> L. <i>Stellaria graminea</i> L. </p> | <p> <i>Lonicera prolifera</i> (Kirchn.) Rehder <i>Lotus corniculatus</i> L. <i>Lupinus perennis</i> L. <i>Lycopus americanus</i> L. <i>Lysimachia</i> <i>astate</i> L. <i>Lysimachia lanceolata</i> Walter <i>Malva neglecta</i> Wallr. <i>Matricaria matricarioides</i> (Less.) Porter <i>Medicago lupulina</i> L. <i>Medicago sativa</i> L. <i>Melilotus alba</i> Medik. <i>Melilotus officinalis</i> (L.) Pall. <i>Melothria pendula</i> L. <i>Mentha arvensis</i> L. <i>Mentha nypacalx</i> <i>Mentha</i> sp. <i>Mimulus ringens</i> L. <i>Mirabilis nyctaginea</i> (Michx.) MacMill. <i>Monarda</i> <i>astate</i> <i>i</i> L. <i>Nepeta cataria</i> L <i>Pteridium aquilinum</i> (L.) Kuhn <i>Oenothera biennis</i> L. <i>Oenothera fruticosa</i> L. <i>Ranunculus acris</i> L. <i>Ratibida pinnata</i> (Vent.) Barnhart <i>Rhamnus cathartica</i> L. <i>Rubus flagellaris</i> Willd. <i>Rudbeckia hirta</i> L. <i>Salix exigua</i> Nutt. <i>Sambucus</i> <i>astate</i> <i>is</i> L. <i>Sambucus racemosa</i> L. <i>Sanicula marilandica</i> L. <i>Saponaria officinalis</i> L. <i>Schizachyrium scoparium</i> (Michx.) Nash <i>Scrophularia lanceolata</i> Pursh <i>Scutellaria baicalensis</i> Georgi. <i>Senecio</i> sp <i>Senna hebecarpa</i> (Fernald) H.S. I and B <i>Silene vulgaris</i> (Moench) Garcke <i>Silphium lacinatum</i> L. <i>Silphium perfoliatum</i> L. <i>Silphium terebinthinaceum</i> L. <i>Solanum carolinense</i> L. <i>Typha latifolia</i> L. <i>Ulmus</i> sp <i>Urtica dioica</i> L. <i>Valeriana officinalis</i> L. <i>Verbascum thapsus</i> L. <i>Verbena astate</i> L. <i>Verbena stricta</i> Vent. </p> |
|---|--|

Table 2. contd.

| | |
|---|---|
| <i>Stylophorum diphyllum</i> (Michx.) Nutt. <i>Symphotrichum ciliolatum</i> (Lindl.) Love <i>Syringa vulgaris</i> L. <i>Tagetes minuta</i> L. <i>Taraxacum officinale</i> F.H. Wigg. <i>Teucrium canadense</i> L. <i>Thalictrum pubescens</i> Pursh <i>Thaspium trifoliatum</i> (L.) A. Gray <i>Thermopsis villosa</i> (Walter) F. and B.G. Schub <i>Thlaspi arvense</i> L. <i>Tradescantia virginiana</i> L. <i>Tragopogon dubius</i> Scop. <i>Trifolium ambiguum</i> Bieb. <i>Trifolium arvense</i> L. <i>Trifolium pratense</i> L. <i>Trifolium repens</i> L. | <i>Verbena urticifolia</i> L. <i>Vicia cracca</i> L. <i>Viola canadensis</i> L. <i>Zanthoxylum americanum</i> Mill. <i>Zizia aptera</i> (A. Gray) Fernald |
|---|---|

antimicrobial activity of *H. perforatum* (Table 1). The degree of antimicrobial activity seems to be affected by the date of collection. Samples harvested in July of 2003 and 2004 did not display antimicrobial activity, whereas, samples collected in August were active. Flowering aerial parts collected earlier in 2004 inhibited our four test microorganisms. However, the aerial parts including immature seed heads collected later in 2004 only inhibited *S. aureus*. Seasonal variations of phyto-chemical production probably occur in this species. Multiple chemical constituents are reported to contribute to the bioactivity of *Hypericum* spp. Hyperforin, a phloroglucinol present in *H. perforatum* and some other species of *Hypericum*, is the primary component responsible for the antimicrobial activity (Avato et al., 2004; Dall'Agnol et al., 2003, 2005; Barnes et al., 2001). Hypericin has also been identified as the active component responsible for killing avian influenza virus H5N1 *in vitro* (Wang et al., 2006).

Lythrum salicaria

Extracts of the flowering spikes and leaves of *L. salicaria* L., (purple loosestrife) showed antimicrobial activity against all four of the test microorganisms (Table 1). This observation is supported by additional reports on the antimicrobial activity of *L. salicaria* extracts against *S. aureus*, *E. coli*, *C. albicans*, *Bacillus cereus*, *Mycobacterium smegmatis* and *Micrococcus luteus* (Rauha et al., 2000; Dulger and Gonuz, 2004). Although *E. coli* and *C. albicans* were included in the study of Dulger and Gonuz (2004), inhibition of these microorganisms was not reported.

Rhus glabra* and *R. typhina

Two separate extracts of *Rhus glabra* L. leaves and

green flower clusters exhibited antimicrobial activity (Table 1). Inhibition zones of the leaf extract ranged from 7 to 15 mm and the extracts of the green flower clusters from 10 to 16 mm. Previous studies reported antibacterial and antifungal activity of extracts prepared from air-dried branches of *R. glabra* (McCutcheon et al., 1992, 1994). *R. glabra* branches showed the broadest spectrum of antimicrobial activity inhibiting 11 microorganisms including four Gram-positive and seven Gram-negative (McCutcheon et al., 1992). Antimicrobial activity from dehydrated unripened and ripened fruits of *Rhus coriaria* L. has also been observed (Nasar-Abbas and Kadir Halkman, 2004).

Plant extracts inhibiting three microorganisms

Desmodium illinoense

Although many *Desmodium* species are used in ethnomedicine, this study is the first report of antimicrobial activity from the genus. *Desmodium illinoense* A. Gray (prairie tick trefoil) showed antimicrobial activity against *S. aureus*, *P. aeruginosa* and *C. albicans* (Table 1). Other species in this genus have been used as an anti-inflammatory, anticatarrhal, and anti-nociceptive (Rathi et al., 2004), analgesic and anticonvulsion (N'gouemo et al., 1996), and antileishmanial (Mishra et al., 2005; Singh et al., 2005).

Scirpus americanus

This study is the first to report antimicrobial activity of a *Scirpus* species (American bulrush). Extracts of *Scirpus americanus* Pers. inhibited the three microorganisms, *S. aureus*, *E. coli* and *P. aeruginosa* (Table 1). We have uncovered no reports on the use of *Scirpus* sp. as an an-

timicrobial in North American.

Plant extracts inhibiting two microorganisms

Eleven extracts from 10 plant species inhibited *S. aureus* and *C. albicans* (Table 1). Those having inhibitory activity were extracts from the leaves of *Clintonia* sp., aerial organs of *Comptonia peregrina* (L.) J.M. Coult., the leaves of *C. peregrina*, flowering organs of *Desmanthus illinoensis* (Michx.) MacM. (Illinois bundle flower), leaves of *Epilobium ciliatum* Raf. (American willow herb), immature flower clusters *Eupatorium maculatum* L. (Joe pye weed), immature floral organs of *Geum virginianum* L. (rough avens), leaves of *Juglans nigra* L. (black walnut) leaves and flowers of *Polygonum coccineum* L. *Cotinus coggygria* Scop. (smoke tree) inhibited *S. aureus* and *P. aeruginosa* but had no activity against *C. albicans*.

Clintonia sp.

This study is the first report of antibacterial activity in a *Clintonia* sp. An extract of *Clintonia* sp. leaves and roots inhibited *S. aureus* and *C. albicans* (Table 1). Fresh leaves of *Clintonia borealis* (Ait.) Raf. (blue bead lily) extracted with ethanol were reported to have antifungal activity against *M. gypseum* and *Trichophyton mentagrophytes* (Jones et al., 2000). Hence, it may not be surprising that we observed inhibitory activity against *C. albicans*.

Comptonia peregrina

Comptonia peregrina (L.) J.M. Coult. (sweet ferns) are aromatic and secrete resin from numerous capitate-stalked glands on the leaves, especially on the lower surface. Our study is the first to report antimicrobial activity from *C. peregrina*. Two separate extracts of *C. peregrina* aerial parts and leaves inhibited *S. aureus* and *C. albicans* with the respective inhibitions zones of 9/15 and 14/12 mm (Table 1). Twenty-seven compounds have been identified in the essential oil of sweet ferns with the following probably accounting for the observed antimicrobial activity of extracts and include cineol, gamma terpinene, and caryophyllene in the highest concentrations (Halim and Collins, 1973; Lawrence and Weaver, 1974).

Cotinus coggygria

Cotinus coggygria Scop. (smoke tree) is a non-native species to the mid-western United States but is commonly grown as a garden shrub. It is closely related to the genus *Rhus* (sumacs). Leaf extracts of *C. coggygria* showed inhibition zones of 13 and 10 mm against *S. aureus* and *P. aeruginosa* (Table 1). In Bulgaria, the leaves are widely used in folk medicine for gastric ulcers, antidiarrhetic, anti-inflammatory, paradontosis (Ivanova et

al., 2005). In some countries extracts are used as antiseptics, antimicrobials, antihemorrhagics and as an aid in wound healing (Demirci et al., 2003; Tzakou et al., 2005). Essential oils of *C. coggygria* leaves, inflorescences and infructescences have been found to be high in monoterpenes such as limonene, myrcene, sabinene and alpha-pinene (Demirci et al., 2003; Tzakou et al., 2005). Medicinal benefits of monoterpenes, especially limonene, are being studied for cancer prevention and treatment while monoterpenols possess antibacterial and antifungal properties (Bowles, 2003).

Desmanthus illinoensis

Desmanthus illinoensis (Michx.) MacMill. (Illinois bundle flower) extracts from flowering aerial parts produced inhibitions zones of 15 and 17 mm against *S. aureus* and *C. albicans* respectively (Table 1). Quercetin, myricitrin and gallic acid esters of myricitrin have been isolated from bundle flower and shown to have antibacterial activity against *Bacillus sphaericus*, *Bacillus thuriengensis*, *Bacillus subtilis* and *Pseudomonas mallophilia* (Nicollier and Thompson, 1983).

Eupatorium maculatum

The flower clusters of *E. maculatum* L. (Joe pye weed) yielded an extract with slight antimicrobial against *C. albicans* and a 9 mm inhibition zone against *S. aureus* (Table 1). *Eupatorium salvia* Colla from Chile is used as an antiseptic/antimicrobial for infected wounds and insect bites (Urzua et al., 1998) and *Eupatorium glutinosum* from Ecuador and Peru is used as an astringent, antirheumatic, and antimicrobial (El-Seedi et al., 2002). The antimicrobial activity of *E. salvia* has been attributed to the presence of diterpenoids (Urzua et al., 1998). Aerial parts of *Eupatorium aschenbornianum* (Rios et al., 2003) and leaves and twigs of *E. glutinosum* (El-Seedi et al., 2002) had antimicrobial activity that was again mainly attributed to the presence of diterpenoids. Our study is the first to report antimicrobial activity of extracts from flower clusters.

Geum sp.

Extracts from *Geum virginianum* L. (rough avens) had antimicrobial activity with inhibition zones of 10 mm against *S. aureus* and *Candida albicans* (Table 1). This is the first report on the antimicrobial activity of this species. However, extracts of other *Geum* species have been reported to have antimicrobial activity. These include *Geum macrophyllum* Willd. var. *macrophyllum* (McCutcheon et al., 1994) and *Geum rivale* L. (Panizzi et al., 2000). Extracts of *Geum macrophyllum* roots showed antifungal activity against nine fungal species although five had incomplete inhibition (McCutcheon et al., 1994). Extracts of flowering aerial parts of *G. rivale* L. had anti-

antimicrobial activity against Gram-positive, Gram-negative and mycetous microorganisms. The crude methanol extract had the most antimicrobial activity (Panizzi et al., 2000).

Juglans nigra

Leaf extracts from *Juglans nigra* L. had antimicrobial activity against *S. aureus* and *C. albicans* (Table 1), supporting similar findings for other *Juglans* species (Omar et al., 2000; Clark et al., 1990; Grujić-vasić et al., 1990; Alkhawajah, 1997; Cha et al., 1998; Lopez et al., 2001; Qa'dan et al., 2005).

As in our study Grujić-vasić et al. (1990) reported antibacterial activity by *J. nigra* against seven microorganisms including the four tested in this study. In contrast, our studies found no activity against either *E. coli* or *P. aeruginosa*. The lack of activity against these two organisms could be associated with the time of collection or the age of tested plants.

One of the compounds believed to contribute to the biological activities of various species of *Juglans* is juglone, a naphthoquinone, which is found in all plant organs in most members of the Juglandaceae (Clark et al., 1990). Juglone (5-hydroxy-1,4-naphthoquinone) was reported to have good activity against the test yeasts and fungi including *C. albicans*, *Saccharomyces cerevisiae*, *Cryptococcus neoformans*, *Aspergillus flavus* and *Aspergillus fumigatus* but having only moderate activity against bacteria indicating that additional phytochemicals contribute to the antimicrobial activity of *Juglans* spp.

Polygonum coccineum

Extracts from many species of *Polygonum* including knotweed and smartweed, have been found to possess antimicrobial activity (Mackeen et al., 1997; Penna et al., 2001; Kumagai et al., 2005). We found flower extract of *Polygonum coccineum* (swamp smartweed) to have antimicrobial activity against *S. aureus* and *C. albicans* (Table 1). Other studies have reported extracts of the leaves, aerial portions, leaves, rhizomes and whole plants of *Polygonum minus* Huds. (Mackeenre, 1997), *Polygonum punctatum* Elliot (Lopez et al., 2001), *P. punctatum* var. *aquatile* (Martins) [(Penna et al., 2001)] and *Polygonum sachalinense* F. Schmidt ex Maxim (Kumagai, 2005) respectively, to have antimicrobial activity. *P. minus* leaf ethanolic extracts have shown antimicrobial activity against *P. aeruginosa* and had suppressive activity on a human cervical carcinoma cell-line (Mackeen et al., 1997). *P. punctatum* methanol extracts of aerial parts exhibited antiviral and antibacterial activity (Lopez et al., 2001). Dichloromethane extracts of *P. punctatum* var. *aquatile* leaves and rhizomes inhibited five microorganisms where methanol extracts inhibited one microorganism and none were inhibited by ethanol or aqueous extracts (Penna et al., 2001). Extracts of whole

plants of *P. sachalinense* have been reported to show antifungal and antimicrobial activity with special interest in its activity against the fish pathogen *Photobacterium damsela* subsp. *piscicida* (Kumagai et al., 2005).

Plant extracts inhibiting a single microorganism

There were 119 plant extracts, representing 98 species having antimicrobial activity against one microorganism. *S. aureus* was inhibited by 108 extracts and *C. albicans* by 11. Several extracts had large inhibition zones over 15 mm and are noted below.

Anaphalis margaritacea

Anaphalis margaritacea L. (Benth and Hook) [pearly everlasting] leaf extract inhibited *S. aureus* with an inhibition zone of 18 mm. Chemical investigations of the flowering aerial parts of *A. margaritacea* have identified flavonoids (Wollenweber et al., 1993), and diterpenes and hydroxylactones as active constituents with two diterpenes having antibacterial activity against *B. cereus*, *P. aeruginosa* and *E. coli* (Ahmed et al., 2004).

Cannabis sativa

Cannabis sativa L. extracts had very good antimicrobial activity against only *S. aureus* with an inhibition zone of 25 mm. This inhibitory zone was nearly equivalent to the controls. Wasim et al. (1995) reported antimicrobial activity from ethanol and petroleum ether extracts against multiple microorganisms. We found antimicrobial activity against only one, a Gram-positive cocci. Essential oils extracted from five fiber varieties of *C. sativa* had antimicrobial activity with the degree of antimicrobial activity varying between cultivars (Novak et al., 2001). The main components of the essential oils reported were alpha-pinene, myrcene, *trans*-beta-ocimene, alpha-terpinolene, *trans*-caryophyllene and alpha-humulene. Alpha-terpinolene was the component that varied the most between cultivars (Novak et al., 2001). Cannabidiol (CBD) has also been identified as a component of hemp oil effective against Gram-positive bacteria and yeast (Leizer et al., 2000). A strong correlation exists between the antimicrobial activity and the level of cannabidiolic acid (CBD) found in this species (Leizer et al., 2000). The *C. sativa* chemotypes grown in northern latitudes are reported to have a higher ratio of CBD to Δ -9-tetrahydro-cannabinol (THC) resulting in stronger antimicrobial activity (Leizer et al., 2000).

Potentilla arguta

Potentilla arguta Pursh leaf extracts had antimicrobial activity against *C. albicans* with an inhibition zone of 16 mm. Other studies have reported antibacterial activity of *P. arguta* as well as antifungal and antiviral properties

(McCutcheon et al., 1992, 1994, 1995). McCutcheon et al. (1992, 1994, 1995) reported that methanol root extracts of *P. arguta* inhibited nine bacterial species, four fungal species and the bovine respiratory syncytial virus, *Paramyxoviridae*.

Smilacina racemosa

The extract from the mature fruits of *Smilacina racemosa* (L.) Desf. (false Solomon's seal) exhibited a 17 mm inhibition zone against *C. albicans*. Methanolic extracts of the rhizomes of *S. racemosa* were reported to have no antibacterial, antifungal or antiviral activity (McCutcheon et al. 1992, 1994, 1995). Our study is the first to report antimicrobial activity of the berries.

Summary

Our study has identified 142 plant extracts from 109 species that have significant antimicrobial activity. The effectiveness of antimicrobial activity could be viewed as significant based on either the number of microorganisms inhibited or the intensity of antimicrobial action based on the size of the zone of inhibition. Twenty-four percent of all the plants investigated had activity against at least one test microorganism. Various extracts from *B. papyrifera*, *C. maculosa*, *E. angustifolium*, *H. perforatum*, *L. salicaria* and *R. glabra* inhibited the growth of all microorganisms used in this study. Two plant species *D. illinoense* and *S. americanus* inhibited three microorganisms, eleven extracts (10 species) inhibited two microorganisms while 119 extracts (98 species) inhibited at least one microorganism.

Results of these studies indicate that further searches and characterizations of plants for antimicrobial compounds are warranted. In addition, research on synergistic combinations of extracts with broad spectrum or a high degree of inhibition against a particular microorganism would seem worthwhile.

As the search for new antimicrobial agents intensifies, plant extracts may provide attractive alternate sources of molecules for consideration. As drug resistance becomes an increasing problem and as consumer demand for products with natural preservative grows, perhaps it is these molecules that may form the basis of future antimicrobial research efforts.

REFERENCES

- Ahmed AA, Hussein TA, Mahmoud AA, Farag MA, Paré PW, Wojcińska M, Karchesy J, Mabry TJ (2004). Nor-ent-kaurane diterpenes and hydroxylactones from *Antennaria geyeri* and *Anaphalis margaritacea*. *Phytochemistry* 65: 2539-2543.
- Alkhawajah AM (1997). Studies on the antimicrobial activity of *Juglans regia*. *Am. J. Chin. Med.* XXV: 175-180.
- Avato P, Raffo F, Guglielmi G, Vitali C, Rosato A (2004). Extracts from St. John's Wort and their antimicrobial activity. *Phytother. Res.* 18: 230-232.
- Barbour EK, Sharif MA, Sagherian VK, Habre AN, Talhouk RS, Talhouk SN (2004). Screening of selected indigenous plants of Lebanon for antimicrobial activity. *J. Ethnopharmacol.* 93: 1-7.
- Barnes J, Anderson LA, Phillipson JD (2001). St. John's Wort (*Hypericum perforatum* L.): A review of its chemistry, pharmacology and clinical properties. *J. Pharm. Pharmacol.* 53: 583-600.
- Battinelli L, Tita B, Evandri MG, Mazzanti G (2001). Antimicrobial activity of *Epilobium* spp. extracts. *Farmaco* 56: 345-348.
- Bauer, AW, Kirby MM, Sherris JC, Turck M (1966). Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Pathol.* 45: 493-496.
- Beuchat LR, Brackett RW, Doyle MP (1994). Antimicrobials occurring naturally in foods. *Food Technol.* 43: 134-142.
- Bowles EJ (2003). *The Chemistry of Aromatherapeutic Oils*. 3rd ed. Griffin Press, South Australia.
- Cha BC, Lee HW, Choi MY (1998). Antioxidative and antimicrobial effects of nut species. *Korean J. Pharmacogn.* 29: 28-34.
- Clark AM, Jurgens TM, Hufford CD (1990). Antimicrobial activity of Juglone. *Phytother. Res.* 4: 11-14.
- Cushman KE, Maqbool M, Gerard PD (2005). Mulch type, mulch depth and rhizome planting depth for field-grown American may-apple. *HortScience* 40: 635-639.
- Cutter C (2000). Antimicrobial effect of herb extracts against *Escherichia coli* O157:H7, *Listeria monocytogenes* and *Salmonella typhimurium* associated with beef. *J. Food Prot.* 63: 601-607.
- Dall'Agnol R, Ferraz A, Bernardi AP, Albring D, Nör C, Sarmiento L, Lamb L, Hass M, von Poser G, Schapoval EES (2003). Antimicrobial activity of some *Hypericum* species. *Phytomedicine* 10: 511-516.
- Dall'Agnol R, Ferraz A, Bernardi AP, Albring D, Nör C, Schapoval EES, von Poser GL (2005). Bioassay-guided isolation of antimicrobial benzopyrans and phloroglucinol derivatives from *Hypericum* species. *Phytother. Res.* 19: 291-293.
- Delaquis PQ, Mazza G (1995). Antimicrobial properties of isothiocyanates in food preservation. *Food Technol.* 49: 73-78.
- Demirci B, Hüsnü Can Başer K, Özek T, Demirci F (2000). Betulenols from *Betula* species. *Planta Med.* 66: 490-493.
- Demirci B, Demirci F, Baser KHC (2003). Composition of the essential oil of *Cotinus coggryia* Scop. from Turkey. *Flavour Fragrance J.* 18: 43-44.
- Dulger B, Gonuz A (2004). Antimicrobial activity of certain plants used in Turkish traditional medicine. *Asian J. Plant Sci.* 3: 104-107.
- El-Seedi HR, Ohara T, Sata N, Nishiyama S (2002). Antimicrobial diterpenoids from *Eupatorium glutinosum* (Asteraceae). *J. Ethnopharmacol.* 81: 293-296.
- Grujić-vasić J, Bosnić T, Tockić S, Bašić F (1990). Relative astringency and antimicrobial activity of the willow herb and walnut tree leaves. *Period. Biol.* 92: 479-480.
- Halim AF, Collins RP (1973). Essential oil analysis of the Myricaceae of the Eastern United States. *Phytochemistry* 12:1077-1083.
- Hao YY, Brackett RE, Doyle MP (1998). Efficacy of plant extracts in inhibiting *Aeromonas hydrophilia* and *Listeria monocytogenes* in refrigerated cooked poultry. *Food Microbiol.* 15: 367-378.
- Ivanova D, Gerova D, Chervenkov T, Yankova T (2005). Polyphenols and antioxidant capacity of Bulgarian medicinal plants. *J. Ethnopharmacol.* 96: 145-150.
- Jones NP, Arnason JT, Abou-Zaid M, Akpagana K, Sanchez-Vindas P, Smith JL (2000). Antifungal activity of extracts from medicinal plants used by First Nations Peoples of eastern Canada. *J. Ethnopharmacol.* 73:191-198.
- Karioti A, Skaltsa H, Lazari D, Sokovic M, Garcia, Harvala C (2002). Secondary metabolites from *Centaurea deusta* with antimicrobial activity. *Z Naturforsch.* 57c: 75-80.
- Kumagai H, Kawai Y, Sawano R, Kurihara H, Yamazaki K, Inoue N (2005). Antimicrobial substances from rhizomes of the giant knotweed *Polygonum sachalinense* against the fish pathogen *Photobacterium damsela* subsp. *piscicida*. *Z Naturforsch* 60: 39-44.
- Lawrence BM, Weaver KM (1974). Essential oils and their constituents. *Planta Med.* 25: 385-388.
- Leizer C, Ribnický D, Poulev A, Dushenkov S, Raskin I (2000). The composition of hemp seed oil and its potential as an important source of nutrition. *J. Nutraceuticals. Funct. Med. Foods* 2: 35-53.
- Lis-Balchin M, Deans SG (1997). Bioactivity of selected plant essential oils against *Listeria monocytogenes*. *J. Appl. Microbiol.* 82: 759-762.

- Lopez A, Hudson JB, Towers GHN (2001). Antiviral and antimicrobial activities of Colombian medicinal plants. *J. Ethnopharmacol.* 77: 189-196.
- Mackeen MM, Ali AM, El-Sharkawy SH, Manap MY, Salleh KM, Lajis NH, Kawazu K (1997). Antimicrobial and cytotoxic properties of some Malaysian traditional vegetables (Ulam). *Int. J. Pharmacogn.* 35: 174-178.
- McCutcheon AR, Ellis SM, Hancock REW, Towers GHN (1992). Antibiotic screening of medicinal plants of the British Columbian native peoples. *J. Ethnopharmacol.* 44: 157-169.
- McCutcheon AR, Ellis SM, Hancock REW, Towers GHN (1994). Antifungal screening of medicinal plants of British Columbian native peoples. *J. Ethnopharmacol.* 44: 157-169.
- McCutcheon AR, Roberts TE, Gibbons E, Ellis SM, Babiuk LA, Hancock REW, Towers GHN (1995). Antiviral screening of British Columbian medicinal plants. *J. Ethnopharmacol.* 49: 101-110.
- Mishra PK, Singh N, Ahmad G, Dube A, Maurya R (2005). Glycolipids and other constituents from *Desmodium gangeticum* with antileishmanial and immunomodulatory activities. *Bioorg. Med. Chem. Lett.* 15: 4543-4546.
- *Moerman DE (2004). *Native American Ethnobotany*. Timber Press, Inc., Portland. pp. 927
- Nasar-Abbas SM, Halkman AK (2004). Antimicrobial effect of water extract of sumac (*Rhus coriaria* L.) on the growth of some food borne bacteria including pathogens. *Int. J. Food Microbiol.* 97: 63-69.
- N'gouemo P, Baldy-Moulinier M, Nguemby-Bina C (1996). Effects of an ethanolic extract of *Desmodium adscendens* on central nervous system in rodents. *J. Ethnopharmacol.* 52: 77-83.
- Nicollier G, Thompson AC (1983). Flavonoids of *Desmanthus illinoensis*. *J. Nat. Prod.* 46: 112-117.
- Novak J, Zitterl-Eglseer K, Deans SG, Franz CM (2001). Essentials oils of different cultivars of *Cannabis sativa* L. and their antimicrobial activity. *Flavour Fragr. J.* 16: 259-262.
- Omar S, Lemonnier B, Jones N, Ficker C, Smith ML, Neema C, Towers GHN, Goel K, Arnason JT (2000). Antimicrobial activity of extracts of eastern North American hardwood trees and relation to traditional medicine. *J. Ethnopharmacol.* 73: 161-170.
- Panizzi L, Catalano S, Miarelli C, Cioni PL, Campeol E (2000). In vitro antimicrobial activity of extracts and isolated constituents of *Geum rivale*. *Phytother. Res.* 14: 561-563.
- Penna C, Marino S, Vivot E, Cruaños MC, Muñoz J, De D, Cruaños J, Ferraro G, Gutkind G, Martino V (2001). Antimicrobial activity of Argentine plants used in the treatment of infectious diseases: Isolation of active compounds from *Sebastiania brasiliensis*. *J. Ethnopharmacol.* 77: 37-40.
- Puupponen-Pimia R, Nohynek L, Meier C, Kahkonen M, Heinonen M, Hopia A, Oksman-Caldentey K-M (2001). Antimicrobial properties of phenolic compounds from berries. *J. Appl. Microbiol.* 90: 494-507.
- Qa'dan F, Thewaini A-J, Ali DA, Afifi R, Elkhawad A, Matalka KZ (2005). The antimicrobial activities of *Psidium guajava* and *Juglans regia* leaf extracts to acne-developing organisms. *Am. J. Chin. Med.* 33: 197-204.
- Rabanal RM, Arias A, Prado B, Hernandez-Perez M, Sanchez-Mateo CC (2002). Antimicrobial studies on three species of *Hypericum* from the Canary Islands. *J. Ethnopharmacol.* 81: 287-292.
- Rathi A, Rao CV, Ravishankar B, De S, Mehrotra S (2004). Anti-inflammatory and anti-nociceptive activity of the water decoction of *Desmodium gangeticum*. *J. Ethnopharmacol.* 95: 259-263.
- Rauha JP, Remes S, Heinonen M, Hopia A, Kähkönen M, Kujala T, Pihlaja K, Vuorela H, Vuorela P (2000). Antimicrobial effects of Finnish plant extracts containing flavonoids and other phenolic compounds. *Int. J. Food Microbiol.* 56: 3-12.
- Rios MY, Aguilar-Guadarrama B, Navarro V (2003). Two new benzofuranes from *Eupatorium aschenbornianum* and their antimicrobial activity. *Planta Med.* 69: 967-970.
- Sakar MK, Tamer AU (1990). Antimicrobial activity of different extracts from some *Hypericum* species. *Fitoterapia* 61: 464-466.
- Singh N, Mishra PK, Kapil A, Arya KR, Maurya R, Dube A (2005). Efficacy of *Desmodium gangeticum* extract and its fractions against experimental visceral leishmaniasis. *J. Ethnopharmacol.* 98: 83-88.
- Skliar, MI, Toribio MS, Oriani DS (2005). Antimicrobial activity of *Centaurea diffusa*. *Fitoterapia* 76: 737-739.
- Small E, Catling PM (1999). *Canadian Medicinal Crops*. NRC Research Press, Ottawa, Ontario Canada. pp 240
- Somaatmadja D, Powers JJ, Handy MK (1964). Anthocyanins.VI. Chelation studies on anthocyanins and other related compounds. *J. Food Prot.* 56: 406-409.
- Tzakou O, Bazos I, Yannitsaros A (2005). Essential oils of leaves, inflorescences and infructescences of spontaneous *Cotinus coggyria* Scop. from Greece. *Flavour Fragrance J.* 20: 531-533.
- Thompson DQ, Stuckey RL, Thompson EB (1987). Spread, impact and control of purple loosestrife (*Lythrum salicaria*) in North American wetlands. *Fish and Wildlife Research 2*, US Department of the Interior, Washington, DC.
- Urzua A, Caroli M, Vasquez L, Mendoza L, Wilkens M, Tojo E (1998). Antimicrobial study of the resinous exudate and of diterpenoids isolated from *Eupatorium salvia* (Asteraceae). *J. Ethnopharmacol.* 62: 251-254.
- Veluri R, Weir TL, Bais HP, Stermitz FR, Vivanco JM (2004). Phytotoxic and antimicrobial activities of catechin derivatives. *J. Agric. Food Chem.* 52: 1077-1082.
- Wani MC, Taylor HL, Wall ME, Coggon P, McPail AT (1971). Plant antitumor agents. Part VI. The isolation and structure of Taxol, a novel antileukemic and antitumor agent from *Taxus brevifolia*. *J. Am. Chem. Soc.* 93: 2325-2327.
- Wasim K, Haq IU, Ashraf M (1995). Antimicrobial studies of the leaf of *Cannabis sativa* L. *Pak. J. Pharm. Sci.* 8: 22-38.
- Wijesekera RB (1991). *Plant derived medicines and their role in global health*. The Medicinal Plant Industry, CRC Press.
- Wollenweber E, Fritz H, Henrich B (1993). Rare flavonoid aglycones from *Anaphalis margaritacea* and two *Gnaphalium* species. *Z Naturforsch* 48: 420-424.
- Yayli N, Yasar C, Gulec A, Usta S, Kolayli K, Coskuncelebi K, Karaoglu S (2005). Composition and antimicrobial activity of essential oils from *Centaurea sessilis* and *Centaurea armena*. *Phytochemistry* 66: 1741-1745.
- Zink DL (1997). The impact of consumer demands and trends on food processing. *Emerging Infect. Dis.* 3: 467-469.