

Full Length Research Paper

Effect of pre-treatment on isolated microspores culture ability in durum wheat (*Triticum turgidum* subsp. *durum* Desf.)

O. Slama Ayed^{1*}, J. De Buyser², E. Picard², Y. Trifa¹ and H. Slim Amara¹

¹Laboratoire d'amélioration génétique des Céréales, Institut National Agronomique de Tunisie, 43 Avenue Charles Nicolle 1082 -Tunis- Mahrajène Tunisia.

²Laboratoire de Morphogenèse Végétale Expérimentale Haploïde, Bat. 360, Université Paris sud XI Orsay- 91405 Cedex, France.

Accepted 11 December, 2009

The production of double haploids through androgenesis is used by breeders to produce homozygous lines in a single generation. Androgenesis can be achieved by isolated microspore culture, which, however, allows the production of green haploids with a very low efficiency. In order to improve the overall green haploid regeneration in durum wheat, we compare the efficiency of eight different pretreatments. These pretreatments were: cold pre-treatment for 5 weeks, 0.3M mannitol for 12 days at 4°C; 0.3M mannitol for 7 days at 4°C; 0.7M mannitol for 5 days at 4°C; PEG₄₀₀₀ 1.5% for 5 days at 4°C; PEG₄₀₀₀ 1% for 15 days at 4°C; PEG₄₀₀₀ 1% for 10 days at 4°C and control without pretreatment. The cold treatment for 5 weeks (4°C) is the most efficient and was used to test the androgenetic ability by microspore culture method of three Tunisian durum wheat cultivars (Razzek, Khiar and Azizi). This pre-treatment improves significantly the embryogenesis induction and green plants regeneration with green/albina ratio ≈ 2 for the cultivar "Khlar". The PEG₄₀₀₀ 1% for 10 days seems promising pre-treatment and this is important to perform new experiments to advance their response.

Key words: Isolated microspore culture, double haploid, durum wheat, pre-treatment.

INTRODUCTION

Double Haploid lines are commonly used in plant breeding programs for the production of homozygous lines in a single generation. Androgenesis is one of the most efficient methods for doubled haploid production and can be performed using anther or isolated microspores culture. In bread wheat (*Triticum aestivum* L.) and other cereals, androgenesis was used to develop number of cultivars that are widely cultivated in many countries (Cistué et al., 2003). However, durum wheat (*Triticum turgidum* subsp. *Durum* Desf.) remain recalci-

trant to this technique with low regeneration rate and very high frequency of albino plants (Jaiti et al., 2000; Labbani et al., 2005; Cistué et al., 2006, 2009). Albino plants can even reach in many cases up to 100% of the regenerated plants (Labbani et al., 2005). Thus, optimization of isolated microspore protocol aiming at maximizing the production of green plants of durum wheat remains of considerable interest. Many physiological factors can influence the microspores culture response such as growth conditions of donor plants, developmental stage of microspores, types of pre-treatments and composition of induction and regeneration media (Touraev et al., 1996; Hu and Kasha, 1997; Liu et al., 2002; Zheng et al., 2003; Cistué et al., 2009). Among them, pre-treatment is the most important factor. The stress treatment allows the conversion of microspores into embryos by diverting microspores from their normal gametophytic pathway into sporophytic development (Touraev et al., 1996; Hu and Kasha, 1999). It also plays an important role in the

*Corresponding author. E-mail: olfa.slama@planet.tn, olfayed@yahoo.fr.

development of the embryos into green seedlings (Shariatpanahi et al., 2006) and consequently improves the yield of green plants regeneration (Kasha et al., 2001; Li and Devaux, 2003). So far, in order to enhance durum wheat microspores reorientation and improve green plants regeneration, various stress pretreatments were applied including cold (Labbani et al., 2005), osmotic shock using mannitol (Cistué et al., 2006; Labbani et al., 2007). It was shown by Labbani et al. (2005) that a long term (five weeks) at 4°C cold pre-treatment of the microspores could be promising for green regeneration in durum wheat. Cistué et al. (2006) reported that anthers pre-treated in 0.7M mannitol for 5 days improved green plants regeneration. Labbani et al. (2007) showed that a pre-treatment consisting in a combined mannitol 0.3 M and cold for 7 days had a strong effect on the number of embryos produced and the regeneration of green plants. In our study, we have tested these different types of stress to compare their efficiency. In order to try to further improve the rate of haploid production through durum wheat microspores culture, we tested also the efficiency of a novel combined pre-treatment cold and PEG for different duration. In solution, PEG is used to induce an osmotic stress. This pre-treatment creates more negative water potential, facilitating a water-restricted environment (Ilić-Grubor et al., 1998). Finally, using the best pre-treatment identified, we have produced haploid plants from three different durum wheat genotypes by isolated microspores culture.

MATERIALS AND METHODS

The experiments presented in this article were performed in Laboratory of Morphogenesis Haploid Vegetable Experimental at Paris XI University-Orsay, France.

Donor plants and growth conditions

Three Tunisian durum wheat cultivars were used as donor plants in this study: Azizi, Khiar and Razzek. Plants were cultivated in pots and grown in a controlled greenhouse, regulated with a photoperiod of 16 h light/ 8 h dark. Light intensity was set at 1150 $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$. Temperature was set at 22°C during the day and 15°C in the night. Humidity was kept at 70%.

Spike pre-treatments

Spikes from Razzek variety were collected when most of the microspores were at the late uninucleate stage and stored in the different pre-treatment solutions in the dark at 4°C.

Spikes were subjected to pretreatments before using them for isolated microspores culture. Pre-treatment were based on cold, cold mannitol and cold PEG₄₀₀₀ treatments. In solution, mannitol and PEG were used to induce stress.

A total of seven pretreatments and control without pre-treatment were used:

- (i) No pre-treatment.
- (ii) Cold pre-treatment for 5 weeks.

- (iii) 0.3 M mannitol for 12 days.
- (iv) 0.3M mannitol for 7 days.
- (v) 0.7M mannitol for 5 days.
- (vi) PEG₄₀₀₀ 1.5% for 5 days.
- (vii) PEG₄₀₀₀ 1% for 10 days.
- (viii) PEG₄₀₀₀ 1% for 15 days.

Microspore isolation

Microspores were isolated and cultivated according to the protocol described by De Buyser et al. (2002). After pre-treatment, the spikes were sterilized with calcium hypochlorite 5% during 10 min followed by three washings with sterile water. The extracted microspores were cultivated in CHB3 medium based on Chu et al. (1990) modified by adding 90 g/l of maltose (Table 1). Microspores were adjusted at density of 50 000 microspores ml^{-1} and were cultivated in 35 × 15 mm Petri dishes. Immature ovaries were added to the culture at density of four per milliliter, immediately preceding the incubation. The inclusion of ovaries in these culture systems seemed to enhance embryogenesis and/or improve the quality of the embryos (Zheng et al., 2002). Plates were sealed and incubated in the dark at 27°C.

Embryo induction and plant regeneration

After embryos were generated and had reached a size of 1 to 2 mm of diameter for about 4 weeks, they were aseptically transferred to solid MS medium (Murashige and Skoog, 1962) without growth regulators at a density of 30 embryos per Petri dish. The dishes were incubated in a growth room at 25°C with 16 h light /8 h dark photoperiod, at a light intensity of 80 - 100 $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$. Approximately 2 weeks after embryos transfer, the numbers of regenerated plantlets (green and albino) were counted.

Data collection

Data were recorded as numbers and as percentages. All percentages were calculated based upon the total number of cultured microspores. Number of induced microspores, dividing cells and proembryos were counted under an inverted microscope.

All data were based on a minimum of 3 replications per experiment for each particular treatment or genotype. Each experiment contains only one variable.

The data were statistically analysed using ANOVA conducted using SAS computer software (1988). Mean comparisons were made with the student test. A difference was considered significant at $P=0.05$.

RESULTS

Spikes were collected when microspores were at the late uninucleate stage (Figure 1a). Subsequently, tillers containing spikes at the desired development stage could be pre-selected on the basis of their morphology (Figure 1b).

Microspore embryogenesis and haploid plant regeneration

In freshly isolated durum wheat microspores cv. "Razzek" (Figure 2a), the first induction occurred 3 days after culture and were observed in the fraction of enlarged microspores

Table 1. Composition of CHB3 medium used for durum wheat isolated microspores culture based on medium of (Chu et al., 1990) modified.

Components	Quantity
Macro elements (g^l⁻¹)	
KNO ₃	1.415
(NH ₄) ₂ SO ₄	0.232
KH ₂ PO ₄	0.200
CaCl ₂ , 2H ₂ O	0.083
MgSO ₄ , 7H ₂ O	0.093
FeCl ₃	0.027
Na ₂ EDTA	0.0373
Micro elements (mg^l⁻¹)	
ZnSO ₄ , 7H ₂ O	5
MnSO ₄ , 7H ₂ O	5
H ₃ BO ₃	5
KI	0.4
Na ₂ MO ₄ , 2H ₂ O	0.0125
CuSO ₄ , 5H ₂ O	0.0125
CoCl ₂ , 6H ₂ O	0.0125
Vitamins (mg^l⁻¹)	
Glycine	1
Thiamine HCl	2.5
Pyridoxine HCl	0.5
Nicotinic acid	0.5
Biotine	0.25
Calcuim panthotenate	0.25
Aminoacids (mg^l⁻¹)	
Glutamine	1000
Myo-Inositol	300
Growth regulators (mg^l⁻¹)	
2,4-D	0.5
Kinetin	0.5
Maltose g ^l ⁻¹	90
pH	5.4

(Figure 2b). These divisions led to the formation of embryogenic microspores that presented a fibrillar cytoplasmic structure (Figures 2c and 2d) which appeared after 7 days of culture in CHB3 medium. Only those were able to develop into proembryos. Ten days after culture, a fraction of proembryo emerged from exine wall (Figure 2e) and can develop into mature embryos after 25 days on culture (Figure 2f).

Embryos floating on the surface of the induction medium (Figure 3a) were formed from individual microspores by direct embryogenesis. These embryos were transferred to the solid regeneration medium and a fraction of them regenerated directly into green (Figure 3b) or albino plantlets (Figure 3c).

Effect of pre-treatments

Eight pre-treatments were compared in the present study

on cv. "Razzek", in order to select those more efficient to induce embryo formation and plant regeneration in durum wheat by isolated microspores culture (Table 2).

Isolated microspores were induced from all kind of pretreatments and also for the control after 3 days of culture except 15 days PEG₄₀₀₀ 1%. This pre-treatment was lethal to microspores. The absence of pre-treatment permits induction of microspores with frequency of 12.16 and 0.002% at 3 and 7 days of culture, respectively.

However, no embryogenesis was obtained without pre-treatment. For all pre-treatments, the viability of microspores decreased during the culture and between the third and seventh day of culture ranged from 4 to 17% (Table 1). No strong differences were found in isolated microspores response to mannitol based pre-treatments. Indeed, the first cell divisions could happen but no microspore could develop into an embryo.

The best results were obtained with cold pre-treatment




Figure 1. Cytological and morphological stages of isolated microspores culture of durum wheat. a: Optimal cytological stage of culture (late uninucleate stage). b: Optimal morphological stage corresponding to optimal cytological stage.




Figure 2. Durum wheat microspore embryogenesis main stages: a. Freshly isolated durum wheat microspores (Gx200); b. Microspores suspension after 3 days in culture : enlarged microspores (1); microspores arrested in the initial stage (2); degenerated microspores (3) (Gx200); c- Dividing microspores after 7 days in culture with fibrillar structure (1)(Gx200) and (1')(Gx400) ; d- microspores with delayed induction (2); degenerated microspores (3)(Gx200); e. Proembryo began to emerge from the exine 10 days after culture initiation (Gx200); f. Embryos formed after 20 - 25 days after culture in medium CHB3 observed with stereoscope.



Figure 3. Durum wheat plant regeneration from microspore embryos: a. Embryos developed after 4 weeks of culture in medium CHB3; b. Green regenerants in MS medium without growth regulators; c. Albino regenerants.

for 5 weeks followed by PEG₄₀₀₀ 1% for 10 days. This last pre-treatment induced a response of the isolated microspores with 46 embryos transferred and 8 plants regenerated. This pre-treatment appears to be promising and it is essential to perform new experiments to improve its response.

The cold pre-treatment improved significantly the microspores response in all steps of culture process. An enhancement of induction of microspores was obtained with a respective frequency of 14.06 and 10.58% for percentages of enlarged microspores at 3 and 7 days of culture. Increased rates of proembryos and embryos (2.82 and 0.12%) were obtained and 8 plants were regenerated per 270 embryos transferred.

Effect of donor plants genotype

The optimized pre-treatment (cold pre-treatment at 4 °C for five weeks) determined in earlier experiments were used in subsequent experiments. In this present study, three cultivars of durum wheat were used: Razzek, Khiar and Aziz, to test the genotypic effects on isolated microspores culture of durum wheat (Table 3).

Isolated microspores were cultured in CHB3 medium (liquid culture) in 35 × 15 mm plates and co-cultured with 6 – 7 ovaries that were derived from the same donor plants as the microspores. Table 3 showed that a strong genotypic variation does exist in the microspores response to *in vitro* culture. While there was no significant genotype difference relative to the frequency of dividing microspores at 3 days culture, difference were observed

at 7 days. Indeed, after 7 days of culture, no additional microspores from Azizi cultivar could initiate their differentiation. While, in other genotypes 7 to 9% of additional microspores were able to develop into the fibrillar structure. The efficiency of durum wheat microspores embryogenesis and regeneration were also genotype-dependent. We report that no response was obtained for experiments with cv. “Azizi”. In contrast, cv. “Razzek” showed the highest rates of embryo formation with 769 embryos transferred but no green plants were regenerated. Among the three genotypes used, the cv. “Khlar” gave the best response in terms of green plant regeneration. This genotype regenerated the highest number of green plants (43 green plants from 414 embryos transferred) with green/albino ratio ≈2 (43 green plants and only 23 albino).

DISCUSSION

The value of *in vitro* culture procedure in breeding programs depends primarily on the efficiency of the method used. Significant progress has been made in plant regeneration via androgenesis but to date no efficient method of plant regeneration has been established that would be universal for most cereals. Adjustments and adaptations for each individual species are necessary (Oleszcuk et al., 2004) and until now, little reports were published for durum wheat.

We report here three important factors on the efficient production of embryos and plants from isolated durum wheat microspores.

Table 2. The effect of pre-treatments on isolated microspore culture response of durum wheat cv. "Razzek".

Pre-treatment	Total number of extracted and cultivated microspores	Number and (%) of enlarged microspores		Number and (%) of pro-embryos	Number and (%) of transferred embryos	Number and (%) of regenerated plantlets	
		3 days of culture	7 days of culture			Albina	Green
No pre-treatment	375000	45600 (12.16) abc	8 (0.002) b	-	-	-	-
5 weeks cold at 4°C	225000	31635 (14.06) ab	23805 (10.58) a	6345 (2.82) a	270 (0.12)	8 (2.96)	-
0.3M Mannitol for 7 days	450000	58680 (13.40) abc	15345 (3.41) b	11 (0.024) b	-	-	-
0.3M Mannitol for 7 days	450000	58680 (13.40) ab	15345 (3.41) b	11 (0.024) b	-	-	-
0.7M Mannitol for 5 days	300000	57810 (19.27) a	7440 (2.48) b	-	-	-	-
PEG ₄₀₀₀ 1.5% for 5 days	150000	7320 (4.88) bc	-	-	-	-	-
PEG ₄₀₀₀ 1% for 10 days	150000	26085 (17.39) ab	6165 (4.11) b	79(0.053) b	46(0.03)	8 (17.4)	-
PEG ₄₀₀₀ 1% for 15 days	150000	-	-	-	-	-	-

PEG: Polyéthylène glycol; M : Molarity. Number and $\leq\%$ in a same column followed by different letter are significantly different at $P=0.05$.

First, stress treatment is essential to induce formation of embryogenic microspores. Many authors reported the importance of pre-treatment for successful isolated microspores culture in cereals and suggest a necessity of specific stress treatment to trigger microspore development from gametophytic to sporophytic pathways (Jähne and Lörz, 1995; Hu and Kasha, 1999; Touarev et al., 2000; Liu et al., 2002 and Zheng et al., 2003). Ramirez et al. (2001) showed that in barley, the osmotic regulation was favourable to switch the microspore program regardless of the nature of the osmoticum.

In durum wheat, different results were obtained with cold pre-treatments and cold mannitol by (Labrani et al., 2005; Cistué et al., 2006; Labrani et al., 2007). In our study, we compare these pre-treatments: cold or cold with osmotic pre-treatments including mannitol and other osmotic pre-treatments using PEG at different concentrations and at different time intervals.

Considering our results obtained from isolated

microspores culture of durum wheat cv. "Razzek", the best result was achieved with cold pre-treatment (4°C) for 5 weeks that appeared to be the most efficient treatment for embryogenesis induction and embryos development of durum wheat. These results are in agreement with previous studies in bread wheat (Ohnoutková et al., 2000; De Buyser et al., 2002) and in durum wheat (Labrani et al., 2005). It was proposed by Duncan and Heberle (1976) that cold pre-treatment slows down degradation processes in the anther tissues thus protecting microspores from toxic compounds released in the decaying anthers. Furthermore, the use of cold pre-treatment improved the capacity of microspores of cv. "Khlar" to regenerate green plants. Hoekstra et al. (1993) showed also the importance of the osmotic pressure during the pre-treatment in order to obtain green plantlets. Our results allow also confirming that embryogenesis and regeneration were strongly dependent on the type of osmoticum used. In barley, the combination of cold (4°C)

and osmotic (mannitol) regulation during pre-treatment was recently found to be more appropriate using isolated microspores culture (Hoekstra et al., 1993). Mannitol has been recognized to penetrate cells passively and slowly (Cram, 1984) because the plasmalemma is poorly permeable to it (Rains, 1989). Wojnarowicz et al. (2004) showed that mannitol induces both a stable osmotic pressure in the pre-treatment medium and a starvation for microspores during 3 days at 4°C. When mannitol is used at high concentration, water is unable to penetrate into the microspores, probably leading to their alteration. In contrast, in durum wheat, Cistué et al. (2006) improved the rate of regeneration with 0.7M mannitol for 5 to 6 days at 24°C. In our work, the absence of embryogenesis that we noted with cold mannitol pre-treatments at different concentrations and at different times may be related to the recalcitrance of the genotype used.

Some genotypes naturally respond better than others to pre-treatment applied (Liu et al., 2002).

Table 3. Comparison of induction, embryogenesis and plant regeneration in isolated wheat microspores cultures of 3 genotypes of durum wheat pretreated with cold (4°C) for 5 weeks.

Genotype	Total number of microspores extracted and cultured	Percentage of induction (enlarged microspores)		Percentage of pro-embryos	Number of embryos transferred	Number of regenerated plantlets	
		3 days culture	7 days culture			Green	Albina
Razzek	600000	14.12 a	9.08 a	2.25 ab	769	-	23
Khlar	450000	17.96 a	7.04 ab	4 a	414	43	23
Azizi	300000	10.18 a	- b	- b	-	-	-

Numbers in the same column followed by different letters are significantly different at 0.05 level by ANOVA test based on a minimum of 3 replications per experiments. The number of regenerated plants was based on the number of transferred embryos.

According to Li and Devaux (2001), the high and low responding genotypes have different requirements for optimal pre-treatment conditions in the same species. Labbani et al. (2005) used two varieties of durum wheat and the quantity and quality of embryos obtained depended on the variety used for the same pre-treatment. Similarly, Li and Devaux (2001) showed that mannitol (0.3M), which is used frequently in barley and which has improved the embryogenesis and regeneration of green plants, proved ineffective in certain recalcitrant barley genotypes. One main problem with some genotypes and some pre-treatments (like mannitol in our case) appeared between one and two weeks of culture. The majority of pro-embryos cannot burst out the exine of microspores (Hu and Kasha, 1999).

We showed that PEG₄₀₀₀ 1% for 10 days pre-treatment has a beneficial effect on isolated microspores response with high frequency of induction at 3 days of culture and regenerated plantlets. The beneficial effect of PEG is in agreement with previous report of Wojnarowicz et al. (2004) that tested mannitol and PEG at different steps of anther culture in barley to elucidate their influence on both, the overall yield of androgenesis and the structure of plastids in relation to albinism. These authors showed that the use of PEG during pre-treatment was more favourable than mannitol for inducing anther response. The high viscosity of the PEG solution and the restricted water supply led to a high number of plantlets. This sugar is not metabolized by plant cells and on their presence, plastids did not accumulate starch at any stage of the protocol but they started to differentiate into chloroplasts in the microspore-derived embryos.

Considering the rate of embryogenesis obtained with cold PEG₄₀₀₀ 1% for 10 days, it is important to perform new experiments with PEG in the aim to confirm their beneficial influence in durum wheat microspores culture and improve their response.

Second important point for the success of microspores culture in durum wheat is obtaining embryogenic microspores with fibrillar structure at 7 days in culture. Only this multicellular structure was embryogenic. In the first week of culture the viability of microspores decreased between 4 and 17% depending on the

evolution of microspores. The microspores were capable of following one of three pathways. A fraction of microspores underwent plasmolysis, another fraction of microspores remained blocked in the initial stage. Another fraction of them increased in size with fibrillar structure. Only the latter types of microspores can be embryogenic. The same types of cells were observed in bread wheat by Zheng et al. (2001). This result is in accordance with previous work. Ramirez et al. (2001) showed in barley that most of the cells which responded to the stress treatment were vacuolate microspores indicating that this developmental stage is responsive for embryogenesis induction in barley. Shariatpanahi et al. (2006) proposed that in some species like wheat, that we still do not know exactly what feature of the microspores can be introduced as universal sign of the embryogenic (totipotent) state. Whereas, microspores do not have a starlike cytoplasmic structure but show an increase in the size of the cell and symmetrical divisions can be embryogenic. Our results suggest also that rates of induced microspores at one week from 7% indicated a possible embryogenic evolution.

Third important factor in durum wheat microspores culture is the genotypic effect during the production of haploid plants. This factor has been reported by several research groups (Labbani et al., 2005; Cistué et al., 2006; Slama et al., 2006; Labbani et al., 2007). As the experiment described in our work, it is possible that only albino plants obtained for cv. "Razzek" and the high frequency of green plants for cv. "Khlar" regenerated with the same method were a characteristic of the vegetal material used. Zamani et al. (2000) reported that embryos formation and green plants regeneration were strongly affected by the genotype of donor plants and the induction medium, while albino production was affected only by the genotype. These two genotypes can be exploited for studies about understanding the mechanism of albinism.

In this work, among the three genotypes tested two were responsive and produced embryos and regenerated plantlets. For cv. "Khlar", from a total of 414 embryos which were transferred on to the regeneration medium, 66 plants were regenerated and 43 of them were green.

Thus, in this cultivar pretreated with cold 4°C for 5 weeks, up to 10% of the transferred embryos were regenerated in green plants and more than 65% of regenerated plantlets were green. The frequency of green regeneration derived from isolated microspores cultured of cv. “Khiar” reported here gave the best frequency of microspore green plant regeneration published until now in durum wheat where albinism seemed to be the major problem (Cistué et al., 2006). In fact, Jaiti et al. (2000) obtained 15 green plants from 1097 plants regenerated from 2824 embryos of durum wheat microspores culture cultivated that represented a rate of 1.31%. In another experience performed by Labbani et al. (2007), 85 green plants were regenerated from 3673 embryos that represented a frequency of 2.31%.

These results indicated that the use of cold pre-treatment improves frequency of green plant regeneration. Yet the number of regenerated plantlets still stay low, some progress has been obtained with durum wheat green regeneration with 43 green plants regenerated and only 23 albina (green /albino ratio ≈2). Green plant regeneration rates obtained in this study were satisfactory with durum wheat where albinism seemed to be the major problem and that considered as a recalcitrant cereal.

The present protocol of microspore culture, in durum wheat, may be widely applicable for increasing yield of green plant regeneration. This result encouraged the potential for using isolated microspores of durum wheat for breeding and biotechnological applications. However, further studies on the optimization of pre-treatments and protocols are recommended.

REFERENCES

- Chu CC, Hill RD, Brule-Babel AL (1990). High frequency of pollen embryoid formation and plant regeneration in *Triticum aestivum* L. on monosaccharide containing media. *Plant Sci.* 66: 255-262.
- Cistué L, Vallés MP, Echàvarri B, Sanz JM, Castillo A (2003). Barley anther culture. Doubled Haploid production in crop plants, ed. Maluszynski M, Kasha KJ, Forester BP and Szarejko I. Kluwer Acad. Publishers pp. 29-34.
- Cistué L, Soriano M, Castillo AM, Vallés MP, Sanz JM, Echàvarri B (2006). Production of doubled haploids in durum wheat (*Triticum turgidum* L.) through isolated microspore. *Plant Cell. Rep.* 25: 257-264.
- Cistué L, Romagosa I, Batlle F, Echàvarri B (2009). Improvements in the production of doubled haploids in durum wheat (*Triticum turgidum* L.) through isolated microspore culture. *Plant Cell Reports* 28(5): 727-735.
- Cram WJ (1984). Mannitol transport and suitability as an osmoticum in root cells. *Physiol. Plant* 61: 396-404.
- De Buyser J, Touraine P, Jaiti F, Picard E (2002). Haplodiploidisation par culture de microspores isolées de blé *in vitro*. *Biotechnologies végétales : techniques de laboratoire*. Livre Robert Haicour Edi Tec et Doc Montréal : AUF 2002: 257-273.
- Duncan EJ, Heberle E (1976). Effect of temperature shock on nuclear phenomena in microspores of *Nicotiana tabacum* and consequently on plantlet production. *Protoplasma* 90: 173-177.
- Hoekstra S, van Zijderveld MH, Heidekamp F, van der Mark F (1993). Microspore culture of *Hordeum vulgare* L.: the influence of density and osmolality. *Plant Cell Rep.* 12: 661-665.
- Hu T, Kasha KJ (1997). Improvement of isolated microspore culture of wheat (*Triticum aestivum* L.) through ovary co-culture. *Plant Cell Rep.* 16: 520-525.
- Hu T, Kasha KJ (1999). Cytological study of pretreatments used to improve isolated microspore cultures of wheat (*Triticum aestivum* L.) c.v. chris. *Genome* 42: 432-441.
- Ilić-Grubor K, Attree SM, Fowke LC (1998). Induction of microspore-derived embryos of *Brassica napus* L. with polyethylene glycol (PEG) as osmoticum in a low sucrose medium. *Plant cell reports* 17: 329-333.
- Jaiti F, El Jaafari S, El Hadrami I, Nachit M, Baum M, De Buyser J, Picard E (2000). Problématique de la culture de microspores chez le blé dur *Triticum turgidum* L. var. durum. L'amélioration du blé dur dans la région méditerranéenne, nouveau défis série A 40: 149-152.
- Kasha KJ, Simion E, Oro R, Yao QA, Hu TC, Carlson AR (2001). An improved *in vitro* technique for isolated microspore culture of barley. *Euphytica* 120: 379-385.
- Labbani Z, Richard N, De Buyser J, Picard E (2005). Plantes chlorophylliennes de blé dur obtenues par culture de microspores isolées: importance des prétraitements. *C. R. Biologies* 328: 713-723.
- Labbani Z, De Buyser J, Picard E (2007). Effect of mannitol pre-treatment to improve green plant regeneration on isolated microspore culture in *Triticum turgidum* ssp. durum cv. *Plant Breed.* 126: 565-568.
- Li H, Devaux P (2001). Enhancement of microspore culture efficiency of recalcitrant barley genotypes. *Plant Cell Reports* 20: 475-481.
- Li H, Devaux P (2003). High frequency regeneration of barley doubled haploid plants from isolated microspore culture. *Plant Sci.* 164: 379-386.
- Liu W, Zheng MY, Konzak CF (2002). Improving green plant production via isolated microspore culture in bread wheat (*Triticum aestivum* L.). *Plant Cell Rep.* 20: 821-824.
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant* 15: 473-497.
- Oleszczuk O, Sowa S, Zimny J (2004). Direct embryogenesis and green plant regeneration from isolated microspores of hexaploid triticale (x Triticosecale Wittmack) cv. Bogo. *Plant Cell Rep.* 22: 885-893.
- Ohnoutková L, Novotny J, Mullerova E, Vagera J, Kucera L (2000). Is a cold pre-treatment really needed for induction of *in vitro* androgenesis in barley and wheat? *Biotechnological approaches for utilisation of gametic cells*. Cost 824 final meeting, Bled, Slovenia, 1 to 5 July 2000. 277: 33-37.
- Rains DW (1989). Plant tissue and protoplast culture: Application to stress physiology and biochemistry. In: Jones HG, Flowers TJ, Jones MB (eds) *Plant under stress*. Cambridge University Press, London pp. 181-196.
- Ramirez C, Testillano PS, Castillo AM, Vallés MP, Coronado MJ, Cistué L and Risuéno MDC (2001). The early microspore embryogenesis pathway in barley is accompanied by concrete ultrastructural and expression changes. *Int. J. Dev. Biol.* 45(S1): S57-S58.
- SAS (1988) SAS Institute SAS/STAT users guide. Release 6.03 SAS Inst, Cary, NC (ed) 1988 p. 1028.
- Shariatpanahi ME, Bal U, Heberle-Bors E, Touraev A (2006). Stresses applied for the re-programming of plant microspores towards *in vitro* embryogenesis. *Physiologia Plantarum* 127: 519-534.
- Shariatpanahi ME, Belogradova K, Hessamvaziri L, Heberle-Bors E, Touraev A (2006). Efficient embryogenesis and regeneration in freshly isolated and cultured wheat (*Triticum aestivum* L.) microspores without stress pretreatment. *Plant Cell Rep.* 25: 1294-1299.
- Slama-Ayed O, Slim-Amara H, De Buyser J, Picard E (2006). Effet du prétraitement des épis sur l'aptitude de la régénération *in vitro* des microspores isolées chez une variété tunisienne de blé dur (*Triticum durum* Desf). *Revue de L'INAT* : 21(1): 123-132.
- Touarev A, Indrianto A, Wratchko I, Vincente O, Heberle-Bors E (1996). Efficient microspore embryogenesis in wheat (*Triticum aestivum* L.) induced by starvation at high temperature. *Sex Plant Rep.* 9: 209-215.
- Touarev A, Tashpulatov A, Indrianto A, Barinova I, Katholnigg H, Akimcheva S, Ribarits A, Voronin V, Zhexsembekova M, Heberle-Bors E (2000). Fundamental aspects of microspore embryogenesis. In *Biotechnological approaches for utilisation of gametic cells*. Cost 824 final meeting, Bled, Slovenia pp. 205-214.
- Wojnarowicz G, Carreda S, Devaux P, Sangwan R, Clément C (2004).

Barley anther culture: assessment of carbohydrate effects on embryo yield, green plant production and differential plastid development in relation with albinism. *J. Plant Physiol.* 161: 747-755.

Zamani I, Gouli E, Roupakias D (2000). Production of doubled haploids in Wheat (*Triticum aestivum* L.). Biotechnological approaches for utilization of gametic cells, Cost 824 final meeting, Bled, Slovenia pp. 43-51.

Zheng MY, Liu W, Weng Y, Polle E, Konzak CF (2001). Culture of freshly isolated wheat (*Triticum aestivum* L.) microspores treated with inducer chemicals. *Plant Cell Rep.* 20: 685-690.

Zheng MY, Weng Y, Liu W, Konzak CF (2002). The effect of ovary-conditioned medium on microspore embryogenesis in common wheat (*Triticum aestivum* L.). *Plant Cell Rep.* 20: 802-807.

Zheng MY, Liu W, Weng Y, Polle E, Konzak CF (2003). Production of doubled haploids in Wheat (*Triticum aestivum* L.) through microspore embryogenesis triggered by inducer chemicals. In M. Maluszynski et al. (eds.), *Doubled Haploid production in crop plants* pp.83-94.