

Full Length Research Paper

## Chemical composition and biological activity of essential oils of Cleopatra mandarin (*Citrus reshni*) cultivated in Egypt

Dalia I. Hamdan, Rehab H. Abdulla, Maged E. Mohamed and Assem M. El-Shazly\*

Pharmacognosy Department, Faculty of Pharmacy, University of Zagazig, Zagazig 44519, Egypt.

Accepted 18 April, 2013

**A comparative study of the essential oils isolated from the leaf and fruit peel of Cleopatra mandarin (*Citrus reshni*) was carried out using gas liquid chromatography (GLC) and GLC/mass analysis (MS). Over 140 components were observed in GLC/MS, of which 123 could be identified. 112 compounds were quantitatively analyzed in the leaf oil with linalool as a major component. The total identified components in the fruit peel oil were 69 and limonene was the most prominent. Cleopatra mandarin volatile components showed high anti-inflammatory activity represented by its effect on tumour necrosis factor- $\alpha$  and nitric oxide. They also showed significant anti-microbial activities against most common Gram positive and Gram negative bacteria and some fungi.**

**Key words:** Cleopatra mandarin, essential oils, gas liquid chromatography (GLC), GLC/mass analysis (MS), anti-inflammatory, antimicrobial.

### INTRODUCTION

Plants belonging to genus *Citrus* are known for their nutritional value, unique flavour and medicinal properties. The members of this genus are characterized by many biologically active secondary metabolites such as flavonoids (Tripoli et al., 2007), limonoids (Manners, 2007), coumarins and furanocoumarins, sterols (Ladaniya, 2008), volatile oils (Espina et al., 2010; Tranchida et al., 2011), organic acids and alkaloids (He et al., 2010). Many *Citrus* species are recognized for their medicinal, physiological and pharmacological activities including antimicrobial (Espina et al., 2010; Singh et al., 2010), antioxidant (Barros et al., 2012; Goulas and Manganaris, 2011), anticancer (Benavente-Garcia and Castillo, 2008; Manthey and Guthrie, 2002), anti-inflammatory (Menichini et al. 2011) and hypoglycaemic (Aruoma et al., 2012) activities.

The original source of Cleopatra mandarin was in India and introduced to Florida from Jamaica in the 19th Cen-

tury. The fruit is orange-red in colour with thin and rough peel and acidic flavour. A survey of the literature revealed very few reports on the identification of secondary metabolites in Cleopatra mandarin. Lota et al. (2001) reported the presence of 29 and 44 volatile components in the fruit peel and leaf oil of Cleopatra mandarin growing in France, respectively. The hydrodistilled oil of the fruit peel gave limonene as the major component (93.6%) followed by myrcene (1.7%), sabinene (1.1%) and linalool (1%). Sabinene (49.7%) was the prominent constituent in the leaf oil followed by linalool (13%) and (*E*)  $\beta$ -ocimene (6.9%) (Lota et al., 2001). Few reports discussed the biological importance of Cleopatra mandarin constituents. The presence of various polymethoxy flavones contributes to the antifungal properties of the plant (Uckoo et al., 2011), in addition, the plant is used for treatment of capillary fragility, haemorrhages and hypertension due to the presence of high percentage

\*Corresponding author. E-mail: [assemels2002@yahoo.co.uk](mailto:assemels2002@yahoo.co.uk) or [a\\_m\\_elshazly@zu.edu.eg](mailto:a_m_elshazly@zu.edu.eg). Tel: +2(0)552303266. Fax: +2(0)552303266.

of hesperidin (Cano and Bermejo, 2011).

There has been no detailed research on the volatile components of Cleopatra mandarin grown in Egypt. Therefore, and in continuing to the investigation of the chemical and biological activities of *Citrus* spp. cultivated in Egypt (El-Readi et al., 2010; Hamdan et al., 2010a, b; Hamdan et al., 2011), the chemical profile of the fruit peel and leaf oils of Cleopatra mandarin were analyzed by GLC and GLC/MS and their anti-inflammatory and antimicrobial activities were also assessed in this study.

## MATERIALS AND METHODS

### Plant

The fresh ripe fruits and leaves of Cleopatra mandarin (*Citrus reshni* Hort. ex Tan.), family Rutaceae were collected on the Research Station of the Faculty of Agriculture (Benha University, Egypt) in March, 2011. The identity of the plants was confirmed by Dr. B. M. Houlyel, Prof. of Pomology, Faculty of Agriculture, Benha University. Voucher specimens were deposited at the Department of Pharmacognosy, Faculty of Pharmacy, Zagazig University, Egypt.

### Isolation of the oil

The fresh leaves and ripe fruit peel (100 g each) of Cleopatra mandarin were separately subjected to hydro-distillation for 6 h using a Clevenger-type apparatus. Both oils were dried over anhydrous sodium sulphate and kept in brown vials in the refrigerator at 4°C until further analyses.

### Gas liquid chromatography (GLC) and GLC/mass analysis (MS)

The volatile oil constituents were analysed by high-resolution capillary GLC and GLC-MS. Oil samples (1 µl each, dissolved in 1 ml n-hexane) were injected (1 µl volume) into a gas chromatograph (TRACE GC ULTRA, Thermo Scientific, Milan, Italy) under the following conditions: column, RTX-5MS<sup>®</sup> fused silica capillary equivalent to DB-5 (30 m × 0.32 mm i.d and 0.25 µm film thickness); He as the carrier gas (2 ml/min); flame ionization detector (FID), temperature (300°C), injection temperature (250°C); oven temperature program: initial temperature at 45°C, 2 min isothermal, 300°C, 4°C/1 min, and then 20 min isothermal; split ratio, 1:15. Retention indices (RI) were calculated with respect to a set of co-injected homologous series of saturated hydrocarbon standards (C10 to C28). Components were quantified as area percentage of total volatiles from GC analyses as shown in Table 1. GLC-MS data were recorded on a Clarus 600 gas chromatograph (Connecticut, USA) equipped with an identical column used for separation and quantification. The capillary column was directly coupled to a quadruple mass spectrometer Clarus 600T. The ionization energy for the mass spectrometer was 70 eV. Split ratio was 1:30; other conditions were identical to those mentioned for GLC.

### Identification of components

Compounds were identified by comparing their spectral data and RI with Wiley Registry of Mass Spectral Data 8th edition, NIST Mass Spectral Library (December 2005), and literature data (Adams, 2007; Hamdan et al., 2010a). Where possible, retention times and

mass spectra were also compared with those of authentic pure samples. Most of the non-identified components are present as traces with relative abundances of less than 0.01%. The identified constituents are listed in the order of their elution in Table 1.

### Estimation of tumor necrosis factor-α (TNF-α)

TNF-α was induced in Raw murine macrophage (RAW 264.7) cells using lipo-polysacchride (LPS). The macrophage cells were treated with LPS alone and with the oil samples (100 µg/ml) for 48 h. The level of TNF-α was quantified in the serum produced from the macrophage cells by ELISA. The assay uses the quantitative sandwich immunoassay technique that uses immobilized monoclonal antibody and biotin-linked polyclonal antibody, both of which are specific against mice TNF-α. Commercially available matched paired antibodies were used (R&D Systems Inc. Minneapolis, MN). 4 µg/ml of anti-TNF-α monoclonal antibody and biotin-labelled anti-TNF-α polyclonal antibody (200 ng/ml) were used. The first (capture) antibody was coated onto 96-well flat bottom microtiter plate (Griener Labortechnik, Kremsmunster, Austria) in phosphate-buffered saline (PBS; Sigma Chemical Company, St. Louis, MO, USA), 50 µl/well and incubated 1 h at 37°C, then overnight at 4°C in humidified chamber. Plates were washed three times with washing buffer and blocked with 200 µl/well blocking buffer and incubated at 37°C for 1.5 h. Triplicate assays on 50 µl aliquots of serum samples were quantified by reference to recombinant human standards (R&D Systems, Inc. USA) added to each plate and incubated for 1 h at 37°C. At the end of the incubation period, the plates were washed three times with washing buffer and diluted second biotin labelled antibody was added for 1 h incubation at 37°C. After washing away any unbound substances, the peroxidase-conjugated streptavidin (Jackson Immunesearch Lab, USA) diluted 1:1000 was added to as 50 µl/well, then the plates were incubated for 1 h at 37°C. After an intensive washing, the enzyme reaction was carried out by adding a 50 µl/well of substrate solution. Color development was stopped by addition of 50 µl/well of stopping buffer (1 M HCl) (Surechem Products, Needham Marker, Suffolk, England). The intensity of the developed color was measured by reading optical absorbance at 450 nm using a microplate reader (FLUOstar OPTIMA, BMG LABTECH GmbH, Offenburg, Germany). Enzyme-linked immune sorbent assay (ELISA) reader-controlling software (Softmax) readily processes the digital data of raw absorbance value into a standard curve from which TNF-α concentration of unknown samples can be derived directly.

### Estimation of nitric oxide (NO)

Assay of nitrite accumulation (as an indicator of NO production) in the supernatants of cultured RAW 264.7 based on the Griess reaction was done according to Green et al. (1982). The cells were treated with the samples (100 µg/ml) for 48 h and compared with control. The concentration of nitrate was measured by reading the absorbance at 540 nm. The NO level of each of the tested cells was expressed using the following equation: NO level of the tested cells ×100/NO level of the control.

### Antimicrobial activity

Cup-plate method (Wood and Washington, 1995) was used to detect the preliminary antimicrobial activity. The test microorganisms: *Staphylococcus aureus* ATCC 6538 (Gram positive bacteria), *Pseudomonas aeruginosa* ATCC 9027, *Klebsiella pneumoniae* ATCC 27736 and *Escherichia coli* ATCC

**Table 1.** Volatile oil components identified in the fruit peel and leaf oils of *Cleopatra mandarin*.

Component	RI	Composition (%)	
		Fruit peel	Leaf
$\alpha$ -Thujene*	943	0.09	0.87
$\alpha$ -Pinene**	947	1.11	2.03
$\alpha$ -Fenchene	957	0.03	0.06
Camphene**	962	tr.	tr.
Sabinene**	977	0.98	22.27
$\beta$ -Pinenene**	999	-	0.03
Myrcene**	994	3.45	2.91
$\alpha$ -Phellandrene**	1002	-	0.27
$\delta$ -2-Carene	1005	0.74	0.05
$\alpha$ -Terpinene*	1013	0.01	1.41
$\rho$ -Cymene	1022	-	0.52
Limonene**	1026	79.46	3.78
(Z)- $\beta$ -Ocimene**	1037	-	0.37
(E)- $\beta$ -Ocimene**	1049	0.45	5.39
$\gamma$ -Terpinene**	1057	-	3.43
cis-Sabinene hydrate	1064	-	0.70
n-Octanol**	1076	3.33	tr.
$\rho$ -Mentha-2,4(8)-diene	1086	0.37	1.17
Linalool**	1092	3.26	23.91
trans-Sabinene hydrate**	1095	-	0.13
trans- $\rho$ -Mentha-2,8-dien-1-ol	1116	0.06	0.08
cis- $\rho$ -Menth-2-en-1-ol*	1121	-	0.27
allo-Ocimene	1129	0.01	0.24
cis- $\rho$ -Menth-2,8-dien-1-ol	1132	0.02	0.07
trans- $\rho$ -Menth-2-en-1-ol	1138	-	0.19
neo-allo-Ocimene	1140	-	0.05
cis- $\beta$ -Terpineol	1145	0.02	tr.
trans-Verbenol	1148	-	0.02
Citronellal	1154	0.02	tr.
Karahanaenone	1156	-	0.01
iso-Isopulegol	1160	-	0.01
neois-Isopulegol	1165	-	0.01
Terpinen-4-ol**	1178	0.52	6.32
trans- $\rho$ -Mentha-1(7),8-diene-2-ol	1184	-	0.04
$\alpha$ -Terpineol**	1190	0.84	1.34
Dihydro carveol	1194	0.05	0.09
n-Dodecane	1199	-	0.02
n-Decanal**	1206	1.01	-
trans-Piperitol	1207	-	0.21
iso-Dihydro carveol	1210	-	0.01
Octanol acetate	1214	0.05	tr.
trans-Carveol	1219	-	0.01
cis-Carveol	1223	-	0.05
Nerol*	1229	0.3	0.09
Thymol, methyl ether	1235	-	0.04
Cumin aldehyde	1238	-	0.05
Geraniol*	1242	0.07	tr.
Piperitone	1253	-	0.01
Methyl citronellate	1258	-	0.06

Table 1. Contd.

Geranial	1273	0.29	0.04
<i>n</i> -Decanol	1275	0.5	-
$\alpha$ -Terpinen-7-al	1283	-	0.03
Limonen-10-ol	1289	0.06	0.04
$\rho$ -Cymen-7-ol	1293	-	0.01
$\rho$ -Mentha-1-en-9-ol	1297	0.03	-
Terpinen-4-ol acetate	1300	0.11	-
<i>n</i> -Tridecane	1301	-	0.01
<i>cis</i> -Piperitol acetate	1326	-	0.02
$\delta$ -Elemene	1336	0.05	1.67
$\alpha$ -Cubebene	1347	-	0.04
$\alpha$ -Terpinyl acetate*	1351	-	tr.
Eugenol	1359	-	tr.
Longicyclene	1368	-	0.03
$\alpha$ -Ylangene	1372	0.03	0.05
$\beta$ -Panasinsene	1382	tr.	0.12
( <i>E</i> )- $\beta$ -Damascenone	1385	0.02	-
$\beta$ -Elemene*	1390	-	1.73
<i>n</i> -Tetradecane	1398	0.01	0.02
$\beta$ -Longipinene	1402	-	0.16
Dodecanal	1408	0.14	-
$\beta$ -Funebrene	1414	0.03	-
( <i>E</i> )-Caryophyllene**	1416	-	3.65
$\beta$ -Cedrene	1424	0.02	0.16
$\gamma$ -Elemene	1431	0.29	-
$\beta$ -Copaene	1433	-	3.67
$\alpha$ -Guaiene	1437	0.01	0.10
Aromadendrene	1441	tr.	0.07
$\alpha$ -Humulene**	1451	0.05	0.78
<i>cis</i> -Muurolo-4(14),5-diene	1460	0.02	0.05
$\gamma$ -Gurjunene	1475	-	0.23
$\gamma$ -Muurolole	1479	0.24	1.01
$\gamma$ -Himachalene	1483	tr.	-
$\beta$ -Selinene	1484	-	0.07
<i>trans</i> -Muurolo-4(14),5-diene	1489	tr.	-
<i>cis</i> - $\beta$ -Guaiene	1490	-	0.06
$\gamma$ -Amorphene	1494	0.01	-
$\alpha$ -Alaskene	1495	-	0.57
$\alpha$ -Muurolole	1500	tr.	0.05
$\gamma$ -Patchoulene	1503	-	0.08
( <i>E,E</i> )- $\alpha$ -Farnesene*	1504	0.04	-
$\delta$ -Amorphene	1507	-	0.17
$\gamma$ -Cadinene	1513	tr.	0.06
$\delta$ -Cadinene*	1523	0.06	0.31
<i>Z</i> -Nerolidol	1527	0.13	0.03
$\alpha$ -Cadinene	1533	-	0.04
<i>trans</i> -Cadina-1(2),4-diene	1537	0.01	-
<i>cis</i> -Sesquisabinene hydrate	1538	-	0.35
$\alpha$ -Calacorene	1542	-	0.03
Elemol*	1549	0.04	0.11
Germacrene B	1555	0.14	1.77
<i>E</i> -Nerolidol*	1564	-	0.12

Table 1. Contd.

Spathulenol*	1575	-	0.07
Caryophellene oxide*	1579	-	0.09
Carotol	1590	tr.	0.06
<i>n</i> -Hexadecane	1597	tr.	0.04
$\beta$ -Atlantol	1604	-	0.04
<i>Z</i> -Bisabolol-11-ol	1615	0.01	0.01
10- <i>epi</i> - $\gamma$ -Eudesmol	1623	-	0.03
$\gamma$ -Eudesmol	1628	0.02	0.37
<i>epi</i> - $\alpha$ -Muurolool	1642	tr.	0.17
$\beta$ -Eudesmol	1648	0.02	0.19
$\alpha$ -Eudesmol	1651	0.03	0.11
$\alpha$ -Cadinol*	1653	tr.	0.12
Khusinol	1679	-	0.02
<i>n</i> -Heptadecane	1700	tr.	0.01
Cedroxyde	1713	-	0.02
Mint sulfide	1733	-	0.01
<i>Z</i> -Lanceol	1758	-	tr.
14-hydroxy- $\alpha$ -Muurolole	1766	-	0.02
Guaiazulene	1772	-	0.02
$\alpha$ -Eudesmol acetate	1793	tr.	tr.
<i>n</i> -Octadecane	1797	tr.	0.02
Nootkatone	1806	tr.	tr.
Monoterpene hydrocarbons	-	87.04	54.31
Oxygen containing monoterpenes	-	5.97	13.48
Sesquiterpene hydrocarbons	-	1.22	24.15
Oxygen containing Sesquiterpenes	-	0.41	4.27
Others	-	5.04	0.08
Total	-	99.68	96.29

In elution order from RTX-5MS® column. RI = identification based on retention index relative to standard *n*-alkanes. tr. = trace (<0.01 %). - = not detected.

\*Previously reported in the leaf oil of Cleopatra mandarin (Lota et al., 2001).

\*Previously reported in the peel oil of Cleopatra mandarin (Lota et al., 2001).

10536 (Gram negative bacteria), *Aspergillus niger* ATCC 16404 and *Candida albicans* ATCC 10231 (fungi) are standard strains obtained from the Department of Microbiology, Faculty of Pharmacy, Zagazig University, Egypt. The nutrient agar or Sabouraud's agar were seeded by about  $10^6$  microbial cells. 100  $\mu$ l of fruit peel and leaf oils were separately dissolved in 500  $\mu$ l dimethyl formamide (DMF). Each cup was filled by about 100  $\mu$ l from each solution. Amoxicillin (500  $\mu$ g/ml) and amphotericin B (500  $\mu$ g/ml) were used as standard antibacterial and antifungal, respectively. The plates were incubated overnight at 37°C for bacteria and 30°C for fungi. Zones of inhibition were measured (mm).

#### Statistical analysis

The Student's unpaired *t*-test was used to detect the statistical significance, where a P value less than 0.05 was considered significant.

## RESULTS AND DISCUSSION

Essential oils obtained from plants have many potential applications, including their use as food additives, preservatives, in perfumes and in pharmaceuticals. These applications are usually due to the medicinal and pharmacological activities of these oils such as antimicrobial, antioxidant and anti-inflammatory activities. Volatile components of the Egyptian variety of Cleopatra mandarin were separated and identified using GLC and GLC/MS and their relative abundance is listed according to their retention indices in Table 1. The oil yields from the fruit peel and leaf were 2 and 0.8% (v/w), respectively. A total of 123 compounds were identified in the fruit peel and leaf oils. The composition of the fruit peel and leaf oils is very different. From the compounds

detected, 67 and 110 components have been identified in the respective oils, representing 99.86 and 96.29% of the oils. In the peel, limonene being the most abundant component (79.46%), this is typical for the orange oils. Only 19 of the identified components were previously reported in the literature and 48 compounds are now reported for the first time as volatile constituents of *Cleopatra* mandarin. On the other hand, we could not find any traces of  $\beta$ -phellandrene, terpinolene, octanal, nonanal, *cis*-limonene-1,2-oxide, octyl acetate,  $\alpha$ -copaene, *trans*- $\alpha$ -bergamotene, neral, germacrene-D and bicyclogermacrene as described by Lota et al. (2001). The main constituents of the leaf oil were linalool (23.91%), sabinene (22.27%), terpinen-4-ol (6.23%) and *E*-( $\beta$ )-ocimene (5.39%). It is worth noting that 81 compounds are now reported for the first time as volatile constituents of *Cleopatra* mandarin leaf oil. The compounds 3-carene,  $\beta$ -phellandrene, terpinolene, *p*-cymene, bicyclogermacrene, 6-methylhept-5-ene-2-one, nonanal, *cis*- and *trans*-limonene-1,2-oxide, (*E*)- $\beta$ -farnesene and manoyl oxide identified previously in the leaf oil of France species (Lota et al., 2001), were not detected in the oil of leaf specimen analyzed in the work. It is evident from the oil yield that seasonal variation, nutrition and temperature have an influence of oil production and its composition.

### Anti-inflammatory activity

The anti-inflammatory activity of *Cleopatra* mandarin oils was determined relative to its ability to inhibit both TNF- $\alpha$  and NO production. TNF- $\alpha$  is an inflammatory mediator which is associated with the development many inflammatory diseases such as rheumatoid, psoriasis and arthritis. Anti-TNF- $\alpha$  is a new approach which is now used in the treatment of many of the former inflammatory diseases (Palladino et al., 2003). NO is another inflammatory regulator which is produced from the amino acid L-arginine through the effect of the nitric oxide synthase. NO is generated through stimulation of many types of cells, especially macrophages, by stimulants such as LPS. If NO is produced in high concentrations out of control, damage of cells occurs due to cell injury. Measuring NO production is a method for assessing the anti-inflammatory effects of essential oils (Kierner et al., 2002). Many classes of natural compounds have been found to act as anti-TNF- $\alpha$  and NO inhibitor agents. These naturally occurring drugs has the advantage of being safer and sometimes more cost-effective than the chemically-synthesized inhibitors. *Citrus* essential oils are considered to have an anti-inflammatory activity through inhibition of NO production in target area (Yang et al., 2009).

Oils isolated from *Cleopatra* mandarin reduced the levels of TNF- $\alpha$  and NO in Raw murine macrophage cell culture (RAW 264.7) induced by LPS. The results indicated that the fruit peel oil at concentration of 100  $\mu$ g/ml, possessed a very high significant inhibitory activity


for LPS-stimulated TNF- $\alpha$  and NO levels to the extent nearly of the control level ( $P < 0.001$ ). The leaf oil also showed high significant inhibitory activity for LPS-stimulated TNF- $\alpha$  and NO levels ( $P < 0.01$ ) as shown in Figure 1. The peel oil showed the higher inhibition activity than leaf oil, respectively as an inhibitor for TNF- $\alpha$ ; however, both oils showed nearly the same activity as NO inhibitors. This inhibition activity can be attributed to the presence and the percentage of limonene in the oil. Limonene is reported to suppress the production of TNF- $\alpha$  and NO, thus becoming a potent anti-inflammatory agent especially in skin inflammatory condition (Yoon et al., 2010). Both TNF- $\alpha$  and NO, results and the good yield of oil by hydro-distillation suggests the ability of using *Cleopatra* mandarin fruit peel oil in skin preparation as an anti-inflammatory potent drug.

### Antimicrobial activity

Bactericidal and fungicidal activities are known properties for volatile oils, particularly those of *Citrus* spp. Many studies confirmed the antimicrobial activity of fruit peel oil of many *Citrus* spp. including *Cleopatra* mandarin (Caccioni et al., 1998; Espina et al., 2010; Fisher and Phillips, 2008). This study investigates the antimicrobial activity of the hydro-distilled essential oil of the fruit peel and the leaf of the Egyptian *Cleopatra* mandarin. The results revealed that all the tested essential oils had moderate antibacterial effect against all tested Gram positive and Gram negative bacteria as shown in Table 2. The activity ranged from 50 to 66% activity of amoxicillin as a standard broad spectrum antibiotic. The antimicrobial activity was nearly the same on Gram positive and Gram negative bacteria; however, the oils showed good activity against *P. aeruginosa* ATCC 9027 which is known for causing infection in lungs and urinary tract. The examined oils showed relatively strong antifungal activities especially on *A. niger* which reached to 82.6% activity of amphotericin B as an antifungal standard (Table 2). The difference between leaf and fruit peel oils in antimicrobial activities on both the bacteria and the fungi cannot be distinguished. These findings encourage the use of these oils as antimicrobial agents topically or internally which can be considered as a reuse and recycling of a waste product from *Citrus* spp. industries.

### Conclusion

Mandarins are one of the fruits which are highly consumed by human due to its nutritional and medicinal values. However, the cultivation and consumption of these fruits generate wastes by-products such as leaves and peel which could bring environmental problems if not properly dealt with. This study was carried out as a step towards reusing and recycling these by-products and



**Figure 1.** The anti-inflammatory activity of hydro-distilled essential oils of the fruit peel and leaf of Cleopatra mandarin. A: The level of tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ) protein in RAW 264.7 cells after treatment with the samples (100  $\mu$ g/ml) for 48 h compared with lipo-polysacchride (LPS) treated cells as measured by ELISA assay. B: Percentage inhibition of TNF- $\alpha$  by leaf and peel oils. C: The level of nitric oxide (NO) in RAW 264.7 cells supernatant after the treatment with the samples (100  $\mu$ g/ml) for 48 h compared with LPS treated cells as measured by Griess reaction according to Green et al. (1982). D: Percentage inhibition of NO by the leaf and peel oils. The data are presented as absorbance (mean  $\pm$ SE) of three replicates ( $n=3$ ).

**Table 2.** The antimicrobial activity of the fruit peel and leaf oils of Cleopatra mandarin.

Tested material	Activity (%)					
	<i>Staphylococcus aureus</i> ATCC6538	<i>Escherichia coli</i> ATCC10536	<i>Klebsiella pneumoniae</i> ATCC27736	<i>Pseudomonas aeruginosa</i> ATCC9027	<i>Candida albicans</i> ATCC10231	<i>Aspergillus niger</i> ATCC16404
Amoxicillin	100.00	100.00	100.00	100.00	0.00	0.00
Amphotericin B	0.00	0.00	0.00	0.00	100.00	100.00
Leaf oil	58.82	52.86	60.20	62.50	60.71	82.61
Fruit peel oil	44.12	55.56	66.00	64.56	53.57	65.22

Oil samples (100  $\mu$ l) dissolved in DMF (500  $\mu$ l) and each cup was filled with 100  $\mu$ l from each extract. The data is represented as an activity percentage of the standard antibacterial amoxicillin (500  $\mu$ g/ml) or antifungal amphotericin B (500  $\mu$ g/ml). The data is the mean of three replicates ( $n=3$ ).

focuses on the identification of new flavors that could have been used in perfumery, food industries and medicinal application.

## ACKNOWLEDGMENTS

The authors thank Mr. Nader Shawky, Department of

Immunology and Microbiology, Faculty of Pharmacy, University of Zagazig for kindly performing the antimicrobial analyses of the oils. They are also grateful to University of Zagazig for providing financial support (Grant No. 5K2/M4 to AE).

## REFERENCES

- Adams RP (2007). Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry, Allured Publishing corporation, Chicago, Illinois, USA.
- Aruoma OI, Landes B, Ramful-Baboolall D, Bourdon E, Neergheen-Bhujun V, Wagner K-H, Bahorun T (2012). Functional benefits of *CITRUS* fruits in the management of diabetes. *Prev. Med.* 54:12-16.
- Barros HR, Ferreira TA, Genovese MI (2012). Antioxidant capacity and mineral content of pulp and peel from commercial cultivars of *citrus* from Brazil. *Food Chem.* 134:1892-1898.
- Benavente-Garcia O, Castillo J (2008). Update on uses and properties of *citrus* flavonoids: new findings in anticancer, cardiovascular, and anti-inflammatory activity. *J. Agric. Food Chem.* 56:6185-6205.
- Caccioni DRL, Guizzardi M, Biondi DM, Agatino R, Ruberto G (1998). Relationship between volatile components of *citrus* fruit essential oils and antimicrobial action on *Penicillium digitatum* and *P. italicum*. *Int. J. Food Microbiol.* 43:73-79.
- Cano A, Bermejo A (2011). Influence of rootstock and cultivar on bioactive compounds in *citrus* peels. *J. Sci. Food Agric.* 91:1702-1711.
- El-Readi MZ, Hamdan D, Farrag N, El-Shazly A, Wink M (2010). Inhibition of P-glycoprotein activity by limonin and other secondary metabolites from *Citrus* species in human colon and leukaemia cell lines. *Eur. J. Pharmacol.* 626:139-145.
- Espina L, Somolinos M, Loreen S, Conchello P, Garcia D, Pageen R (2010). Chemical composition of commercial *Citrus* fruit essential oils and evaluation of their antimicrobial activity acting alone or in combined processes. *Food Control.* 22:896-902.
- Fisher K, Phillips C (2008). Potential antimicrobial uses of essential oils in food: is *citrus* the answer?. *Trends in Food Sci. Technol.* 19:156-164.
- Goulas V, Manganaris GA (2011). Exploring the phytochemical content and the antioxidant potential of *Citrus* fruits grown in Cyprus. *Food Chem.* 131:39-47.
- Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR (1982). Analysis of nitrate, nitrite, and <sup>15</sup>N nitrate in biological fluids. *Anal. Biochem.* 126:131-138.
- Hamdan D, El-Readi MZ, Nibret E, Sporer F, Farrag N, El-Shazly A, Wink M (2010a). Chemical composition of the essential oils of two *Citrus* species and their biological activities. *Pharmazie* 65:141-147.
- Hamdan D, El-Readi MZ, Tahrani A, Herrmann F, Kaufmann D, Farrag N, El-Shazly A, Wink M (2010b). Chemical composition and biological activity of *Citrus jambhiri* Lush. *Food Chem.* 127:394-403.
- Hamdan D, El-Readi MZ, Tahrani A, Herrmann F, Kaufmann D, Farrag N, El-Shazly A, Wink M (2011). Secondary metabolites of ponderosa lemon (*Citrus pyriformis*) and their antioxidant, anti-inflammatory, and cytotoxic activities. *Z. Naturforsch.* 66c:385-393.
- He D, Shan Y, Wu Y, Liu G, Chen B, Yao S (2010). Simultaneous determination of flavanones, hydroxycinnamic acids and alkaloids in *citrus* fruits by HPLC-DAD-ESI/MS. *Food Chem.* 127:880-885.
- Kiemer AK, Muller C, Vollmar AM (2002). Inhibition of LPS-induced nitric oxide and TNF- $\alpha$  production by  $\alpha$ -lipoic acid in rat Kupffer cells and in RAW 264.7 murine macrophages. *Immunol. Cell Biol.* 80:550-557.
- Ladaniya MS (2008). *Citrus* Fruit Biology, technology and evaluation, Academic press, London, UK. 1-11.
- Lota M-L, de Rocca Serra D, Tomi FI, Casanova J (2001). Chemical variability of peel and leaf essential oils of 15 species of mandarins. *Bioch. Syst. Ecol.* 29:77-104.
- Manners GD (2007). *Citrus* limonoids: analysis, bioactivity, and biomedical prospects. *J. Agric. Food Chem.* 55:8285-8294.
- Manthey JA, Guthrie N (2002). Antiproliferative activities of *citrus* flavonoids against six human cancer cell lines. *J. Agric. Food Chem.* 50:5837-5843.
- Menichini F, Loizzo MR, Bonesi M, Conforti F, De Luca D, Statti GA, de Cindio B, Menichini F, Tundis R (2011). Phytochemical profile, antioxidant, anti-inflammatory and hypoglycemic potential of hydroalcoholic extracts from *Citrus medica* L. cv Diamante flowers, leaves and fruits at two maturity stages. *Food Chem. Toxicol.* 49:1549-1555.
- Palladino MA, Bahjat FR, Theodorakis EA, Moldawer LL (2003). Anti-TNF- $\alpha$  therapies: the next generation. *Nat. Rev. Drug Discov.* 2:736-746.
- Singh P, Shukla R, Prakash B, Kumar A, Singh S, Mishra PK, Dubey NK (2010). Chemical profile, antifungal, anti-aflatoxigenic and antioxidant activity of *Citrus maxima* Burm. and *Citrus sinensis* (L.) Osbeck essential oils and their cyclic monoterpene, d-limonene. *Food Chem. Toxicol.* 48:1734-1740.
- Tranchida PQ, Bonaccorsi I, Dugo P, Mondello L, Dugo G (2011). Analysis of *Citrus* essential oils: state of the art and future prospective. A review. *Flavour Fragr. J.* 27:98-123.
- Tripoli E, Guardia ML, Giammanco S, Majo DD, Giammanco M (2007). *Citrus* flavonoids: Molecular structure, biological activity and nutritional properties: A review. *Food Chem.* 104:466-479.
- Uckoo RM, Jayaprakasha GK, Patil BS (2011). Rapid separation method of polymethoxyflavones from *citrus* using flash chromatography. *Se. Pu. Technol.* 81:151-158.
- Wood GL, Washington JA (1995). Antibacterial susceptibility tests: dilution and disk diffusion methods. *Manual of Clinical Microbiology* ed. Murray PR, Baron JE, Pfaller MA, Tenover FC and Tenover RH. Washington DC. American Society for Microbiology 1327-1341.
- Yang EJ, Kim SS, Oh TH, Baik JS, Lee NH, Hyun CG (2009). Essential oil of *citrus* fruit waste attenuates LPS-induced nitric oxide production and inhibits the growth of skin pathogens. *Int. J. Agric. Biol.* 11:791-794.
- Yoon WJ, Lee NH, Hyun CG (2010). Limonene suppresses lipopolysaccharide-induced production of nitric oxide, prostaglandin E<sub>2</sub>, and pro-inflammatory cytokines in RAW 264.7 macrophages. *J. Oleo Sci.* 59:415-421.