

Full Length Research Paper

Attenuation of *t*-Butylhydroperoxide induced oxidative stress in HEK 293 WT cells by tea catechins and anthocyanins

Kerio L. C.¹, Bend J. R.², Wachira F. N.^{1,3*}, Wanyoko, J. K.¹ and Rotich M. K.⁴

¹Tea Research Foundation of Kenya, P. O. Box 820, 20200, Kericho, Kenya.

²Department of Pathology, Schulich School of Medicine and Dentistry, University of Western Ontario, 1400 Western Road London, N6G 2V4, Canada.

³Biochemistry Department, Egerton University, P. O. Box 536, Egerton, Kenya.

⁴Chemistry Department, Egerton University, P. O. Box 536, Egerton, Kenya.

Accepted 25 November, 2011

The health promoting properties of catechins and anthocyanins have been of great interest to researchers in the recent past due to their significant *in vitro* and *in vivo* antioxidant activities. Most research on anthocyanins has been based on berry anthocyanins. These potentially health enhancing pigments are also found in some Kenyan tea cultivars. An *in vitro* study was carried out to determine the effects of pure catechins (EGCG and EC) and tea anthocyanin extract from cultivar TRFK 306/1 on *t*-Butylhydroperoxide (*t*-BHP) induced oxidatively stressed HEK 293 cells. The effects of the catechins and tea anthocyanin extract on untreated cells (without *t*-BHP) cells and cells treated (with *t*-BHP, 500 μ M) were determined by measuring cytotoxicity (lactate dehydrogenase (LDH) leakage from the cells) and depletion of cellular glutathione (GSH). Cells were preincubated with the antioxidants for 30 min before addition of *t*-BHP (500 μ M) and additional incubation for 6 h. The results showed that epigallocatechin gallate (EGCG) and epicatechin (EC) as well as tea anthocyanin extract significantly ($p < 0.001$) attenuated *t*-BHP induced LDH leakage in a concentration dependent manner in treated cells. One way ANOVA analysis showed significant ($p < 0.001$) differences in the various effective concentrations of the catechins and tea anthocyanin extract used. Intracellular GSH content was also increased in a dose dependent manner. From these results, it is concluded that anthocyanin rich tea from selected Kenyan cultivars may have cytoprotective effects against oxidative stressors.

Key words: Anthocyanins, catechins, oxidative stress, lactate dehydrogenase, glutathione, *t*-BHP, attenuate, antioxidant, reactive oxygen species (ROS).

INTRODUCTION

Kenya is one of the most important producers of black tea in Africa. In fact, Kenya supplies 22% of the world's black tea and this crop is a major foreign exchange earner for the country (TBK, 2010), contributing about 26% of all foreign exchange earnings and 4% of the gross domestic product (GDP). In 2010, tea earnings in Kenya was Kshs 98 billion compared to Kshs 69 billion

earned in 2009 (TBK, 2010). Tea production in 2010 was 399 million Kg compared to 314 million Kg in 2009, accounting for much of the increase in revenue. Despite high production, local consumption of tea in Kenya is very low. For example, the estimated national per capita consumption of tea was only 0.46 Kg made tea/person/year in 2009 (TBK, 2010). To spur local use of Kenyan tea, it may be necessary to add nutraceutical value to the existing black tea products or to diversify into other products that can use tea and tea products as a raw material. Some specialty tea products are particularly sought after because they are considered to be healthful,

*Corresponding author. Email: fwachira@yahoo.com. Tel: +254 0722 644279.

often because they contain or are believed to contain a higher content of biologically active antioxidants. Alternatively, teas can be more appealing to consumers because they contain colour additives or flavours.

The pharmacological properties of the tea plant (*Camellia sinensis*) have historically been ascribed to its polyphenols particularly the catechins; catechin (+)-C, epicatechin (EC), epigallocatechin (EGC), epigallocatechin gallate (EGCG), gallic catechin (GC) and epicatechin gallate (ECG) in green tea; and theaflavins (TFs) and thearubigins (TRs) in black tea. Tea catechins have been reported to exhibit the following pharmacological properties: radical scavenging of reactive oxygen species (ROS) and reactive nitrogen species (RNS) *in vitro* (Muzolf et al., 2008; Lee et al., 2008); metal chelation (Khokhar and Owusu, 2003; Tang et al., 2002); anticarcinogenic (Roy et al., 2005; Seely et al., 2005); antiinflammatory (Karori et al., 2008); antimicrobial (Almajano et al., 2008); antioxidant (Gramza et al., 2006; Luczaj and Skrzydlewska, 2005; Maurya and Rizvi, 2008; Karori et al., 2007); antidiabetic (Cabrera et al., 2006); pro-apoptotic in monocytes (Kawai et al., 2004) and protection of cells against oxidative stress-induced apoptosis (Yao et al., 2007). The most abundant and bioactive green tea catechin is EGCG (Cabrera et al., 2006). The antioxidant effectiveness of a tea therefore depends on the tea variety in concert with its content of EGCG (Katalinic et al., 2006). Though pharmacological properties of tea have traditionally been attributed to catechins in green tea and theaflavins in black tea (Leung et al., 2001), anthocyanins which occur in grapes, have also occur in some tea plants (Terahara et al., 2001; Kerio et al., 2011).

In the plant kingdom, anthocyanins occur ubiquitously conferring red, blue and purple colours to fruits, vegetables and some grains (Kong et al., 2003; Prior, 2003). Anthocyanins are most abundant in berries (*Vaccinium* sp.), grapes, apples, purple cabbage, eggplant, black carrots, purple fleshed sweet potato and grains like black rice and purple corn. In plants anthocyanins are localized in the cell vacuole (epidermis and peripheral mesophyll cells), in the leaves, stems, roots, flowers and fruits. In nature, over 600 different anthocyanins have been identified to date, though only six are predominant; cyanidin, peonidin, petunidin, pelargonidin, delphinidin and malvidin. A recent study we carried out on anthocyanin rich tea germplasm revealed that malvidin was the most predominant in Kenyan processed black and green tea (Kerio et al., 2011). The different anthocyanidins found in nature differ in their substitution patterns in the B-ring (hydroxylation and methylation), which also affects their biological activities. Anthocyanins have been found to exhibit superior antioxidant activities to some other polyphenols although just like the catechins in the tea plant, they have been reported to have a wide range of biological effects which include antioxidant (Bae and Suh, 2007; Orak, 2007;

Choi et al., 2007), anti-inflammatory (Dai et al., 2007; Arli and Cau, 2007) antimicrobial (Viskeliš et al., 2009; Heinonen et al., 2007), antiatherosclerotic (Mazza, 2007) and anticarcinogenic (Wang and Stoner, 2008) activities. Anthocyanins have also been implicated in induction of apoptosis in cancer cells (Hafeez et al., 2008; Lee et al., 2009) as well as with the chemoprotection of cells against oxidative stress-induced apoptosis (Elisia and Kitts, 2008), improvement of vision (Lee et al., 2005) and neuroprotective effects (Tarrozia et al., 2007). Anthocyanins have also been linked with the modulation of oxidative stress in cells which has long been recognized as a mediator of cell death either by necrosis or apoptosis (Yao et al., 2008; Elisia and Kitts, 2008). Anthocyanins do this by preventing the accumulation of intracellular ROS that cause loss of mitochondrial membrane potential ($\Delta\Psi_m$) (Haidara et al., 2002; Gogvadze et al., 2006). Apoptosis is an important physiological process of cell death that plays a critical role in tissue homeostasis (Krysko et al., 2008) but that also has been implicated in a number of pathological conditions for example Parkinson's Syndrome and Alzheimer's disease (Jellinger and Bancher, 1998; Erdem et al., 1998).

Though numerous potential pharmacological properties have been described for anthocyanins, little work has been carried out to establish the bioactivity of tea-derived anthocyanins. A cell culture assay based on human embryonic kidney (HEK 293) cells was used in this study to determine the ability of a tea anthocyanin extract and catechins to attenuate oxidative stress. Oxidative stress in cultured cells results in damage to three major categories of essential macromolecules, causing lipid peroxidation (Duthie et al., 2005) protein oxidation (Halliwell and Whiteman, 2004) and DNA damage due to formation of oxidized nucleotide derivatives (Lazze et al., 2003). To detoxify ROS, cells largely depend on innate antioxidants like glutathione (GSH), dietary antioxidants such as Vitamin C and E and enzymic antioxidants including superoxide dismutase, catalase, and the peroxiredoxins. The innate antioxidant glutathione (GSH), the oxidized form (glutathione disulphide) of which is a marker of oxidative stress, is a tripeptide (γ -glutamyl-cysteinyl-glycine) responsible, in significant part, for maintaining the redox balance of most cells, where it is the most abundant non-protein thiol (Lapenna et al., 1998). GSH can scavenge ROS directly or act as a co-factor for enzymatic defenses (Franco and Cidlowski, 2009). Glutathione is found almost exclusively in a reduced state in cells and the enzyme glutathione reductase, which is constitutively active and inducible upon oxidative stress, converts glutathione disulphide (GSSG) to GSH (Shih et al., 2005). The ratio of GSH:GSSG within cells is often used as a measure of cellular oxidative stress (Atakisi et al., 2010). If oxidative stress results in irreparable cell damage, then apoptosis (programmed cell death) are activated but if cells are

exposed to extremely high levels of ROS, they also become necrotic (Krysko et al., 2008).

In the present study, HEK 293 cells were treated with *t*-BHP (500 μ M) for 6 h to induce oxidative stress, to evaluate the ability of pretreatment with pure tea catechins and tea anthocyanin extract to attenuate the *t*-BHP-mediated cytotoxicity and GSH depletion.

MATERIALS AND METHODS

Anthocyanin extraction and purification

Five grams of ground steamed tea leaf from tea clone TRFK 306/1 samples were weighed into 250 ml conical flasks covered with foil and mixed with 50 ml methanol-1.5 M hydrochloric acid MeOH/HCl; (99/1v/v) and magnetically stirred at 900 rpm for 4 h at room temperature. The resultant solution was filtered and evaporated to dryness using a Rotavapour (Buchi Rotavapour R-300, Switzerland) under reduced pressure at 35°C. The extract was dissolved in 10 mL distilled water and passed through a 0.45 μ m membrane filter and kept in an ice bath until the next phase of isolation.

The extracts were passed through reverse phase (RP) C₁₈ solid phase extraction (SUPELCO, SPE) cartridges (Sigma-Aldrich, USA) previously activated with 10% MeOH/HCl and washed with 0.01% HCl in distilled water. Anthocyanins were adsorbed onto the column while sugars, acids and other water-soluble compounds were eluted from the column with 0.01% HCl in distilled water. Anthocyanins were then recovered using methanol acidified with 10% v/v formic acid. The cartridges were subsequently eluted with ethyl acetate (Fischer Scientific, UK) to remove phenolic compounds other than anthocyanins. The purified extracts were stored at -20°C until further analysis. The extracted anthocyanins crystallized upon solvent removal.

Catechins

The catechins (-)-epicatechin and (-)-epigallocatechin gallate were purchased from Sigma Aldrich, USA.

Incubation of HEK 293 cells

HEK 293 cells (Eton Bioscience, USA) were cultured in 75 cm³ flasks in minimum essential medium MEM (Invitrogen, Canada) and 10% FBS containing 0.5% gentamycin until confluent. Then 4 \times 10⁵ cells/mL were seeded in 60 mm Petri dishes for treatment prior to assay for GSH/GSSG or Western Blot or 4 \times 10⁵ cells/ml were seeded in 96 well plates for treatment prior to assay for lactate dehydrogenase release. The cells were incubated overnight at 37°C in a humidified atmosphere containing 95% air and 5% CO₂ prior to treatment.

LDH leakage

Overall cell damage was determined by LDH leakage into the extracellular medium according to the method of Korzeniewski and Callewaert, (1983). Cells were first pre-treated with varying concentrations of either the catechins (10-50 μ M) or the tea anthocyanin extract (10-100 μ M) or vehicle (blank) for 30 min prior to incubation with 500 μ M *t*-BHP (Sigma-Aldrich, Canada) for 6 h. Released LDH was analyzed at 490 nm using a Titertek Multiskan Microplate Reader (Phoenix AZ, USA). Cell damage was expressed

as the fraction of LDH released into the culture medium relative to the total cellular LDH activity determined after lysis of cells with 0.1% v/v Triton X-100 (Sigma-Aldrich, USA).

Analysis of intracellular GSH

The intracellular concentrations of GSH were determined according to the method of Habig et al. (1974). Cells were plated overnight in 2 ml of minimum essential medium (MEM) (Invitrogen, Canada) and treated with the antioxidants (catechins or anthocyanin) for 30 min after adding fresh media. After incubation with 500 μ M *t*-BHP for 6 h, the cells were harvested from the wells by gentle scraping, placed in ice, and centrifuged at 500 \times g for 5 min. Cell pellets were re-suspended in 250 μ l lysis buffer (10 mM Tris, 130 mM NaCl, 10 mM NaF, 10 mM NaH₂PO₄, 10 mM Na₂P₂O₇, and 2 mM EDTA) and quickly frozen in liquid nitrogen. The remaining aliquots were centrifuged for 2 min at 14,000 \times g, and the supernatants were used for GSH content measurement based on a GST enzymatic method of Habig et al., (1974). Sample supernatant (50 μ l) was added to wells of a 96-well plate and made up to 190 μ l with 0.1 M potassium phosphate buffer (pH 7.6). To each sample reaction, 5 μ l of 40 mM DNFB, and 0.2 unit of glutathione S-transferase were added, followed by incubation in the dark at room temperature for 20 min. The formation of S-2, 4-dinitrophenyl glutathione was measured by monitoring the absorbance at wavelength 340 nm with a Titertek Multiskan Microplate Reader (Phoenix AZ, USA). GSH content was expressed as nmol GSH/1000 cells.

Statistical analysis

The LDH assays were carried out in triplicate with three different batches of cultured cells and the data were subjected to one way analysis of variance. Student's *t*-test was used to determine significance between treatments. Differences between treatments were considered to be statistically significant if *p*<0.05. The GSH assays were carried out twice with different batches of cells and the means of the two experiments are reported here.

RESULTS

Effect of tea catechins and anthocyanins on LDH leakage from HEK 293 cells

Tea catechins and anthocyanins were compared for their ability to attenuate *t*-BHP induced oxidative stress in HEK 293 cells. Epicatechin (EC) and epigallocatechin gallate (EGCG) were used as representative tea catechins in the assays while the tea anthocyanin extract was obtained from steamed leaves of the tea clone TRFK 306/1, as described above. The molarity of the tea anthocyanin extract solution was obtained from the Relative Molecular Mass (RMM) of anthocyanins (449.2 g) as Cyanidin-3-glucoside equivalents. HEK 293 cells were subjected to oxidative stress by treatment with 500 μ M *t*-BHP, a cell-permeant organic oxidant that generates ROS and increased the amount of LDH released from these cells by disrupting cell membrane integrity. The effects of catechins and tea anthocyanin extract in attenuating the *t*-BHP induced oxidative stress was evaluated by determining by comparing the release of LDH to the

medium cells treated with the tea constituents and *t*-BHP vs cells treated only with *t*-BHP.

Lactate dehydrogenase leakage is recognized as a sensitive measure of plasma membrane damage and cytotoxicity caused by reactive oxygen species (ROS) which can eventually lead to cell death. Pre-treatment of the cells for 30 min with the catechins (-)-EC alone or before addition of 500 μ M *t*-BHP and incubation for 6 h resulted in lowering of LDH leakage to 4.71% ($p < 0.001$), 1.57% ($p < 0.001$) and 0.16% ($p < 0.001$) for 10, 25 or 50 μ M (-)-EC, respectively without *t*-BHP compared to 500 μ M *t*-BHP while the cells that were oxidatively stressed after pre-treatment showed a reduction of LDH leakage to 14.6% ($p > 0.05$), 9.1% ($p > 0.05$) and 5.2% ($p < 0.05$) compared to 500 μ M *t*-BHP (Figure 1). Treatment of cells with 500 μ M *t*-BHP for 6 h caused a rapid release of LDH into the cytosol which was attenuated in a dose dependent manner by pre-treatment of cells with increasing Epicatechin levels. One way ANOVA results showed significance ($p < 0.001$) between the various concentrations of epicatechin used showing that EC did protect the cells under oxidative stress. To examine the attenuation of LDH leakage by EGCG, a similar treatment was carried out. It was observed that LDH leakage in the controls (without 500 μ M *t*-BHP) was lowered in a dose dependent manner from 2.58% to 1.67% ($p < 0.01$), 1.11% ($p < 0.01$) and 0.11% ($p < 0.01$) while for the cells that were treated (with 500 μ M *t*-BHP) LDH leakage was lowered from 15.1% to 12.2% ($p > 0.05$), 10.9% ($p > 0.05$) and 2.8% ($p < 0.01$), respectively compared to *t*-BHP treatment (Figure 2). One way ANOVA results showed significant cytoprotective effects ($p < 0.001$) against oxidative stress between the various concentrations of EGCG tested.

In an assay to evaluate the effect of tea anthocyanin extract on LDH leakage in HEK 293 cells, the first group of cells was treated with anthocyanins only and the second group was pre-treated with increasing anthocyanin concentrations for 30 min before induction of oxidative stress with *t*-BHP (500 μ M) (Figure 3). LDH leakage in the controls (without 500 μ M *t*-BHP) was lowered to 9.35% ($p < 0.001$), 3.7% ($p < 0.001$), 2.0% ($p < 0.000$) and 0.23% ($p < 0.0001$) for 10, 25, 50 and 100 μ M, respectively. In the oxidatively stressed cells (with 500 μ M *t*-BHP), LDH leakage was dose dependently lowered from 20.63% to 16.79% ($p > 0.05$), 13.92% ($p < 0.01$), 7.23% ($p < 0.001$) and 2.41% ($p < 0.0001$) for the same tea anthocyanin extract concentrations, respectively. ANOVA results showed significant differences ($p < 0.001$) between the various concentrations of tea anthocyanin extract used. These results demonstrated that the anthocyanins were effective cytoprotectants against oxidative stress.

Effect of catechins and tea anthocyanins on intracellular GSH content

The effect of catechins and tea anthocyanin extract on

reduced glutathione (GSH) content of HEK 293 cells was also determined in this study. GSH is an important high capacity cellular antioxidant, widely distributed among living cells. It is also involved in many biological functions such as amelioration of *t*-BHP-induced rat hepatotoxicity (Hwang et al., 2005). though not statistically significant, EGCG marginally increased cellular GSH content in *t*-BHP treated cells from 0.043 GSH nmol/mg protein for *t*-BHP only to 0.058, 0.067 and 0.074 GSH nmol/mg protein, respectively for 10, 25 and 50 μ M EGCG (Figure 4). Tea anthocyanin extract at concentrations of 10, 50 and 100 μ M similarly, marginally increased GSH concentration in the *t*-BHP treated cells. (Figure 5) but not significantly. The data presented for EGCG and tea anthocyanin extract were obtained from the mean of two independent experiments that were repeated with similar results.

DISCUSSION

Interest in the protective function of naturally occurring antioxidants in biological systems has risen in the recent past. For a long time, tea has been used as a healthy beverage owing to its pharmacological properties which are largely ascribed to the catechins contained. Catechins are known to quench free radicals, a characteristic associated with the hydroxyl groups in the B-ring. Evidence has shown that increased hydroxylation increases the antioxidant power of the catechins with EGCG in green tea being the most potent of all (Saffari and Sadrzadeh, 2004). Epigallocatechin-3-gallate (EGCG) is the most abundant compound found in green tea and has been associated with various health benefits as nutritional supplements for various ailments. The potential health benefits ascribed to EGCG include cancer chemoprevention (Hsuuw and Chan, 2007), antioxidant (Fu and Koo, 2006), improving cardiovascular health (Hirai et al., 2007) and neuroprotective (Mandel et al., 2004) among others. However, increasing evidence has also shown that EGCG is a pro-oxidant in higher concentrations and enhances production of ROS which subsequently lead to cell death (Bandeled and Osheroff, 2008; Lambert et al., 2010; Rohde et al., 2011).

Recently, anthocyanin containing tea cultivars have been developed and characterized (Kerio et al., 2011). The anthocyanins/anthocyanidins identified in the tea cultivars were cyanidin-3-O-galactoside, cyanidin-3-O-glucoside, cyanidin, delphinidin, peonidin, pelargonidin and malvidin (Kerio et al., 2011) but the most predominant in the TRFK 306 cultivars is malvidin. Anthocyanins are pigmented plant polyphenols, whose colour in a plant depends on the pH of the cell sap. One purpose of this preliminary study was to compare the antioxidant potency of a mixed extract of tea anthocyanins with the two pure tea catechins, EC and EGCG. All three treatments attenuated the oxidative stress induced by 500 μ M *t*-BHP in HEK 293 cells by




Figure 1. The effect of pre-incubating HEK 293 cells with 0-50 μM epicatechin for 30 min on LDH leakage following incubation with or without *t*-BHP (500 μM) for 6 h. DMSO is the vehicle control (0 μM epicatechin); C1, C2 and C3 contain 10, 25 or 50 μM epicatechin (EC), respectively. Each column represents the mean and SD of three independent experiments. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ compared with *t*-BHP treatment alone (student's *t*-test).




Figure 2. The effect of pre-incubating HEK 293 cells with 0-50 μM Epigallocatechin gallate (EGCG) for 30 min on LDH leakage following incubation with or without *t*-BHP (500 μM) for 6 h. DMSO is the vehicle control (0 μM epicatechin); C1, C2 and C3 contain 10 μM , 25 μM or 50 μM EGCG, respectively. Each column represents the mean and SD of three independent experiments. ** $p < 0.01$ and *** $p < 0.001$ compared with *t*-BHP treatment alone.

reducing LDH leakage, an assay for cell damage. The integrity of cellular membranes is critical for normal cell function. Peroxidative decomposition of the membranes leads to impairment of normal cellular functions. Others have shown that *t*-BHP enhances lipid membrane peroxidation that contributes to cytotoxicity of HEK 293 cells (Kim et al., 2011; Hwang et al., 2005; Wang et al., 2000).

The results of this study revealed that catechins (EC and EGCG) as well as tea anthocyanin extract were able

to partially protect the cells from the oxidative damage caused by *t*-BHP (500 μM). The effectiveness of the anthocyanidin extract might be attributed to its major constituent, malvidin. Malvidin is a dimethylated compound that is non-polar in nature. It is postulated that this anthocyanidin will have increased partitioning into the lipid phase of the membrane thereby increasing the antioxidant capacity of this tea anthocyanin. In addition, anthocyanin antioxidants are unique in their mode of action which may be attributed to their ability for electron




Figure 3. The effect of pre-incubating HEK 293 cells with 0-100 µM tea anthocyanin extract for 30 min on LDH leakage following incubation with or without *t*-BHP (500 µM) for 6 h. DMSO is the vehicle control (0 µM tea anthocyanin extract); A1, A2 and A3 contain 10, 25, 50 or 100 µM tea anthocyanin extract, respectively. Each column represents the mean and SD of three independent experiments. ** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$, compared with *t*-BHP treatment alone.




Figure 4. Effect of EGCG on the concentration of GSH in the treated cells and on the depletion of GSH induced by *t*-BHP (500 µM, 6 h) in HEK 293 cells. C1, C2 and C3 contain 10, 25 or 50 µM EGCG, respectively. Each column represents the mean of two independent experiments.

delocalization and to form resonating structures following changes in pH (Bagchi et al., 2006). Anthocyanins have been shown to be novel antioxidants and potent inhibitors of lipid peroxidation in comparison with other classic antioxidants. For example, grape seed proanthocyanidin extract was found to have superior antioxidant efficacy compared to vitamins C, E and β -carotene (Bagchi et al., 2003).

GSH has been established as a very important endogenous molecule for protection against the cytotoxicity of reactive electrophilic metabolites by converting them to covalent S-glutathione conjugates (Habig et al., 1974). GSH is also a very important antioxidant, essential for the maintenance and balance of thiol (-SH) groups on intracellular proteins. Antioxidants

(vitamin C, catechins and other polyphenols) have been shown to increase the GSH pool and decrease GSSG content as well as attenuate hydroperoxide induced oxidative stress by stimulating the activity of two enzymes involved in GSH/GSSG balance that is glutathione peroxidase (GPx) and glutathione reductase (GRx) (Wijeratne et al., 2005). Increase in GSH content and decrease in cellular hydroperoxides (oxidants) are related phenomena. Dietary antioxidants for example catechins in tea and lately anthocyanins in the same plant, assist the cells to quench the oxidant evaluated in this study caused by *t*-BHP, complementing cellular antioxidants such as GSH and preventing their depletion. In cells, antioxidants often cause cyclic stimulation of glutathione peroxidase (GPx) and glutathione reductase (GRx), the




Figure 5. Effect of tea anthocyanin extract on the concentration of GSH in the treated cells and on the depletion of GSH induced by *t*-BHP (500 μ M, 6 h) in HEK 293 cells. A1, A2 and A3 contain 10 μ M, 50 μ M or 100 μ M tea anthocyanin extract. Each column represents the mean of two independent experiments.

latter which regenerates GSH from GSSG to be used in other antioxidant processes in the mitochondria or they improve mitochondrial function by detoxifying hydroperoxides (Shih et al., 2007). For Example, EGCG has been shown to modulate GSH metabolism in rat astrocytes (Ahmed et al., 2002). A key finding in this study is that anthocyanins and EGCG not only maintain a good redox status in basal conditions for GSH, but also do so in the *t*-BHP treated cells that have undergone oxidative insult. Our *t*-BHP treatment resulted in depletion of GSH possibly as a result of increased the levels of cellular hydroperoxides and high amounts of ROS responsible for oxidative stress (Alia et al., 2005). The 500 μ M *t*-BHP used in this study caused oxidation of cellular GSH and subsequently resulted in an increase in the levels of GSSG, an inhibitor of GPx that can saturate GRx. (Yang et al., 2006). *t*-BHP-induced ROS formation affects GRx activity indirectly by oxidation of the pyridine nucleotide NADPH, a limiting cofactor for GRx. On the positive side, anthocyanins and catechins behave as electron donors to the mitochondrial electron transport chain or as mitochondrial respiring substrates that support the reduction of GSSG formed during oxidative stress (Weissel et al., 2006). The ability of dietary antioxidants including catechins and anthocyanins to function as direct free radical scavengers is related to their electron donating ability which is a direct function of the number of hydroxyl groups in the B-ring and the structural orientation of the compound (Castaneda-Ovando et al., 2009). Studies on the chemistry of EGCG as an antioxidant have shown that the trihydroxyphenyl B-Ring is the more active site of the antioxidant reaction than is the galloyl moiety (Nagle et al., 2006). In anthocyanins, the ring orientation has been found to determine the ease by which a hydrogen atom from a hydroxyl group can be donated to a free radical and its ability to support a free electron by delocalization of its own electrons (Castaneda-Ovando et al., 2009).

A study on *t*-BHP induced hepatotoxicity in rats demonstrated the protective effects of anthocyanins extracted from dried hibiscus flowers, primarily delphinidin-3-glucoside and cyanidin-3-glucoside (Wang et al., 2000) which decreased cytotoxicity (LDH release) and lipid peroxidation at concentrations of 0.1 and 0.2 mg/mL. Similar attenuation of oxidative stress-mediated toxicity was found *in vivo* after the oral administration of 100 or 200 mg/kg anthocyanins to rats for 5 days before a single dose of 0.2 mmol/kg of *t*-BHP. Although the catechin EGCG, also has cytoprotective properties it can also causes oxidative stress, as demonstrated by the effect of EGCG on GSH metabolism in cultured rat astrocytes (Ahmed et al., 2002). Similarly, several other studies have shown that EGCG, at high concentrations has pro-oxidative activities. In pancreatic beta cells, treatment with EGCG reduced cell viability and increased apoptotic cell death, H_2O_2 and ROS production, particularly at 50 and 100 μ M although enhanced cell death was found at concentrations as low as 5 μ M after 48 h (Suh et al., 2010). Recently, there was a case reported in Denmark of liver toxicity in an individual who drank 4-6 cups of green tea per day for six months (Rohde et al., 2011) which was ascribed to EGCG, the major catechin in green tea. Cell culture studies have revealed that EGCG could inhibit intercellular communication via gap junctions in normal rat liver epithelial cells (Kang et al., 2008). Indeed, EGCG has been shown to poison both isomers of topoisomerase II by a redox-dependent mechanism similar to that for 1, 4-benzoquinone (Bandeled and Osherof, 2008).

The lowered levels of LDH leakage and apparent increase in cellular GSH content (due to attenuated depletion) in cells treated with the tea anthocyanin extract in this study reveal potential benefits of these compounds in the normal body cells. The tea anthocyanin extract was able to partially attenuate *t*-BHP-induced oxidative stress in the HEK 293 cells, effectively demonstrating the

cytoprotective properties of these antioxidants in cultured cells.

Conclusion

Oxidative stress results from an excess of intercellular ROS and RNS free radicals which are associated with several degenerative diseases, including cancer and chronic inflammation (Halliwell, 2002; Wang and Stoner, 2008; Tang et al., 2007). Therefore, it seems reasonable that consumption of reasonable amounts of anthocyanin rich teas, containing both anthocyanins and catechins, is likely to attenuate the adverse effects of oxidative stress in the body by decreasing oxidation of essential cellular proteins, lipids and nucleic acids while sparing GSH depletion. In this study, it is shown that the tea anthocyanin extract at the highest concentrations tested (100 μ M), did not cause any obvious oxidative stress on the cells. This shows the potential of the tea anthocyanins for cytoprotection against oxidative stressors. It should however be stressed that although regardless of the class of chemical antioxidants selected for this purpose, ingestion of high concentrations over time is likely to be associated with negative effects, as has been reported in experimental animals and humans for EGCG, particularly when taken on an empty stomach which results in enhanced absorption. Additional work is however required to prove that anthocyanin rich tea cultivars have less potential for negative effects because of the presence of both anthocyanin and catechins antioxidants.

ACKNOWLEDGEMENTS

Authors thank the Graduate Student Exchange Program organized by the Canadian Bureau for International Education (CBIE) funded by the Department of Foreign Affairs and International Trade Canada (DFAIT) for funding this work. We also thank The Tea Research Foundation of Kenya for providing the tea samples used in this study and for permission to publish these results.

REFERENCES

- Alia M, Ramos S, Mateus R, Bravo L, Goya L (2005). Response of the antioxidant defense system to *tert*-butyl hydroperoxide and hydrogen peroxide in a human Hepatoma cell line (HepG2). *J. Biochem. Mol. Toxicol.* 19: 119-128.
- Ahmed I, John A, Vijayasathy C, Robin MA, Raza H (2002). Differential modulation of growth and glutathione metabolism in cultured rat astrocytes by 4-hydroxynonenal and green tea polyphenol, epigallocatechin-3-gallate. *Neurotoxicology*, 23: 289-300.
- Almajano PM, Rosa C, Jimenez L, Gordon H (2008). Antioxidant and antimicrobial activities of tea infusions. *Food Chem.*, 108: 55-63.
- Arlı G, Cau NO (2007). Determination of naturally occurring antioxidant and anti-inflammatory compounds in fresh fruits by HPLC. *Toxicol. Lett.*, 172: S221-S222.
- Bae SH, Suh HJ (2007). Antioxidant activities of five different mulberry cultivars in Korea. *Food Sci. Technol.*, 40: 955-62.
- Bagchi D, Roy S, Patel V, He G (2006). Safety and whole body antioxidant potential of a novel anthocyanins-rich formulation of edible berries. *Mol. Cell. Biochem.*, 281: 197-209.
- Bagchi D, Sen CK, Ray SD, Das DK, Bagchi M, Preuss HG, Vinson JA (2003). Molecular mechanisms of cardioprotection by a novel grape seed proanthocyanidin extract-A review. *Mutation Res.*, 524: 87-89.
- Bandle OJ, Osheroff N (2008). (-)-epigallocatechin gallate, a major constituent of green tea, poisons type II topoisomerases. *Chem. Res. Toxicol.*, 21: 936-943.
- Cabrera C, Artacho R, Gimenez R (2006). Beneficial effects of green tea-A review. *J. Am. Coll. Nutr.*, 25: 79-99.
- Castaneda-Ovando A, Pacheco-Hernandez MDL, Paez-Hernandez ME, Rodriguez JA, Galan-Vidal CA (2009). Chemical studies of anthocyanins: A review. *Food Chem.*, 113: 859-871.
- Choi E, Chang H, Cho J, Hyan S (2007). Cytoprotective effects of anthocyanins against doxorubicin-induced toxicity in H9c2 cardiomyocytes in relation to their antioxidant activities. *Food Chem. Toxicol.*, 45: 1873-1881.
- Dai J, Patel JD, Mumper RJ (2007). Characterization of blackberry extract and its antiproliferative and anti-inflammatory properties. *J. Med. Food*, 10: 258-265.
- Duthie SJ, Gardner PT, Morrice PC, Wood SG, Pirie L, Bestwick CC, Milne L, Duthie GG (2005). DNA stability and lipid peroxidation in vitamin E deficient rats *in vivo* and *in vitro*. Modulation by the dietary anthocyanin cyanidin-3-glycoside. *Eur. J. Nutr.*, 44: 195-203.
- Elisia I, Kitts DD (2008). Anthocyanins inhibit peroxyl radical induced apoptosis in CaCO-2 cells. *Mol. Cell Biochem* 312: 139-145.
- Erdem S, Mendell JR, Sahenk Z (1998). Fate of Schwann cells in CMT1A and HNPP: Evidence for apoptosis. *J. Neur. Exp. Neur.*, 57: 635- 642.
- Franco R, Cidlowski JA (2009). Apoptosis and glutathione; beyond an antioxidant. *Cell Death Diff.*, 16: 1303-1314.
- Fu Y, Koo MW (2006). EGCG protects HT-22 cells against glutamate-induced oxidative stress. *Neur. Res.*, 10: 23-30.
- Gogvadze V, Orrenius S, Zhivotovsky B (2006). Multiple pathways of cytochrome c release from mitochondria in apoptosis. *Biochim. Biophys. Acta.*, 1757: 639-647.
- Gramza A, Khokhar S, Yoko S, Swiglo AG, Hes M, Korczak J. (2006). Antioxidant activity of tea extracts in lipids and correlation with polyphenol content. *Eur. J. Lipid Sci. Tech.*, 108: 351-362.
- Habig WH, Pabst MJ, Jakoby WB (1974). Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J. Biol. Chem.* 249: 7130-7139.
- Haidara K, Morel I, Abalea V, Barre MG, Denizeau F (2002). Mechanism of *tert*-butylhydroperoxide induced apoptosis in rat hepatocytes: involvement of mitochondria and endoplasmic reticulum. *Biochim. Biophys. Acta.*, 1542: 173-185.
- Hafeez BB, Siddiqui IA, Asim M, Malik A, Afaq F, Adhami VM, Saleem M, Din M, Mukhtar H (2008). A dietary anthocyanidin delphinidin induces apoptosis of human prostate cancer PC3 cells *in vitro* and *in vivo*: Involvement of nuclear factor- κ B signalling. *Cancer Res.* 68: 8564-8572.
- Halliwell B, and Whiteman M. (2004). Measuring reactive species and oxidative damage *in vivo* and in cell culture- how should you do it and what do the results mean? *British J. Pharm.*, 142: 231-255.
- Halliwell B (2002). Effect of diet on cancer development: Is oxidative DNA damage a biomarker? *Free Rad. Biol. Med.*, 32: 968-974.
- Heinonen M (2007). Antioxidant activity and antimicrobial effect of berry phenolics- A Finnish perspective. *Mol. Nutr Food Res.* 51: 684-691.
- Hirai M, Hotta Y, Ishikawa N, Wakida Y, Fukuzawa Y, Isobe F, Nakano A, Chiba T, Kawamura N (2007). Protective effects of EGCG or GCG, a green tea catechin epimer against postischemic myocardial dysfunction in guinea-pig hearts. *Life Sci.*, 80: 1020-1032.
- Hsuw Y, Chen W (2007). Epigallocatechin gallate dose-dependently induces apoptosis or necrosis in Human MCF-7 cells. *Ann. N.Y. Acad. Sci.*, 1095: 428-440.
- Hwang JM, Wang CJ, Chou FP, Tseng TH, Hsieh YS, Hsu JD, Chu, CY (2005). Protective effect of baicalin on *tert*-butyl hydroperoxide-induced rat hepatotoxicity. *Arch. Toxicol.*, 79: 102-109.
- Jellinger KA, Bancher C (1998). Neuropathology of Alzheimer's disease: A critical update. *J. Neural Trans.*, 54: 77-95.

- Kang NJ, Lee KM, Kim JH, Lee BK, Kwon JY, Lee KW, Lee HJ (2008). Inhibition of gap junctional intercellular communication by the green tea polyphenol (-)-epigallocatechin gallate in normal rat liver epithelial cells. *J. Agric. Food Chem.*, 56: 10422-10427.
- Karori SM, Ngure RM, Wachira FN, Wanyoko JK, Mwangi JN (2008). Different types of tea products attenuate inflammation induced in *Trypanosoma brucei* infected mice. *Parasit Int.*, 57: 325-333.
- Karori SM, Wachira FN, Wanyoko JK, Ngure RM (2007). Antioxidant capacity of different types of tea products. *Afr. J. Biotech.*, 6: 2287-2296.
- Katalinic V, Milos M, Kulisic T, Jukic M (2006). Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food Chem.*, 94: 550-557.
- Kerio LC, Wachira FN, Wanyoko JK, Rotich MK (2011). Characterization of anthocyanins in Kenyan teas: Extraction and identification. *Food Chem.*, 131(1): 31-38.
- Kim SM, Kang K, Jho EH, Jung YH, Nho CW, Um BH, Pan CH (2011). Hepatoprotective effects of flavonoid glycosides from *Lespedeza cuneata* against oxidative stress induced by *tert*-butyl hydroperoxide. *Phytotherapy Res.* (Epub ahead of print).
- Khokhar S, Owusu RK (2003). Iron binding characteristics of phenolic compounds: some tentative structure-activity relations. *Food Chem.*, 81: 133-140.
- Kong JM, Chia LS, Goh NK, Chia TF, Brouillard R (2003). Analysis and biological activities of anthocyanins. *Phytochemical*, 64: 923-933.
- Korzeniewski C, Callewaert DM (1983). An enzyme-release assay for natural cytotoxicity. *J. Immun. Methods* 64: 313-20.
- Krysko D, Berghe TV, D'Herde K, Vandenaabeele P (2008). Apoptosis and necrosis: Detection, discrimination and phagocytosis. *Methods* 44: 205-221.
- Lambert JD, Kennet MJ, Sang S, Reut KR, Ju J, Yang CS (2010). Hepatotoxicity of high oral dose (-)-epigallocatechin-3-gallate in mice. *Food Chem. Toxicol.*, 48: 409-416.
- Lapenna D, de Goia S, Ciofani G, Mezzetti A, Uchino S, Calafiore AM, Napolitano AM, Di Ilio C, Cucurulo F (1998). Glutathione-related antioxidant defenses in human atherosclerotic plaques. *Circulation* 97: 930.
- Lazze M, Pizzala R, Savio M, Stivala L, Prosperi E, Bianchi RL (2003). Anthocyanins protect against DNA damage induced by *t*-BHP in rat smooth muscle and hepatoma cells. *Mutation Res.*, 535: 103-115.
- Lee V, Chen C, Liao Y, Tzen J, Chang C (2008). Structural determination and DPPH radical scavenging activity of two acylated flavonoids tetraglycosides in oolong tea (*Camellia sinensis*). *Chem. Pharm. Bull.*, 56: 851-853.
- Lee J, Lee HK, Kim CY, Hong Y, Choe CM, You T, Seong GJ (2005). Purified high dose of anthocyanoside oligomer administration improves nocturnal vision and clinical symptoms in myopia subjects. *British J. Nutr.*, 93: 895-899.
- Lee S, Park S, Park JH, Shin, DY, Kim GY, Rya, CH, Choi YH (2009). Induction of apoptosis in human leukaemia U937 cells by anthocyanins through downregulation of Bcl-2 and activation of caspases. *Internatl. J. Onc.*, 34: 1077-1083.
- Leung LK, Su Y, Chen R, Zhang Z, Huang Y, Chen ZY (2001). Theaflavins in black tea and catechins in green tea are equally effective antioxidants. *J. Nutr.* 131: 2248-2251.
- Luczaj W, Skrzydlewska E (2005). Antioxidative properties of black tea - A review. *Prev. Med.* 40: 910-918.
- Mandel S, Weinreb O, Amit T, Youdim MBH (2004). Cell signalling pathways in the neuroprotective actions of the green tea polyphenol (-)-epigallocatechin-3-gallate: Implications for neurodegenerative diseases. *J. Neurochem.*, 88: 1555-1569.
- Maurya PK, Rizvi SI (2008). Protective role of tea catechins on erythrocytes subjected to oxidative stress during human aging. *Nat. Prod. Res.*, pp. 1-8.
- Mazza G. (2007). Anthocyanins and heart health. *Ann. Ist. Super San.* 43: 369-374.
- Muzolf M, Szymusiak H, Anna G, Rietjens MCM, Tyrakowska B (2008). pH-dependent radical scavenging capacity of green tea catechins. *J. Agric. Food Chem.*, 56: 816-832.
- Nagle DG, Ferreira D, Zhou Y (2006). Epigallocatechin-3-gallate (EGCG): Chemical and biomedical perspectives-Molecules of interest. *Phytochemical*, 67: 1849-1855.
- Orak HH (2007). Total antioxidant activities, phenolics, anthocyanins, polyphenol oxidase activities of selected red grape cultivars. *Scientia Hort.*, 111: 235-41.
- Prior RL (2003). Fruits and vegetables in the prevention of cellular oxidative damage. *Am. J. Clin. Nutr.*, 78: 570S-578S.
- Rohde J, Jacobsen C, Kromann-Andersen H (2011). Toxic hepatitis triggered by green tea. *Ugeskr Laeger*, 173: 205-206
- Royr AM, Baliga MS, Katiyar SK (2005). Epigallocatechin-3-gallate induces apoptosis in estrogen receptor-negative human breast carcinoma cells via modulation in protein expression of p53 and Bax and caspase-3 activation. *Mol. Cancer Ther.*, 4: 81 - 90.
- Saffari Y, Sadraze SMH (2004). Green tea metabolite EGCG protects membranes against oxidative damage in vitro. *Life Sci.*, 74: 1513-1518.
- Seely D, Mills EJ, Wu P, Verma S, Guyatt, GH (2005). The effects of green tea consumption on incidence of breast cancer and recurrence of breast cancer: A systematic review and meta-analysis. *Integr. Cancer Ther.* 4: 144 - 155.
- Shih P, Yeh C, Yen G (2007). Anthocyanins induce the activation of Phase II enzymes through the antioxidant response element pathway against oxidative stress induced apoptosis. *J. Agric. Food Chem.*, 55: 9427-9435.
- Shih PH, Yeh CT, Yen GC (2005). Effects of anthocyanidin on the proliferation and induction of apoptosis in human gastric endocarcinoma cells. *Food Chem. Toxicol.*, 55: 1557-1566.
- Suh KS, Chon S, Oh S, Kim JW, Kim YS, Woo JT (2010). Pro-oxidative effects of green tea polyphenol (-)-epigallocatechin-3-gallate on the HIT-T15 pancreatic beta cell line. *Cell Biol. Toxicol.*, 26: 189-199.
- Tang Y, Zhao DY, Elliott S, Zhao W, Curriel TJ, Beckman BS, Burrow ME (2007). Epigallocatechin-3-gallate induces growth inhibition and apoptosis in human breast cancer cells through surviving suppression. *Internat. J. Oncol.*, 31: 705-711.
- Tang SZ, Kerry JP, Sheehan D, Buckley DJ (2002). Antioxidative mechanisms of tea catechins in chicken meat systems. *Food Chem.*, 76: 45-51.
- Tarozzia A, Morroni F, Hrelia S, Ageloni C, Machesi A, Cantelli FG, Hrelia P. (2007). Neuroprotective effects of anthocyanins and their *in vivo* metabolites in 5H-SY5Y cells. *Neurosci. Lett.*, 424: 36-40.
- Tea Board of Kenya (2010). Tea industry performance in 2009. The Tea Board of Kenya.
- Terahara N, Takeda Y, Nesumi A, Honda T (2001). Anthocyanins from red-flower tea (Benibana-cha) *Camellia sinensis*. *Phytochem.* 56: 359-361.
- Viskalis P, Rubiskiene M, Jasutiene I, Sarkinas A, Daubaras R, Cesoniene L (2009). Anthocyanins, antioxidative and antimicrobial properties of American Cranberry (*Vaccinium macrocarpon* Ait) and their press cakes. *J. Food Sci.*, 74: 157-161.
- Wang C, Wang J, Lin W, Chu C, Chou F, Tseng T (2000). Protective effects of Hibiscus anthocyanins against *tert*-butyl hydroperoxide-induced hepatotoxicity in rats. *Food Chem. Toxicol.*, 38: 411-416.
- Wang L, Stoner GD (2008). Anthocyanins and their role in cancer prevention. *Cancer Lett.*, 269: 281-290.
- Weissel T, Baum M, Eisenbrand G, Dietrich H, Will F, Stockis JP, Kulling S, Rufer C, Johannes C, Jackowski C (2006). An anthocyanins/phenolic-rich fruit juice reduces oxidative DNA damage and increases glutathione levels in healthy probands. *Biotechnol. J.*, 1: 388-397.
- Wijeratne SS, Cuppet SL, Schlegel V (2005). Hydrogen peroxide induced oxidative stress damage and antioxidant enzyme response in Caco-2 human colon cells. *J. Agric Food Chem.*, 53: 8768-8774.
- Yang MS, Chan HW, Yu LC (2006). Glutathione peroxidase and glutathione reductase activities are partially responsible for determining the susceptibility of cells to oxidative stress. *Toxicology*, 226: 126-130.
- Yao K, Ye P, Zhang L, Tang X, Zhang Y (2007). Epigallocatechin gallate protects against oxidative-stress induced mitochondria-dependent apoptosis in human lens epithelial cells. *Mol. Vis.*, 14: 217-223.