

## Full Length Research Paper

## Assessment of Ochratoxin A intake due to consumption of coffee and cocoa derivatives marketed in Abidjan (Côte d'Ivoire)

Pierre Manda\*, Aholia Jean Baptiste Adepo, Jean Verdier Ngbé and Djédjé Sébastien Dano

Laboratoire de Toxicologie et Hygiène Agro-industrielle, UFR Sciences Pharmaceutiques et Biologiques, Université Félix Houphouët Boigny, B.P.V. 34 Abidjan Côte d'Ivoire.

Received 8 August, 2016; Accepted 30 September, 2016

Ochratoxin A is a mycotoxin produced by moulds belonging to the *Aspergillus* and *Penicillium* species, and is a common contaminant of various food products. Ochratoxin A is known for its nephrotoxic and carcinogenic effects; protective actions to protect consumer health must be established. The aim of this study was to assess the level of contamination of manufactured coffee and cocoa-based products marketed in Abidjan to estimate ochratoxin A intake related to the consumption of these goods. We quantified ochratoxin A in 87 samples by high-performance liquid chromatography with fluorescence detection. Coffee samples contained an average of  $4.0 \pm 3.4$   $\mu\text{g}/\text{kg}$  ochratoxin A and cocoa-based industrial products  $0.9 \pm 0.6$   $\mu\text{g}/\text{kg}$ . OTA intakes were assessed using data obtained from the analysis and the estimated Ivorian adult daily consumptions. The average estimated intake for the Ivorian population is 0.15 ng/kg bw/d for coffee and 0.02 ng/kg bw/d for cocoa, lower than the tolerable daily intakes of 5 and 14 ng/kg bw/d, respectively set by the European Commission and World Health Organization. However, in view of the toxicity of OTA, it is imperative to establish legislation on mycotoxin in Côte d'Ivoire

**Key words:** Product derivatives, coffee, cocoa, ochratoxin A.

### INTRODUCTION

Ochratoxin A (OTA) is a mycotoxin produced by moulds belonging to the *Aspergillus* and *Penicillium* species. OTA is a common contaminant of cereals, dried fruits, coffee and cocoa, as well as manufactured food products such as chocolate, wine, beer, and bread (Miraglia and Brera, 2002; Sangare et al., 2006). OTA-contaminated foods may be pathogenic to humans and animals (Rutqvist et al., 1978). OTA appears to be involved in

Balkan endemic nephropathy (Pfohl-Leszkowicz et al., 2002; Castegnaro et al., 2006), and is genotoxic, immunotoxic, and carcinogenic (IARC, 1993; Pfohl-Leszkowicz and Manderville, 2007; 2012; Manderville and Pfohl-Leszkowicz, 2008; Mantle et al., 2010). The health risk due to OTA contamination of food necessitates establishing specific actions to protect consumer health worldwide. In Côte d'Ivoire, there is no

\*Corresponding author. E-mail: mandapierre@yahoo.fr. Tel: 00225 05698726.

regulation on mycotoxin and particularly ochratoxine A in foodstuffs. This exported cocoa ochratoxin A contamination assessment target is to answer the legal dispositions and regulations in exporting countries. In Europe, regulations (EU 1881/2006 and 105/2010) have set the maximal limits for some contaminants in foodstuffs, e.g. the limits of OTA. In 2006, the European Food Safety Authority (EFSA) established a weekly tolerable OTA intake of 120 ng/kg of body weight (bw). The European Commission (EC, 2010) stated that it does not appear to be necessary to establish a maximal OTA content for cocoa and cocoa-derived products to protect consumer health because these foodstuffs do not contribute significantly to OTA exposure and high OTA contents have rarely been detected in these food products. To guarantee the sanitary quality of cocoa, it is important for the cocoa producing countries to develop post-harvesting treatment guidelines to minimize OTA-contamination in cocoa. Indeed, analysis of cocoa bean samples from different producing countries revealed varying levels of OTA contamination between regions and even within the same area (Amézquetà et al., 2005; Dongo et al., 2008; Copetti et al., 2010). Côte d'Ivoire is the largest cocoa-producing country (1,450,000 tons in 2012) and the 8<sup>th</sup> largest coffee-producing country in the world. Most of the Ivorian production is exported to Europe and North America; only a small amount is kept for home consumption. The aim of this study was to assess the level of contamination of coffee and cocoa-based products marketed along with the estimation of the dietary intake in the Ivorian adults.

## MATERIALS AND METHODS

### Collection of samples

Altogether, 87 samples, including 50 cocoa-based products (cocoa powder 100%; chocolate drink, chocolate bars) and 37 coffee samples (ground coffee; instant coffee), were collected from five supermarkets in Abidjan (Table 1) in accordance with EC N°401/2006 of 23 February 2006 specifying the methods of sampling and analysis for the official control of the levels of mycotoxins in foodstuffs (method for sampling coffee and coffee products: sampling at retail stage). Coffee and cocoa-derived product samples were collected during three distinct harvest seasons (January, March, and July). The coffee content of the coffee samples was greater than 60%, and the cocoa content of the cocoa-based products was greater than 25% (CODEX STAN 87-1981, rev. 1 – 2003).

### Reagents and standards

A standard 100 µg/L solution of OTA, phosphate buffered saline (PBS), and an OTA-specific immunoaffinity column were obtained from R-Biopharm (Lyon, France). Acetonitrile and methanol were of analytical quality (Carlo Erba réactifs Val de Reuil, France). Glacial acetic acid and sodium bicarbonate of analytical quality were obtained from Sigma-Aldrich (St. Louis, MO, USA). Reference coffee (T 1771) was obtained from Fapas (York, United Kingdom).

### Determination of OTA-validation

OTA extraction was performed under alkaline conditions as previously described (Amézquetà et al., 2005). Three different samples of each matrix spiked with OTA and were analysed on the same day. The limit of detection (LOD) and limit of quantification (LOQ) were 0.05 and 0.2 µg/kg for the cocoa matrix and 0.1 µg/kg for the coffee, respectively. The average recoveries were 95.23±1.2 and 95.22, respectively for cocoa and coffee. Recoveries were consistent with relative standard deviations (RSDs) of less than 3%, demonstrating the precision of the analytical procedure. The method is valid according to Directive 2002/26/CE, which indicates that recoveries of 70-110% are acceptable for concentrations of OTA between 1 and 10 µg/kg. Solutions containing 1, 5, 10, 20 and 40 µg/kg OTA were measured to construct dose-response curves for OTA analysis with a linearity coefficient ( $r^2$ ) of 0.9995 and 0.9998 respectively for cocoa and coffee.

### Estimation of OTA intake

OTA intake was estimated from data of the International Coffee and Cocoa Organization (2014) related to the consumption of coffee and cocoa derivatives. The estimated mean cocoa consumption in Côte d'Ivoire is 0.048 kg of chocolate per capita per year, and that for coffee 0.13 kg per capita per year. The dietary intake has been calculated for a 70 kg adult. OTA intake was calculated according to the following formula (WHO, 2003):

OTA intake =  $T_{OTA} \times C$  ( $T_{OTA}$ : OTA intake content found in food in µg/kg and C: individual daily food consumption in kg/day).

### Statistical analysis

Data are expressed as the mean ± standard error of the mean (SEM). The occurrence of OTA in samples was compared using a Wilcoxon matched-pair test, and statistical significance was established as  $p < 0.05$ .

## RESULTS AND DISCUSSION

### Amount of OTA in the coffee samples

OTA was detected in 30 of 37 (81%) coffee samples. The average OTA concentration of contaminated coffee samples was  $0.92 \pm 0.6$  µg/kg. The average concentration of OTA was  $6.4 \pm 3.8$  µg/kg in ground coffee and  $0.8 \pm 0.8$  µg/kg in instant coffee. Twelve of 21 ground coffee samples contained less than 5 µg/kg OTA. All instant coffee samples were in compliance with EC regulation No1881/2006, that is,  $< 10$  µg/kg). Coffee was the most highly contaminated substance in our study with levels occasionally reaching 20 µg/kg (Table 1). Since the first detection of OTA in green coffee by Levi et al., 1974) other studies have confirmed the presence of this mycotoxin in various coffee-based products, such as roasted coffee and instant coffee (Leoni et al., 2000). OTA has been identified in manufactured coffee from various countries, such as Vietnam (Leong et al., 2007), France (NARI, 2004), India (Gopinandhan et al., 2008), the Netherlands (Van der Stegen et al., 2001), and the Czech Republic (Malir et al., 2014). The reported average

**Table 1.** Level of OTA contamination (in  $\mu\text{g}/\text{kg}$ ) of coffee and cocoa-based samples marketed in Côte d'Ivoire

Products	Total	Samples		OTA ( $\mu\text{g}/\text{kg}$ )	
		Positive (%)	< 1 $\mu\text{g}/\text{kg}$	Mean	Range
Cocoa powder	19	19 (100)	9	1.34 $\pm$ 0.9	0.14-6.3
Chocolate bars	11	10 (90)	9	0.56 $\pm$ 0.48	LD-2.4
Chocolate drink	20	20 (100)	14	0.73 $\pm$ 0.36	0.17-2.1
Cocoa (local)	16	6 (37)		0.75 $\pm$ 0.45	0.09-3.0
Cocoa (imported)	34	33 (97)		1.0 $\pm$ 0.7	LD-6.3
<b>Total cocoa</b>	<b>50</b>	<b>49 (98)</b>	<b>32</b>	<b>0.92 <math>\pm</math> 0.6</b>	<b>LD-6.3</b>
Ground coffee	21	21 (100)	n<5 $\mu\text{g}/\text{kg}$ * 12	6.4 $\pm$ 3.8	0.6 -20.3
Instant coffee	16	9 (56)	n<10 $\mu\text{g}/\text{kg}$ * 16	0.80 $\pm$ 0.8	LD-3.7
Coffee (local)	20	20 (100)		5.5 $\pm$ 3.5	0.9-20.3
Coffee (imported)	17	10 (58)		2.2 $\pm$ 2.6	LD-11.2
<b>Total coffee</b>	<b>37</b>	<b>30 (81.08)</b>		<b>4.0 <math>\pm</math> 3.4</b>	<b>LD-20.3</b>

\*EC 1881/2006 and 105/2010 regulations set the maximal limits for some contaminants in foodstuffs.

levels were variable. Almeida et al., (2007) detected OTA in 81 of 82 instant coffee samples identified in Brazil, with contamination levels varying from 0.17 to 6.3  $\mu\text{g}/\text{kg}$ . In Spain, Coronel et al. (2011) reported an average of 2.2  $\pm$  0.8  $\mu\text{g}/\text{kg}$  (1.21 to 4.21  $\mu\text{g}/\text{kg}$ ) for 79 roasted coffee samples. The average contamination level of instant coffee samples was lower than that reported by Pittet et al. (1996) (1.1  $\mu\text{g}/\text{kg}$ ) and of Prado et al. (2000) (1.7  $\mu\text{g}/\text{kg}$ ).

Twenty of the coffee products were manufactured in Côte d'Ivoire and 17 were imported. The average OTA content of the Ivoirian coffee was 5.5  $\pm$  3.5 whereas that of the imported coffee was 2.2  $\pm$  2.6  $\mu\text{g}/\text{kg}$ . This difference was significantly different ( $p=0.035$ ; Table 1).

Because coffee roasting is not a heavy manufacturing process, there are several local coffee merchants of ground coffee that represent the "under grade or fragments" coffee category; high contamination was observed in this category. Dembélé et al., (2008) reported the average OTA content of 100 coffee samples, depending on the grade collected, from the two seaports of Côte d'Ivoire as follows: grade 1: 7  $\mu\text{g}/\text{kg}$ ; grade 2: 9.7  $\mu\text{g}/\text{kg}$ ; grade 3: 8.5  $\mu\text{g}/\text{kg}$ , and fragments: 40  $\mu\text{g}/\text{kg}$ . The authors established a linear correlation between the merchantability criteria and the content of OTA in coffee. The higher category (grade I or II) coffee is generally exported. Instant coffee which requires a more sophisticated industrial process is made at the local level by only one "multinational" manufacturer with higher quality standards for raw materials. Instant coffee samples did not generally exceed 10  $\mu\text{g}/\text{kg}$ . Several authors reported that coffee manufacturing operations reduce OTA content. Coffee roasting has been the most studied.

Roasting leads to a significant reduction of OTA, between 40 and 90% of the initial content (Van der Stegen et al., 2001). The detection of mycotoxigenic fungi in the postharvest processing by PCR-DGGE could also be a promising tool in order to reduce OTA production in coffee beans (Durand et al., 2013; Nganou et al., 2012)

#### Amount of OTA in the cocoa samples

OTA was detected in 49 out of the 50 (98%) cocoa-based products. The average OTA concentration of the cocoa-based samples was 0.92 $\pm$ 0.6  $\mu\text{g}/\text{kg}$ . Of the 50 cocoa-based samples, 32 exhibited OTA concentrations of less than 1  $\mu\text{g}/\text{kg}$ . Cocoa powder had the highest average OTA content (1.3  $\pm$  0.9  $\mu\text{g}/\text{kg}$ ) followed by chocolate bars (0.7  $\pm$  0.4  $\mu\text{g}/\text{kg}$ ) and chocolate drink with 15% cocoa (0.56  $\pm$  0.48  $\mu\text{g}/\text{kg}$ ) (Table 1). A high prevalence of OTA in chocolate products has been reported in the last decade worldwide. The results obtained for OTA levels of cocoa powder samples in our study are similar to those reported in the literature. OTA has been detected in 92 to 100% of evaluated cocoa powder, with average levels between 0.18 and 7.8  $\mu\text{g}/\text{kg}$  (Gilmour and Lindblom, 2008; Miraglia and Brera, 2002; Turcotte and Scott, 2011; Copetti et al., 2012). The prevalence of OTA reported in the literature for chocolate is also high [60 to 100% in chocolate (0.08 to 0.88  $\mu\text{g}/\text{kg}$ )] (Gilmour and Lindblom 2008; Turcotte and Scott, 2011).

Thirty-four of the 50 cocoa samples were imported, coming from France (25 samples), Ghana (5 samples), Spain (3 samples), and England (1 sample). The average OTA content of the imported cocoa-based products and

**Table 2.** Food intake of OTA estimated from quantity of food consumed per Ivorian adult

		Coffee	Cocoa
Consumed quantity	(g/day/individual)	2.6	0.13
Estimated intake (mean)	(ng/day/kg of bw)	0.15	0.0017
TDI	5 ng/kg of bw (Health Canada) 14.3 ng/kg bw/d (JEFCA 1995; 2001)		

those manufactured in Côte d'Ivoire was  $1.0 \pm 0.7$  and  $0.8 \pm 0.5$   $\mu\text{g}/\text{kg}$ , respectively. There was no significant difference between the level of OTA contamination in imported and locally manufactured products (Table 1). The high OTA content observed in the Ivorian samples evaluated in our study contrast with the low level of contamination previously reported for Ivorian products. Indeed, Coulibaly et al. (2013) analysed 1895 samples of cocoa collected from all the producing regions of Côte d'Ivoire between December 2013 and June 2014. The overall average level of OTA contamination was  $1.4 \pm 1.3$   $\mu\text{g}/\text{kg}$ . The average was  $0.64 \pm 0.53$   $\mu\text{g}/\text{kg}$  for cocoa for export (grade I and II), and that for non-exportable cocoa (under grade) was  $1.9 \pm 1.9$   $\mu\text{g}/\text{kg}$ . Several authors reported that cocoa manufacturing operations reduce OTA content. The operations of roasting, shelling, and the addition of additives were identified as operations that may reduce OTA content by 90% (Copetti et al., 2012).

### OTA intake

The risks to public health related to the presence of a toxin in food are evaluated by estimating the quantity of the substance actually ingested and comparing this quantity with the toxicity reference values (TRV) established for the same substance. These metrics include the acceptable daily intake (ADI), the provisional tolerable weekly intake (PTWI) or safety limit (SL). Thus, the High Council of Public Hygiene of France (HCPHF, 1999) and the Scientific Committee on Food (SCF, 1998) established a tolerable daily intake (TDI) of 5 ng/kg bw/d for OTA in 1999 and 1998, respectively. Similarly, the Joint FAO/WHO Expert Committee on Food Additives (JECFA, 2001) proposed a TDI of 14.3 ng/kg bw/d in 1995 and then again in 2001. The average estimated intake for the Ivorian population is 0.14 ng/kg bw/d for coffee and 0.0017 ng/kg/bw d for cocoa (Table 2). These values are lower than either of the recommended ADI (5 and 14 ng/kg bw/d). Coulibaly et al. (2013) estimated a dietary intake of OTA, from the consumption of cocoa-based products, for the Ivorian consumer of  $0.004 \pm 0.003$  ng/kg bw/d for exportable beans and  $11.0 \pm 10.0$  ng/kg bw/d for non-exportable beans. A study performed in France by the National Agronomic Research Institute (2004) also provided data on the levels of OTA contamination of coffee samples (0.04  $\mu\text{g}/\text{kg}$ ) and cocoa-

based (chocolate: 0.25  $\mu\text{g}/\text{kg}$ ) foods. These values correspond to the very low levels of OTA exposure of 0.076 and 0.011 ng/kg bw/d, respectively, for adults (15 years and more) for coffee and cocoa.

### Conclusion

Our results show that the prevalence of OTA contamination of coffee and cocoa-based products marketed in Côte d'Ivoire is high. OTA is detectable in greater than 80% of products with means levels of 4.0  $\mu\text{g}/\text{kg}$  in coffee and 0.9  $\mu\text{g}/\text{kg}$  in cocoa-based products. OTA intake estimated from the consumption of these products is, however, very low. In light of the toxicity of OTA, it is imperative to foster best practices for harvesting, pod breaking, drying, and storing cocoa beans to minimize their contamination. In addition, health authorities must enforce controls regarding foodstuffs.

### Conflict of Interests

The authors have not declared any conflict of interests.

### REFERENCES

- Almeida AP, Alaburda J, Shundo L, Ruvieri V, Navas SA, Lamardo LCA, Sabino M (2007). ochratoxin A in brazilian instant coffee. *Braz. J. Microbiol.* 38(2):300-303.
- Amézquetá S, González-Peñas E, Murillo M, Lopez de Cerain AL (2005). Occurrence of ochratoxin A in cocoa beans: effect of shelling. *Food Addit. Contam.* 22(6):590-595.
- Castegnaro M, Canadas D, Vrabcheva T, Petkova-Bocharova T, Chernozemsky IN, Pfohl-Leszkiwicz A (2006). Balkan endemic nephropathy: role of ochratoxins A through biomarkers. *Mol. Nutr. Food Res.* 50(6):519-529.
- Copetti MV, Iamanaka BT, Pereira JL, Lemes DP, Nakano F, Taniwakib MH (2012). Co-occurrence of ochratoxin A and aflatoxins in chocolate marketed in Brazil. *Food Control* 26(1):36-41.
- Copetti MV, Pereira JL, Iamanaka BT, Pitt JL, Taniwaki MH (2010). Ochratoxigenic fungi and ochratoxin A in cocoa during farm processing. *Int. J. Food Microbiol.* 143(1-2):67-70.
- Coronel MB, Marin S, Cano G, Ramos AJ, Sanchis V (2011). Ochratoxin A in Spanish retail ground roasted coffee: occurrence and assessment of the exposure in Catalonia. *Food Control* 22(3-4):414-419.
- Coulibaly A, Godi HMB, Ardjouma D, Kouassi MB, Abba T (2013). Cocoa beans and cocoa derivatives from Côte d'Ivoire: investigating ochratoxin A level and assessing dietary intake adults. *Sustain. Agric. Res.* 2(1):173-180.

- Dembélé A, Coulibaly A, Traoré SK, Mamadou K, Silue N, Abba T (2008). Détermination du niveau de contamination de l'ochratoxine A (OTA) dans les cerises de cafés verts à l'exportation. *Int. J. Biol. Chem. Sci.* 2(1):33-41.
- Dongo L, Bandyopadhyay R, Kumar M, Ojiambo PS (2008). Occurrence of ochratoxin A in Nigerian ready for sale cocoa beans. *Agric. J.* 3:4-9.
- Durand N, El Sheikh AF, Suarez-Quiros ML, Gonzales-Rios O, Nganou DN, Fontana-Tachon A, Montet D (2013). Application of PCR-DGGE to the study of dynamics and biodiversity of yeasts and potentially ota producing fungi during coffee processing. *Food Control* 34(2):466-471.
- European Communities (EC) 2006. Règlement (CE) No 401/2006 de la Commission du 23 Février 2006 Portant Fixation des Modes de prélèvement d'échantillons et des méthodes d'Analyse pour le contrôle Officiel des teneurs en mycotoxines des denrées alimentaires. *Journal Officiel de l'Union Européenne*: L70/12.
- Gilmour M, Lindblom M (2008). Management of ochratoxin A in the cocoa supply chain: a summary of work by the CAOBisCo/ECA/FCC Working Group on ochratoxine A. In *Mycotoxins: Detection Methods, Management, Public Health and Agricultural Trade*, edited by R. Leslie, R. Bandyopadhyay, and A. Visconti, 231-243. Wallingford: CABI.
- Gopinandhan TN, Kannan GS, Panneerselvam P, Velmourougane K, Raghuramulu Y, Jayarama J (2008). Survey on ochratoxin A in indian green coffee destined for export. *Food Addit. Contam. Part B, Surveillance* 1(1):51-57.
- Higher Public Health Council of France (HPHCF) (1999). Les mycotoxines dans l'alimentation : évaluation et gestion du risque, eds TEC & DOC.
- National Agronomic Research Institute (NARI) (2004). Etude de l'alimentation Totale Française. Mycotoxines, Minéraux et Éléments Traces. Rapport INRA/DGAL. *Coordinateur Jean-Charles Leblanc*. p. 72.
- International Agency for Research on Cancer (IARC) (1993). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, IARC Working Group. Vol. 56. Lyon, France: WHO.
- International Coffee and Cocoa Organization (2014). Available at : [http://www.ico.org/trade\\_statistics.asp?section=Statistics](http://www.ico.org/trade_statistics.asp?section=Statistics).
- Joint FAO/WHO Expert Committee on Food Additives (JECFA) (2001). Safety Evaluation of Certain Mycotoxins in Food. Fifty-Sixth Meeting of the Joint FAO/WHO Expert Committee on Food Additives WHO Food Additives Series 47 – FAO Food and Nutrition Paper –IPCS-International Programme on Chemical Safety. Geneva, Switzerland: WHO.
- Leong SL, Hien LT, An TV, Trang NT, Hocking AD, Scott ES (2007). Ochratoxin A—producing aspergilli in vietnamese green coffee beans. *Lett. Appl. Microbiol.* 45(3):301-306.
- Leoni LAB, Soares LMV, Oliveira PL (2000). Ochratoxin A in brazilian roasted and instant coffees. *Food Addit. Contam.* 17(10):867-870.
- Levi CP, Trenk HL, Mohr HK (1974). Study of the occurrence of ochratoxin A in green coffee beans. *J. Assoc. Off. Anal. Chem.* 57(4):866-70.
- Nganou DN, Durand N, Tatsadjieu NL, Meile JC, El Sheikh AF, Montet D, Mbufung CM (2012). Determination of coffee origin by using 28S rDNA fingerprinting of fungal communities by PCR-DGGE: Application to the Cameroonian coffee. *Int. J. Biosci.* 2(5):18-30.
- Malir F, Ostry V, Pfohl-Leschowicz A, Toman J, Bazin I, Roubal T (2014). Transfer of ochratoxin A into tea and coffee beverages. *Toxins* 6(12): 3438-3453.
- Manderville R, Pfohl-Leschowicz A (2008). Bioactivation and DNA adduction as a rationale for ochratoxin A carcinogenesis. *World Mycotoxin J.* 1(3):357-367.
- Mantle PG, Faucet-Marquis V, Manderville RA, Sciqualli B, Pfohl-Leschowicz A (2010). Structures of covalent adducts between DNA and ochratoxin A: a new factor in debate about genotoxicity and human risk assessment. *Chem. Res. Toxicol.* 23(1):89-98.
- Miraglia M, Brera C (2002). Assessment of dietary intake of ochratoxin A by the population of EU member states. *Rapport Task Force* 3(2).
- Pfohl-Leschowicz A, Manderville RA (2007). Ochratoxin A: an overview on toxicity and carcinogenicity in animals and humans. *Mol. Nutr. Food Res.* 51(1):61-99.
- Pfohl-Leschowicz A, Manderville RA (2012). An Update on direct genotoxicity as a molecular mechanism of ochratoxin A carcinogenicity. *Chem. Res. Toxicol.* 25(2):252-262.
- Pfohl-Leschowicz A, Petkova-Bocharova T, Chernozemsky IN, Castegnaro M (2002). Balkan endemic nephropathy and the associated urinary tract tumours: review on aetiological causes and the potential role of mycotoxins. *Food Addit. Contam.* 19(3):282-302.
- Pittet A, Tornare D, Huggett A, Viani R (1996). Liquid chromatographic determination of ochratoxin A in pure and adulterated soluble coffee using an immunoaffinity column cleanup procedure. *J. Agric. Food Chem.* 44 (11):3564-3569.
- Prado G, Oliveira MS, Abrantes FM, Santos LG, Veloso T, Barroso RES (2000). Incidência de ochratoxina A em café torrado e moído em café solúvel consumido na cidade de belo horizonte, MG. *Cienciae Tecnologia de Alimenta* 2:192-196.
- Rutqvist L, Björklund NE, Hult K, Hökby E, Carlsson B (1978). Ochratoxin A as the cause of spontaneous nephropathy in fattening pigs. *Appl. Environ. Microbiol.* 36(6):920-925.
- Sangare-Tigori B, Moukha S, Kouadio HJ, Betbeder AM, Dano DS, Creppy EE (2006). Co-occurrence of aflatoxin B1, fumonisin B1, ochratoxin A and zearalenone in cereals and peanuts from Côte d'Ivoire. *Food Addit. Contam.* 23(10):1000-1007.
- Scientific Committee for food (SCF) (1998). European Commission DG XXIV Unit B3. Opinion on ochratoxin A
- The Commission of the European Communities (EC) (2006). Commission Regulation (EC) No1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Off. J. Euro. Union* L364:4-24.
- The Commission of the European Communities (EC) (2010). Commission Regulation (EU) No 105/2010 of 5 February 2010 amending Regulation (EC) No 1881/2006 setting maximum levels for certain contaminants in foodstuffs as regards ochratoxin A. *Off. J. Euro. Union* L035 :7-8.
- Turcotte AM, Scott PM (2011). Ochratoxin A in cocoa and chocolate sampled in Canada. *Food Addit. Contam.* 28(6):762-6
- Van der Stegen GHD, Essens PJM, Van der Lijn J (2001). Effect of roasting conditions on reduction of ochratoxin A in coffee. *J. Agric. Food Chem.* 49(10):4713-4715.
- World Health Organization (WHO) (2003). *GEMS/Food regional diets: regional per capita consumption of raw and semi-processed agricultural commodities*. Report of global environment monitoring system/food contamination monitoring and assessment programme. Geneva, Switzerland: Food Safety Department.