

Full Length Research Paper

Sero-prevalence of human immunodeficiency virus (HIV) 1 and 2 infections in Uyo metropolis, Akwa Ibom State

Akinjogunla, O. J.* and Adegoke, A. A.

Department of Microbiology, University of Uyo, Akwa Ibom, Nigeria.

Accepted 12 February, 2009

The seroprevalence of Human immunodeficiency virus (HIV), the causative agents of Acquired Immune Deficiency Syndromes (AIDS) was investigated from January 2008 to June 2008. Three hundred and sixteen (316) blood samples were screened presumptively for presence of HIV1 and HIV-2 using a third generation ELISA kit called "Determine". Out of the 316 samples screened, IgM antibodies to HIV were found present in 108 (34.2%) patients of which 40(12.7%) were males and 68(21.5%) were females. The prevalence of HIV infection in the study population was 34.2% and the prevalence of the viral infection in relation to sex was 39.2% for males and 31.8% for females. The prevalence of viral infection also varied with the highest prevalence rate recorded in age group 31 – 45 years with 43 (13.6%) while the age group 1 - 15 years had the lowest percentage positivity with 5 (1.58%). The findings of this study emphasize the more urgent need for intervention measures like sex education, health education and preventive education among the general populace.

Key words: AIDS, seropositivity, seroprevalence, HIV, Uyo.

INTRODUCTION

Human immunodeficiency virus (HIV) is the aetiological agent of the Acquired Immunodeficiency Syndromes (Georges and Georges-Courboj, 1990; Talaro and Talaro, 1996; Prescott et al., 1999). HIV is one of the human T-cell lymphotropic retroviruses (human T-cell leukemia virus types I and II), of the family Retroviridae and sub family Lentivirinae (Gallo et al., 1983; Fleming, 1984; Abimiku et al., 1994; Murray et al., 1998). The human immunodeficiency virus (HIV) epidemic continues to be a burden globally and presents serious public health problems in developing countries especially in Nigeria. HIV-II antibodies were present in the samples taken in Africa as early as 1966, and convincing evidence from Europe has linked HIV-1 to HIV-like diseases occurring in the late 1950s. The investigation conducted by Olaleye et al. (1993) showed HIV 1 to be more prevalent in Nigeria. It has also reported that HIV 1 is more pathogenic and is found in Asia, East and Central Africa, Europe and United States, while HIV-2 is mainly limited

to West Africa. The AIDS pandemic claimed an estimated 2.1 million (range 1.9 - 2.4 million) lives in 2007 of which an estimated 330,000 were children under 15 years. Globally, an estimated 33.2 million people lived with HIV in 2007, including 2.5 million children and estimated 2.5 million (range 1.8 – 4.1 million) people were newly infected in 2007, including 420,000 children (Pennap et al., 2006). Sub-Saharan Africa remains by far the worst affected region. In 2007, it contained an estimated 68% of all people living with AIDS and 76% of all AIDS deaths. Unlike other regions, most people living with HIV in sub-Saharan Africa in 2007 (61%) were women. Adult prevalence in 2007 was an estimated 5.0%, and AIDS continued to be the single largest cause of mortality in this region (Weiss, 1993; Holmes et al., 2003; Weiss, 2007). The majority of HIV infections are acquired through unprotected sex, blood transfusion, contaminated hypodermic needles, exchange between mother and baby during pregnancy, childbirth, or breastfeeding, or other exposure to one of the above bodily fluids (Johnson and Laga, 1988; Ngaly and Ryder, 1988; Deschamps et al., 1996; Kuby, 1997). HIV/AIDS is also the leading cause of death in sub-Saharan African and it is the fourth biggest killer worldwide.

*Corresponding author. E-mail: papajyde2000@yahoo.com.
Tel.: +2348064069404.

Table 1. Shows the gender and age-specific prevalence of HIV infection in the study participants.

Age group	No. of patients screened	Sex distribution		No. of positive	No. of negative
		Male	Female		
1-15	28	11	17	05	23
16-30	85	28	57	24	61
31-45	110	39	71	43	67
46-60	59	16	43	18	41
60 above	34	08	26	12	22

Nigeria's epidemic is characterized as one of the most rapidly increasing rates of new HIV/AIDS cases.

Several factors have contributed to the rapid spread of HIV in Nigeria and these include sexual networking practices such as polygamy, a high prevalence of untreated sexually transmitted infections, low condom use, poverty, low literacy, poor health status, low status of women (Pennap et al., 2006). The Joint United Nations programme on HIV/AIDS estimated that 3.5 millions Nigerian adults and children were living with HIV/AIDS as at the end of 2001. HIV prevalence in adults increased from 1.8% in 1991 to 5.8% in 2001. There are scanty data on the prevalence of HIV/AIDS at the Local Government Areas of Nigeria. This study was therefore aimed at investigating the prevalence of HIV/AIDS in Uyo local Government Area, Akwa Ibom.

MATERIALS AND METHODS

Study area and population

This study was designed to cover individuals from Uyo metropolis. Uyo is capital city of Akwa Ibom State in South-South (S/S) part of Nigeria. Demographic information about each person screened was obtained. Such information included sex, age, occupation and address. All information about persons screened was kept confidential.

Samples collection

Five milliliter of peripheral blood samples were collected by venous puncture using sterile syringes and needles following informed consent from January to June, 2008. Using this method, the arm of each individual was first tied with a tourniquet and the position of the vein disinfected with cotton soaked with methylated spirit. A separate needle and syringe was used for each sample collection. The blood samples were transferred into sterile universal bottles. Serum was extracted by centrifugation of blood specimens at 3000 rpm for 10 min and stored frozen at -20°C until analyzed in the laboratory.

Detection of virus

For the detection of the presence of HIV-1 and/or HIV-2 antibodies in the blood samples collected, a World Health Organization (WHO) approved kit called 'Determine' which is an ELISA based kit was used. The kit was designed primarily to test for the presence of HIV-1 and/or HIV-2 antibodies in the blood. This ELISA based kit is both sensitive and specific (99 - 100%) and was used according to the

manufacturer's instructions. The kit is made up of strips impregnated with selenium colloid HIV antigen conjugate at one end and two reaction windows labelled as patient and control windows respectively at the other end. The kits were stored at 4°C

Interpretation of test results

When two red bars appeared in both control and patient windows of the strip, the test was interpreted as positive. However, the appearance of only one red bar in the control window of the strip and the absence of the red bar in the patient window was interpreted as a negative test.

Statistical analysis

The prevalence of HIV/AIDS was determined from the proportion of positive individuals to the total number of individual under consideration, and it is expressed in percentage.



RESULTS AND DISCUSSION

Out of the total 316 samples screened, IgM antibodies to HIV were found present in 108 (34.2%) patients of which 40(12.6%) were males and 68(21.5%) were females. (Table 1).The prevalence of HIV infection in the study population of Uyo was 34.2%. There was a significant difference in the sex of the individuals with the viral infection and the prevalence of the viral infection in relation to sex was 39.2% for males and 31.8% for females respectively (Table 2). The prevalence of viral infection also varied with age, the highest prevalence rate was recorded in age group 31 – 45 years with 43(13.6%) out of 316 subjects being positive. The age group 1 - 15 years had the lowest percentage positivity (Table 2). This study revealed that highest incidence of HIV infection in males and females was found in the age group of 31 - 45 years (Table 1). The trend in the prevalence of HIV infection was high in January and very low in April (Figures 1 and 2).

The sero-prevalence of HIV infection from January 2008 to June 2008 was 34.17%, which is an indicative of a higher prevalence among different groups of populations. The prevalence rate recorded in this study is in conformity with the research conducted by Pennap et al. (2006). The results of this study assumed significance and is also an indicative of an emerging epidemic in Uyo. As the World enters the third decade of the HIV/AIDS epi-

Table 2. The incidence of HIV with sex.

Sex	Total No. (%)	No. Positive (%)	No. Negative (%)
Male	102 (32.3)	40 (39.2)	62 (60.8)
Female	214 (67.7)	68 (31.8)	146 (68.2)
Total	316 (100.0)	108 (34.2)	208 (65.8)

**Figure 1.** Trends in HIV prevalence among male screened.**Figure 2.** Trend in HIV prevalence among female patients screened.

demic, the evidence of its impact is undeniable, robbing countries of both human and natural resources. Nigeria, like other countries of the world is also affected, with an estimated average infection of about 6% out of a population of 150 million. In this study the HIV infection was found to be significantly higher, that is 39.2% in male than in female, that is 31.8% and this is also in agreement with the study conducted by Kreiss et al. (1993). This high percentage recorded among the male could be attributed to more risky sex behavior practiced by males and females due to prevailing socio-cultural-economic scenario. One study done in Malawi showed the highest prevalence was in a younger age group (25 - 34 years) in 1994 and shifted to an older age group (35 - 44 years) in 1998. In view of the above it was apparent that now more cases were being detected in older age groups. From the

present study, it is very clear that in Uyo the HIV infection is taking roots in the older age group population which is sexually active as well as the ones most likely involved in risky nonsexual practices such as ear piercing, tattooing, intravenous drug usage, scarification marks, barbing, manicure/pedicure with shared unsterilized instruments. Other contributing factor to sustained spread of HIV/AIDS was that pre-marital and extra-marital sexual contacts were common in Uyo.

The findings of this study emphasize the more urgent need for intervention measures like sex education, health education and preventive education among the general populace to reduce spread of HIV infection we hereby recommend voluntary screening of everyone in order to establish their HIV status. All women who are at high risk of HIV infection should go for HIV testing. Finally, all levels of government should create employment opportunities an alternative and decent means of living for those driven into commercial sex business due to poverty.

ACKNOWLEDGEMENTS

We are grateful to the support given to us by Dr. (Mrs.) B.C Adebayo Tayo in Department of Microbiology, University of Uyo. Also, we are grateful to our numerous participants who kindly responded to the questionnaires.

REFERENCES

- Abimiku AG, Zwanndo G, Kyari N, Opajobi S, Ibanga S, Guyit A (1994). HIV-1 not HIV-2 is present in Nigeria: Need for consideration in vaccines plans. *Vaccine Res.* 3(2); 101-103.
- Deschamps MM, Pape JW, Hafner A, Johnson WD. (1996). Heterosexual transmission of HIV in Haiti. *Annals Intern. Med.* 125 (4): 324-330.
- Fleming A (1984). HTLV from Africa to Japan, *The Lancet* 1: 279.
- Gallo RC, Sliski A, Wong-Staal F (1983). Origin of Human T-cell leukemia/lymphoma Virus. *The Lancet* 2: 962-963.
- Georges AJ, Georges-courboj MC (1990). Epidemiology of HIV infections in Africa. *Rev. Prat.* 40(23): 2131-2135.
- Holmes CB, Losina E, Walensky RP, Yazdanpanah Y, Freedberg KA. (2003). Review of human immunodeficiency virus type 1-related opportunistic infections in sub-Saharan Africa. *Clin. Infect. Dis.* 36 (5): 656-662.
- Johnson AM, Laga M (1988). Heterosexual transmission of HIV. *AIDS* 2 (suppl. 1): S49-S56.
- Kreiss JK, Koech D, Plummer FA (1993). AIDS virus infection in Nairobi prostitutes: Spread of the epidemic to East Africa. *New Engl. J. Med.* 314: 414- 418.
- Kuby J (1997). *Immunology*, 3rd ed. W.H. Freeman and Company, New York, p. 664.

- Murray PR, Rosenthal KS, Pfaller MA. (1998). Retroviruses. Med. Microbiol. pp. 507-519.
- N'Galy B, Ryder RW (1988). Epidemiology of HIV infection in Africa. J. Acquir. Immune Defic. Syndr. 1 (6): 551-558.
- Olaleye OD, Bernstein L, Kkueozor CC (1993). Prevalence of Human Immunodeficiency virus types 1 and 2 infection in Nigeria. J. Infect. Dis. 16: 710-714.
- Pennap GRI, Makut MD, Gyar SD, Owuna G (2006). Sero-prevalence of HIV/AIDS in Keffi and Environs. Nig. J. Microbiol. 20(3):1114-1146.
- Prescott LM, Harley JP, Klein DA. (1999). Acquired Immune Deficiency Syndrome (AIDS). In: Microbiology 4th edition, pp. 746-753.
- Talaro K, Talaro A (1996). Acquired Immunodeficiency Syndromes Foundation in Microbiology, pp. 787-798.
- Weiss RA (2007). Male circumcision as a preventive measure against HIV and other sexually transmitted diseases. Curr. Opin. Infect. Dis. 20 (1): 66-72.
- Weiss RA. (1993). How does HIV cause AIDS? Sci. J. 260: 1273-1279.