

Full Length Research Paper

Length-weight relationships, relative condition factor and relative weight of three fish species from beach seine fishing grounds in Iranian coastal waters of Caspian Sea

Moradinasab, Gh.^{1*}, Raeisi, H¹, Paighambari, S.Y.¹, Ghorbani, R¹ and Bibak, Z.¹

Department of Fisheries, Faculty of Fisheries and Environment, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran.

Accepted 25 April, 2012

The aim of this study is to record the length-weight relationship, relative condition factor (K_{rel}) and relative weight (W_r) for three fish species in Iranian coastal waters of the Caspian Sea. Fish sampling was carried out in the beach seine fishing grounds in autumn and winter seasons for two years (2007 and 2009). 14104 specimens were measured and weighed. The values of the exponent b in the length-weight relationships (LWRs) were 2.8449 for *Cyprinus carpio*, 2.8844 for *Liza aurata* and 2.9077 for *Rutilus frisii kutum*. Relative condition factor (K_{rel}) values were ranged from 1.017 ± 0.002 to 1.071 ± 0.002 . In addition, Relative weight (W_r) ranged from 0.929 ± 0.002 to 1.740 ± 0.004 .

Key words: Length-weight relationship, relative condition factor, relative weight, Caspian Sea.

INTRODUCTION

The Caspian Sea is the largest lake in the world that Iran is associated with that via the coasts of Guilan, Mazandaran and Golestan provinces (Alizadeh, 2004). This lake has a rich diversity of aquatic species. One of the common methods of fishing in this lake is the beach seine fishing which accounts for about 60% of total fishing in the area (Iranian Fisheries Statistic Yearbook, 2008). *Rutilus frisii kutum* (Kamensky, 1901), *Liza aurata* (Risso, 1810) and *Cyprinus carpio* (Linnaeus, 1758) are the most important fish species in fishing composition and they account for more than 95% of the beach seine fishing.

In terms of economic, employment and income, the three species mentioned before play a significant role in people's lives and livelihoods (Abdolmalaki and Ghaninejad, 2005). *R. frisii kutum* is the most valuable species in this fishing method and it is widely reproduced

artificially in reproduction and breeding farms in Iran, and release into the Caspian Sea every year (Razavi sayad, 1999). This species lives only in the basin southern Caspian Sea and so far, it has not been reported its length-weight relationships in Fish base and neither the other two species (*L. aurata* and *C. carpio*) in Iran (Froese and Pauly, 2012).

In this study, length-weight relationship (LWR), relative condition factor (K_{rel}) and relative weight (W_r) for these three species is discussed.

MATERIALS AND METHODS

Data collection

Fish specimens were collected monthly from 130 fishing grounds beach seine with a net mesh size of 28 to 33 mm STR (48° 52' E, 38° 19' N to 53° 45' E, 36° 25' N, Iran) in autumn and winter seasons for two years (2007 and 2009). There was no sampling in, spring and summer because they are close seasons for beach seine fishing in Iran. Numbers of different sizes were separated of combination fishing randomly and then the fresh specimens measured and weighed using the total length (TL, nearest 0.1 cm)

*Corresponding author. E-mail: Moradinasab88@yahoo.com.
Tel: +98 911 7098 500. Fax: +98 0171 2245886.

Table 1. Descriptive statistics and length-weight parameters for the three fish species in the fishing grounds beach seine Iranian coastal waters of Caspian Sea.

Species	n	Length (cm)				WLR parameters and statistics						r ²
		Mean	S.E	Min	Max	a	SE (a)	95% CL(a)	b	SE (b)	95%CL (b)	
<i>Rutilus frisii kutum</i>	4870	41.95	0.094	20.5	62.5	0.0179	0.022	0.0161-0.0198	2.9077	0.0139	2.8803-2.9350	0.91
<i>Cyprinus carpio</i>	2090	46.08	0.181	22.7	72.4	0.0314	0.029	0.0275-0.0358	2.8449	0.0175	2.8104-2.8793	0.93
<i>Liza aurata</i>	7144	36.55	0.075	21.3	60.5	0.0157	0.019	0.0144-0.0172	2.8844	0.0124	2.8599-2.9088	0.93

n, number of individuals sampled; S.E, standard error; Min, minimum, Max, maximum; a, intercept; b, slope; CL 95%, confidence limits; r², coefficient of determination.

and body weight (BW, nearest 1 g), respectively. Total data were pooled together in each species without sexing.

Data analysis

The length-weight relationships were estimated by using following equation (Froese, 2006):

$$W = a L^b$$

Where W is the body wet weight (g), L is the total length (cm), a is the intercept of the regression and b is the regression coefficient (slope). The parameters a and b of the length-weight relationships were estimated by the least-squares method based on logarithms:

$$\log(W) = \log(a) + b \log(L)$$

A t-test was used for comparison the b values obtained from the linear regressions with isometric values (Sokal and Rohlf, 1987):

$$t_s = \frac{(b - 3)}{S_b}$$

Where t_s is the t-test value, b the slope and S_b the standard error of the slope (b). The comparison between the obtained values of t-test and the respective tabled critical values allowed the determination of the b values statistically significant, and their inclusion in an isometric ($b=3$) or allometric range (negative allometric; $b<3$ or positive allometric; $b>3$).

In addition, for each individual, the relative condition

factor (K_{rel}) and the relative weight (W_r) were calculated by following the equations (Le cren, 1951; Froese, 2006; Wege and Anderson, 1978):

$$t_s = \frac{(b - 3)}{S_b}$$

Where t_s is the t-test value, b the slope and S_b the standard error of the slope (b). The comparison between the obtained values of t-test and the respective tabled critical values allowed the determination of the b values statistically significant, and their inclusion in an isometric ($b=3$) or allometric range (negative allometric; $b<3$ or positive allometric; $b>3$).

In addition, for each individual, the relative condition factor (K_{rel}) and the relative weight (W_r) were calculated by following the equations (Le cren, 1951; Froese, 2006; Wege and Anderson, 1978):

$$K_{rel} = \frac{W}{aL^b}$$

$$W_{rm} = 100 \frac{W}{a_m L^b}$$

$$W_r = \frac{W}{W_s}$$

Where W is the body wet weight (g), L is the total length (cm), a and b are the parameters of length-weight relationships, W_s is a standard weight representing the 75th percentile of observed weights at that length, a_m is geometric mean a and b_m is geometric mean b . Statistics

were performed using the R software version 2.11.0.

RESULTS AND DISCUSSION

A total of 14104 specimens of three fish species were collected from the fishing grounds beach seine Iranian coastal waters of Caspian Sea during the present study. The number of individuals sampled (n), the length and weight ranges, parameters a and b of the length-weight relationships, the standard error of b value and the determination coefficient (r^2) for the three species are given in Table 1.

In this study, relative weight was obtained for *R. frisii kutum* 0.929 ± 0.002 . Several factors influence in the growth of the fish such as hereditary characteristics, food reserves, environmental factors, pollution, etc. Investigations carried out indicate that growth *R. frisii kutum* in the recent years has decreased (Abdolmalaki and Ghaninejad, 2005). This can be mainly attributed to artificial breeding and restocking programs carried out every year by the Iranian Fisheries Organization (Table 2).

Over fishing and the elimination of larger individuals of this species due to the use of inappropriate fishing gears have also contributed to this situation. Considering that there is no selection involved in choosing male and female spawners for artificial breeding programs, the

Table 2. Relative condition factor (K_{rel}) (\pm S.E) and Relative weight (W_r) (\pm S.E) for the three fish species in the fishing grounds beach seine Iranian coastal waters of Caspian Sea.

Species	Relative condition factor (K_{rel})			Relative weight (W_r)		
	Min	Max	Mean (S.E)	Min	Max	Mean (S.E)
<i>Rutilus frisii kutum</i>	0.43	2.57	1.017 \pm 0.002	0.38	2.32	0.929 \pm 0.002
<i>Cyprinus carpio</i>	0.45	2.91	1.029 \pm 0.004	0.69	4.30	1.594 \pm 0.006
<i>Liza aurata</i>	0.38	3.28	1.071 \pm 0.002	0.62	5.25	1.740 \pm 0.004

Min, minimum; Max, maximum; S.E, standard error.

Table 3. Length-weight relationships obtained from other parts of the world for *L. aurata* and *C. carpio*.

Species	Location and references	Type of length	Length (cm)	Sex	a	b
<i>Liza aurata</i>	Portugal; Algarve (Borges et al., 2003)	TL	20.1 - 40.5	Unsexed	0.0078	3.006
	Croatia; Eastern Adriatic, (Dulcic and Glamuzina, 2006)	TL	21.5 - 44.2	Unsexed	0.0181	2.952
	Greece; G. Saronikos, (Stergiou and Moutopoulos, 2001)	SL	15.5 - 21.0	Mixed	0.0078	3.230
<i>Cyprinus carpio</i>	Japan; Shioda Plain, Nagano Prefecture, (Carlander, 1969)	TL	31.5 - 57.0	Unsexed	0.0060	3.210
	Turkey; Lake Iznik, Marmara, (Tarkan et al., 2006)	TL	14.2 - 48.8	Unsexed	0.0250	2.830
	Spain; Araquil River, Navarra, (Miranda et al., 2006)	TL	7.1 - 59.0	Mixed	0.0120	3.070

gene bank of this species is gradually shifting.

Thus providing the required conditions for natural spawning in rivers where this species migrates to and or the semi-natural breeding of *R. frisii kutum* in earthen ponds where the selection of breeders is allowed even if only of a small size is stressed more than before (Khanipour and Valipour, 2009).

A result of present study isn't in agreement with reports of Stergiou and Moutopoulos (2001) and Carlander (1969) and is in agreement with reports of Tarkan et al (2006) (Table 3).

In studies of population dynamics high condition factor values shows of favorable environmental conditions (such as: habitat and prey availability) and low values indicate less favorable environmental conditions (Blackwell et al., 2000). Relative condition factor (K_{rel}) is commonly factor for indicate the condition of fish species. In our study, *Liza aurata* (1.071 \pm 0.002) had best performance, while K_{rel} value in *Rutilus frisii kutum* (1.017 \pm 0.002) was lowest across caught species (Table 2).

Table 3 indicates a and b parameters of weight-length relationships of selected species obtained from other parts of the world. Our results mostly agreed with the sturgeon species studies given in Table 3. The difference of a and b can be affected area, sex, season, degree of stomach fullness, gonad maturity, health, habitat, nutrition (Tesch, 1971).

ACKNOWLEDGEMENTS

We thank Iranian Fisheries Organization and Beach

seine fishing cooperative societies for cooperation.

REFERENCES

- Abdolmalaki Sh, Ghaninejad D (2005). Report on stock assessment and composition of the commercial bony fishes on the southern Caspian Sea. Fisheries Research Institute. Bandar Anzali, p. 132.
- Alizadeh H (2004). Introduction to the Features of the Caspian Sea. Norbakhsh publ., p. 120.
- Blackwell BG, Brown ML, Willis DW (2000). Relative Weight (W_r) Status and Current Use in Fisheries Assessment and Management. Rev. Fish. Sci., 8: 1-44.
- Borges TC, Olim S, Erzini K (2003). Weight-length relationship for fish species discarded in commercial fisheries of the Algarve (southern Portugal). J. Appl. Ichthyol., 19(6): 394-396.
- Carlander KD (1969). Handbook of freshwater fishery biology. Volume 1. The Iowa State University Press. Ames. Iowa. DOI <http://dx.doi.org/>.
- Dulcic J, Glamuzina B (2006). Length-weight relationships for selected fish species from three eastern Adriatic estuarine systems (Croatia). J. Appl. Ichthyol., 22: 254-256.
- Froese R (2006). Cube law, condition factor and Length-Weight relationships: history, meta-analysis and recommendations. J. Appl. Ichthyol., 22: 241-253.
- Froese R, Pauly D (2012). Fish base. World Wide Web Electronic Publication. Available at <http://www.fishbase.org>. (accessed on 16 February 2012).
- Khanipour AA, Valipour A (2009). Kutum jewel of the Caspian Sea. Iranian Fisheries Research Organization. Tehran. Iran, p. 97.
- Iranian Fisheries Statistic Yearbook (2008). Statistic catch fishes the basin southern Caspian Sea. Budget and code office- Statistic group and fishing developing study. Tehran Iran, p. 56.
- Le cren ED (1951). The length-weight relationships and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). Anim. Ecol., 20: 201-219.
- Miranda E, Oscoz J, Leunda PM, Escala MC (2006). Weight-length relationships of cyprinid fishes of the Iberian Peninsula. J. Appl. Ichthyol., 22: 297-298.
- Razavi sayad B (1999). Introduction to the ecology of the Caspian Sea.

- Iranian Fisheries Research Organization (IFRO). p. 90.
- Sokal RR, Rolf FJ (1987). Introduction to Biostatistics. 2nd Edition. Freeman. New, York, p. 363.
- Stergiou KI, Moutopoulos DK (2001). A review of length-weight relationships of fishes from Greek marine waters. Naga ICLARM Q. 24(1 and 2): 23-39.
- Tarkan AS, Gaygusuz O, Acipinar H, Gürsoy C, Ozulug M (2006). Length-weight relationships of fishes from the Marmara region (NW-Turkey). J. Appl. Ichthyol., 22: 271-273.
- Tesch FW (1971). Age and growth. In: Ricker, W. E. ed., Methods for Assessment of Fish Production in Freshwaters. Blackwell Scientific Publications. Oxford, pp. 98-100.
- Wege GJ, Anderson RO (1978). Relative weight (W_r): A new index of condition of largemouth bass. In: New approaches to management of small impoundments. G. Novinger J. Dillard (Eds). Am. Fish. Soc. Spec. Publ.. Bethesda. MD, 5: 79-91.